

## RESEARCH PAPER

# Effect of plant neutrophil elastase inhibitor on leucocyte migration, adhesion and cytokine release in inflammatory conditions

C Oliveira<sup>1</sup>, RA Navarro-Xavier<sup>2</sup>, EA Anjos-Vallota<sup>3</sup>, JO Martins<sup>4</sup>, VLF Silveira<sup>5</sup>, LRC Gonçalves<sup>6</sup>, MS Araújo<sup>1</sup>, G Motta<sup>1</sup>, P Sannomiya<sup>7</sup> and MLV Oliva<sup>1</sup>

<sup>1</sup>Department of Biochemistry, Universidade Federal de São Paulo, Rua Três de Maio, São Paulo, Brazil, <sup>2</sup>Laboratory of Immunopharmacology and <sup>3</sup>Laboratory of Inflammation, Instituto Oswaldo Cruz, Fiocruz, Manguinhos, Rio de Janeiro, Brazil, <sup>4</sup>Department of Immunology, Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, Brazil, <sup>5</sup>Department of Biological Sciences, Universidade Federal de São Paulo, Diadema, São Paulo, Brazil, <sup>6</sup>Laboratory of Pathophysiology, Instituto Butantan, São Paulo, Brazil and <sup>7</sup>Instituto do Coração (InCor), LIM-11, Universidade de São Paulo, São Paulo, Brazil

## BACKGROUND AND PURPOSE

The serine and cysteine peptidase inhibitor, BbCI, isolated from *Bauhinia bauhinoides* seeds, is similar to the classical plant Kunitz inhibitor, STI, but lacks disulphide bridges and methionine residues. BbCI blocks activity of the serine peptidases, elastase ( $K_{iapp}$  5.3 nM) and cathepsin G ( $K_{iapp}$  160.0 nM), and the cysteine peptidase cathepsin L ( $K_{iapp}$  0.2 nM). These three peptidases play important roles in the inflammatory process.

## EXPERIMENTAL APPROACH

We measured the effects of BbCI on paw oedema and on leucocyte accumulation in pleurisy, both induced by carrageenan. Leucocyte–endothelial cell interactions in scrotal microvasculature in Wistar rats were investigated using intravital microscopy. Cytokine levels in pleural exudate and serum were measured by ELISA.

## KEY RESULTS

Pretreatment of the animals with BbCI ( $2.5 \text{ mg} \cdot \text{kg}^{-1}$ ), 30 min before carrageenan-induced inflammation, effectively reduced paw oedema and bradykinin release, neutrophil migration into the pleural cavity. The number of rolling, adhered and migrated leucocytes at the spermatic fascia microcirculation following carrageenan injection into the scrotum were reduced by BbCI pretreatment. Furthermore, levels of the rat chemokine cytokine-induced neutrophil chemo-attractant-1 were significantly reduced in both pleural exudates and serum from animals pretreated with BbCI. Levels of interleukin-1 $\beta$  or tumour necrosis factor- $\alpha$ , however, did not change.

## CONCLUSIONS AND IMPLICATIONS

Taken together, our data suggest that the anti-inflammatory properties of BbCI may be useful in investigations of other pathological processes in which human neutrophil elastase, cathepsin G and cathepsin L play important roles.

## Abbreviations

BbCI, *Bauhinia bauhinoides* serine and cysteine peptidase inhibitor; CINC-1, cytokine-induced neutrophil chemo-attractant-1; IL-1 $\beta$ , interleukin-1beta;  $K_{iapp}$ , apparent inhibitory constant; TNF- $\alpha$ , tumour necrosis factor-alpha

## Correspondence

Maria Luiza V. Oliva,  
Universidade Federal de São  
Paulo-Escola Paulista de  
Medicina, Departamento de  
Bioquímica, Rua Três de Maio  
100, 04044-020, São Paulo, SP,  
Brazil. E-mail:  
olivaml.bioq@epm.br

## Keywords

*Bauhinia*; bradykinin; cytokines;  
elastase; inflammation; intravital  
microscopy; Kunitz inhibitor;  
paw oedema; pleurisy

## Received

23 April 2010

## Accepted

18 May 2010

## Introduction

A variety of pathological conditions is associated with the release of human neutrophil elastase, a serine peptidase which plays an important role in inflammatory processes. This peptidase is a very damaging enzyme because it degrades insoluble elastin into soluble peptides (Owen and Campbell, 1999; Pham, 2006). Cathepsin G, another serine peptidase, which also has broad substrate specificity, acts in cooperation with human neutrophil elastase. Both peptidases are released during inflammation and can be extremely destructive if their activities are not controlled (Owen and Campbell, 1995). The major endogenous inhibitors of these peptidases are the  $\alpha_1$ -peptidase inhibitor ( $\alpha_1$ -PI), secretory leucoprotease inhibitor (SLPI) and  $\alpha_2$ -macroglobulin, and when the enzyme/inhibitor balance is in favour of these neutrophil enzymes, their deleterious actions become more obvious (Sallenave, 2000; Moraes *et al.*, 2003).

One sign of acute inflammation is neutrophil [polymorphonuclear (PMN) cells] migration across endothelium and towards the affected tissue. However, the mechanism used by PMN cells to cross the endothelium and enter into the interstitium of inflamed tissue is not entirely clear despite the number of investigations on this subject (Skrzydowska *et al.*, 2005).

One of the major events in cell migration is leucocyte chemotaxis, which involves two independent, but interrelated processes, such as mobility and directionality that are regulated by extracellular chemo-attractant stimuli (Wu *et al.*, 2005). Many different classes of compounds can act as leucocyte chemo-attractants, including lipids, antimicrobial peptides, chemokines and cytokines (Zabel *et al.*, 2006). In mammalian systems, the complement fragment C5a and the superfamily of small secreted proteins and pro-inflammatory cytokines regulate leucocyte recruitment, infiltration and trafficking, as well as their function (Zlotnik and Yoshie, 2000; Pease and Williams, 2006).

Numerous events are involved in controlling chemo-attractant release. For instance, C5a is generated by serine peptidase activities to which plasmin contributes importantly (Moraes *et al.*, 2003; Kohl *et al.*, 2006), interleukin (IL)1- $\beta$  is generated by both cathepsin G and elastase activities (Kasama *et al.*, 2005) and IL-8 [the rat orthologue is cytokine-induced neutrophil chemo-attractant-1 (CINC-1)] is generated by IL1- $\beta$  (Campbell *et al.*, 2003).

Inflammation plays an important role in host defence, and this uncontrolled process results in many human diseases, such as ischaemia-

reperfusion injury, allergic reactions and tumorigenesis among others (Coussens and Werb, 2002; Nathan, 2002). The lack of specific therapeutic agents has impaired effective treatment for these inflammatory conditions. Thus, a new approach for a better understanding of the inflammatory processes may provide novel therapeutic targets or treatment strategies.

The *Bauhinia* genus belongs to the subfamily Caesalpinoideae and comprises more than 600 species widely distributed in the tropical and subtropical forests (Vaz and Tozzi, 2005). Many proteins have been isolated from their seeds, and particularly in *bauhinioides* species there are two inhibitors named *Bauhinia bauhinioides* kallikrein inhibitor (BbKI) and *B. bauhinioides* cruzipain inhibitor (BbCI) showing distinct structural characteristics and biochemical properties (Oliva *et al.*, 1999a,b; Oliva *et al.*, 2001; de Oliveira *et al.*, 2001; Oliva and Sampaio, 2008). BbCI inhibits the activity of different serine peptidases, such as cathepsin G ( $K_{iapp}$  160.0 nM); human neutrophil elastase ( $K_{iapp}$  5.3 nM); porcine pancreatic elastase ( $K_{iapp}$  2.5 nM); and cysteine peptidases, such as cruzipain ( $K_{iapp}$  1.2 nM), cruzain ( $K_{iapp}$  0.3 nM) and cathepsin L ( $K_{iapp}$  0.2 nM) (de Oliveira *et al.*, 2001). The unexpected and uncommon property of inhibiting two classes of enzymes was confirmed by studies with the recombinant protein (Araujo *et al.*, 2005). Moreover, BbCI lacks cysteine and methionine residues conferring resistance to protein oxidation in contrast to  $\alpha_1$ -PI, SLPI and  $\alpha_2$ -macroglobulin, which are oxidized at methionine in the P<sub>1</sub> position of the reactive site, impairing the efficacy of peptidase inhibition (Travis and Salvesen, 1983; Jung *et al.*, 1995).

Therefore, in the present series of experiments, we have evaluated the effects of BbCI on elastase, cathepsin G and cathepsin L activities in acute inflammatory models *in vivo*, and have assessed some of the chemical mediators involved.

## Methods

### Animals

All animal care and experimental procedures were in accordance with the ethical principles in animal research, as adopted by the Brazilian College of Animal Experimentation and approved by the Federal University of São Paulo Ethical Committee for Animals. In this study, adult male Wistar albino rats, weighing 200–250 g, housed at  $22 \pm 1^\circ\text{C}$ , under 12 h light–dark cycle time were used. The animals had a standard pellet diet and tap water *ad libitum*. The animals were acclimatized to the laboratory for

at least 1 h before testing. They were used only once throughout the experiments.

### *Inhibitor purification and inhibitory activity*

BbCI was obtained from *B. bauhinoides* seeds using standard protein purification techniques previously described (de Oliveira *et al.*, 2001). N-terminal sequence analysis indicated a single protein structure corresponding to SVLDTKGEPVSNAADAYLPVPVSJGEFLALAKVG, confirming the homogeneity of the inhibitor isolated. The activity of BbCI was confirmed on human neutrophil elastase, cathepsin G and on cathepsin L.

### *Carrageenan-induced paw oedema and coaxial perfusion*

The rats were anaesthetized with sodium pentobarbital (40 mg·kg<sup>-1</sup>, i.p.). Anaesthetic supplementation (20 mg·kg<sup>-1</sup>) was provided as required to maintain light anaesthesia during the experiment. Carrageenan (100 µg) was injected in the right subplantar region. Paw volumes were determined up to 4 h by plethysmography (H. Basile, Milan, Italy). Each measurement was repeated three times, and the mean value was calculated (Silveira *et al.*, 1995; Oliva *et al.*, 2000). The results were expressed as increases in rat paw volume (%) from 0 to 4 h following carrageenan injection. The inflamed animals were pretreated with BbCI 2.5 mg·kg<sup>-1</sup> or sterile saline injected intravenously (tail vein), 30 min before oedema was induced by carrageenan injection.

For the determination of the local levels of bradykinin (BK) in inflamed (injected with carrageenan) or non-inflamed groups, rats were submitted to a paw coaxial perfusion as described by Wohlers *et al.* (2005). Briefly, before the carrageenan injection, the paw was cannulated and perfused at 53 µL·min<sup>-1</sup> with a perfusion solution (NaCl 158 mM, KCl 5.65 mM, CaCl<sub>2</sub> 0.54 mM, MgCl<sub>2</sub> 2.90 mM, NaHCO<sub>3</sub> 178 mM, glucose 1.0 g·L<sup>-1</sup>) kept in a bath at 36.5°C. Paw perfusate samples were collected during 0–1.5 and 1.5–2.5 h after carrageenan injection, and placed into ice-cold tubes containing 80% ethanol. The samples were centrifuged, and supernatant was evaporated in a flow of nitrogen and stored at –70°C. BK levels were determined by radioimmunoassay (RIA) as described below.

### *RIA*

Kinins were extracted from perfusate by ethanol precipitation (1:4 v/v) and left for 10 min at –70°C. The samples were then freeze-dried and dissolved in 200 µL of buffer (phosphate 0.01 M, NaCl 0.14 M containing 0.1% egg albumin, EDTA 30 mM, orthophenanthroline 3 mM, 0.1% NaN<sub>3</sub>), pH 7.0. Aliquots (50 µL) were incubated with 100 µL anti-BK

antibody (1:80,000) and 100 µL of [<sup>125</sup>I-Tyr]-BK for 20 h at 4°C. Then, 400 µL of 0.1% bovine γ-globulin in buffer B (phosphate 0.01 M, 0.14 M NaCl containing 0.1% NaN<sub>3</sub>), pH 7.0, and 800 µL of 25% polyethyleneglycol 6000 (w/v) solution were added to the samples, which were incubated for 10 min at 4°C. After incubation, the samples were centrifuged at 2,000×g for 20 min at 4°C, and the supernatants were removed. The [<sup>125</sup>I] in the pellets was counted and the BK released was calculated (Shimamoto *et al.*, 1982; Gozzo *et al.*, 2002).

### *Carrageenan-induced pleurisy*

This experiment included the inflamed and non-inflamed groups. Both groups were submitted to anaesthetic procedure. At 30 min before injection of 100 µg of carrageenan into the pleural space, the inflamed group was pretreated with BbCI (2.5 mg·kg<sup>-1</sup>) or saline by the tail vein. The non-inflamed rats received the same volume of saline in the pleural space and BbCI (2.5 mg·kg<sup>-1</sup>) or saline injected via the tail vein.

Carrageenan was injected into the pleural space between the third and fourth ribs on the right side of the mediastinum according to the method of Vinegar *et al.* (1982). Four hours later, the animals were killed and carefully bled from the carotid arteries. The exudate was removed by aspiration from the pleural cavity and collected in 5.0 mL of PBS buffer (NaCl 137 mM, KCl 2.7 mM, Na<sub>2</sub>HPO<sub>4</sub> 8.1 mM, KH<sub>2</sub>PO<sub>4</sub> 1.5 mM, pH 7.2), according to the procedure previously described by Harada *et al.* (1996). Any exudates contaminated with blood were discarded.

Total leucocytes in blood were determined using a Neubauer chamber after dilution with Turk's solution (1:20 v/v) before and 4 h after saline or carrageenan injection. Leucocytes in the exudate at the pleural cavity were determined after 4 h of pleurisy. Differential leucocyte counts were carried out on stained slides under oil immersion microscopy. A total of 100 cells were counted and classified as neutrophils, eosinophils, monocytes and lymphocytes based on morphological criteria.

### *Intravital microscopy of the microcirculation and haemodynamic parameters*

The group of inflamed rats were pretreated with BbCI 2.5 mg·kg<sup>-1</sup> or saline injected via the tail vein, 30 min before the injection of 100 µg of carrageenan into the scrotal chamber, under light ether anaesthesia. Two non-inflamed groups, pretreated with BbCI 2.5 mg·kg<sup>-1</sup> or saline, were also submitted to anaesthesia. Rolling, adhesion and migration of leucocytes were analysed 2 h after the local injection of carrageenan.

The animals were anaesthetized with sodium pentobarbital ( $40 \text{ mg}\cdot\text{kg}^{-1}$ , i.p.), and the internal spermatic fascia of the wall of the scrotal chamber was exteriorized for microscopic examination *in situ* (Fortes *et al.*, 1991). This procedure was performed through a longitudinal incision of the skin and dartos muscle in the midline over the ventral aspect of the scrotum, then opening the cremaster muscle to expose the internal fascia. The procedure does not require extensive surgical manipulation for the observation of the vascular network, and provides a valuable means for trans-illuminating the tissue for quantitative studies of the microcirculation. The animals were kept on a special board thermostatically controlled at  $37^\circ\text{C}$ , which included a transparent platform on which the tissue to be trans-illuminated was placed. The preparation was kept moist and warmed by irrigating the tissue with warmed ( $37^\circ\text{C}$ ) Ringer–Locke's solution, NaCl 154 mM, KCl 5.6 mM,  $\text{CaCl}_2$  2 mM,  $\text{NaHCO}_3$  6 mM, glucose 5 mM, pH 7.2 containing 1% gelatin. The rate of the solution outflow onto the exposed tissue was controlled to keep the preparation in continuous contact with a film of liquid. A 500-line television camera was incorporated with a triocular Zeiss microscope (Carl Zeiss Co, München-Hallbergmoos, Germany) to facilitate the observation of the enlarged image ( $\times 3400$ ) on the video screen. Images were recorded on a video recorder using an  $\times 40$  long-distance objective with a 0.65 numerical aperture. An image-splitting micrometer was adjusted to the phototube of the microscope (Baez, 1973). The image splitter sheared the optical image into two separate images and displaced one with respect to the other. By rotating the image splitter in the phototube, the shearing was maintained in a direction at right angles to the axis of the vessel. The displacement of one image with the other allowed measurement of the vessel diameter. Vessels selected for study were third-order venules, defined according to their branch order location within the microvascular network (Gore and Bohlen, 1977; Rhodin, 1986). These vessels corresponded to post-capillary venules and their diameters ranged from 12 to  $16 \mu\text{m}$ .

Centerline red blood cell velocity was measured using an optical Doppler velocimeter (Microcirculation Research Institute, Texas A&M University, College Station, TX, USA) previously calibrated against a rotating glass disk coated with red blood cells. Venular blood flow velocity was calculated using the formula ( $V_{\text{mean}} = \text{centerline velocity}/1.6$ ). Venular wall shear rate ( $\gamma$ ) was calculated using the Newtonian definition:  $\gamma = 8 (V_{\text{mean}}/D_v)$ , where  $D_v$  = vessel diameter) (Davis, 1987; Panes *et al.*, 1996).

Leucocytes rolling and adhering to the endothelium (stickers) and migrating into the adjacent

perivascular tissue were evaluated using recorded images (Farsky *et al.*, 1995). Two observers independently scored counts. The number of rolling leucocytes was presented as the mean number of cells passing at a designated line perpendicular to the venular axis in 10 min. A leucocyte was considered to be adherent to the venular endothelium if it remained stationary for more than 30 s (Farsky *et al.*, 1995). Adherent cells were counted during a 10 min period in a  $100 \mu\text{m}$  segment of the vessel. A given section of the vascular bed was tested only once for determination of the number of stickers. Three to five determinations were performed on a single animal. These were averaged for each animal. The number of leucocytes accumulating in the connective tissue adjacent to a post-capillary venule was determined in a standard area of  $1000 \mu\text{m}^2$ . This area was defined on the video screen,  $10 \mu\text{m}$  in tissue corresponding to 3.4 cm on the screen (magnification  $\times 3400$ ). Three to five different fields were evaluated on a single animal to avoid variability based upon sampling. Data were then averaged for each animal.

#### *Enzyme immunoassay for cytokines and chemokine*

The concentrations of tumour necrosis factor (TNF)- $\alpha$ , IL-1 $\beta$  and CINC-1 were determined in pleural exudates and serum samples by ELISA using commercially available kits, according to the manufacturer's instructions (R&D Systems Inc., Minneapolis, MN, USA). The sensitivity of the assay was  $15 \text{ pg}\cdot\text{mL}^{-1}$ .

#### *Data analysis*

Data were expressed as mean  $\pm$  SEM and analysed by a computerized package for statistical analysis. A statistically significant difference was determined by the Tukey–Kramer multiple comparison test. The Kruskal–Wallis test followed by Dunn's multiple comparisons test was used when data were not normally distributed. The significance was set at  $P < 0.05$ .

#### *Materials*


ELISA kits for rat TNF- $\alpha$ , IL-1 $\beta$  and CINC-1 were obtained from R&D Systems Inc. All other chemicals used were of reagent grade.  $\lambda$ -Carrageenan type IV and urethane were purchased from Sigma Chemical Company (St Louis, MO, USA). The other chemicals were of analytical grade.

## **Results**

#### *Effect of BbCI on paw oedema and kinin release*

Figure 1 shows the development of paw oedema induced by carrageenan, which was monitored





**Figure 1**


Effect of BbCI on carrageenan-induced paw oedema. Rats were pretreated intravenously with saline ( $n = 12$ ) or BbCI ( $2.5 \text{ mg} \cdot \text{kg}^{-1}$ , i.v.  $n = 12$ ) 30 min before the local injection of carrageenan ( $100 \mu\text{g} \cdot 0.1 \text{ mL}^{-1}$ ). Oedema was expressed as increases in rat paw volume (%) from 0 to 4 h following carrageenan injection. Data are expressed as mean  $\pm$  SEM (\*)  $P < 0.01$ .

during 4 h after the subplantar injection of  $100 \mu\text{g}$  carrageenan in rats. Relative to the saline-inflamed group, which showed a time-dependent increase in paw oedema volume, in the inflamed rats pretreated with BbCI ( $2.5 \text{ mg} \cdot \text{kg}^{-1}$ ), there was a significant reduction in paw oedema at the second (24%), third (44%) and fourth (40%) hours after carrageenan injection.

Compared to non-inflamed rats, the local injection of carrageenan induced significant increases in the levels of BK obtained from the perfusion fluid of the rat paw (Figure 2). Values were higher at 0–1.5 h than at 1.5–2.5 h intervals after carrageenan injection. In contrast, BbCI-treated inflamed rats exhibited a significant reduction in the levels of BK at 0–1.5 h interval. Values attained at 1.5–2.5 h were similar to those observed in saline-treated inflamed group.

#### Effect of BbCI on leucocyte–endothelial interactions and haemodynamic parameters

Rats pretreated with BbCI or saline were submitted to a local injection of carrageenan (inflamed group) or saline (non-inflamed group), and leucocyte–endothelial interactions at the microcirculation of the spermatic fascia were evaluated 2 h thereafter. Results, illustrated in Figure 3, showed that saline-treated inflamed rats exhibited significant increases in the number of rolling, adherent and migrated leucocytes compared to saline-treated non-inflamed rats. In contrast, BbCI-treated inflamed rats presented a significant reduction on leucocyte–endothelial interactions, including rollers, stickers




**Figure 2**

Effect of BbCI on BK release. Rats were pretreated intravenously with saline (S) or with  $2.5 \text{ mg} \cdot \text{kg}^{-1}$  BbCI 30 min before the local injection of saline (non-inflamed) or carrageenan ( $100 \mu\text{g} \cdot 0.1 \text{ mL}^{-1}$ ). BK levels were determined by RIA in paw perfusate samples collected during 0–1.5 and 1.5–2.5 h after the local injection of saline or carrageenan. Data are expressed as mean  $\pm$  SEM for six animals in each group. \* $P < 0.001$  versus saline-treated inflamed rats; # $P < 0.001$  and # $P < 0.01$  versus non-inflamed rats; § $P < 0.001$  versus corresponding value at 1.5 h.

and migrated cells. Values attained matched those observed in saline- or BbCI-treated non-inflamed animals. Representative photomicrographs are shown in Figure 4. There were no differences in the mean arterial pressure, blood flow velocity and venular wall shear rate between saline- and BbCI-treated inflamed rats (Table 1).


#### Effect of BbCI on carrageenan-induced pleurisy

The effect of BbCI on leucocyte recruitment was evaluated by measuring total and differential cell counts in pleural exudate samples. As summarized in Table 2, there were no differences in the number of leucocytes [PMN and mononuclear (MN) cells] in the pleural cavity of saline- or BbCI-treated non-inflamed rats. In contrast, treatment of inflamed rats with BbCI reduced the number of leucocytes harvested from the pleural cavity, compared to saline-treated inflamed rats. The number of PMN leucocytes was significantly reduced to 39% of those observed in saline-treated inflamed rats. There were no differences in the number of MN cells between saline- and BbCI-treated inflamed rats. Results, summarized in Table 3, showed that there were equivalent increases in the number of total blood leucocyte counts in saline- and BbCI-treated inflamed rats during the 4 h pleurisy. The number of



**Figure 3**

Effect of BbCI on leucocyte–endothelial interactions. Number of rolling leucocytes/10 min (A), adherent leucocytes to the endothelium of a 100  $\mu\text{m}$  segment of post-capillary venules (B) and number of migrated leucocytes to a standard area (1000  $\mu\text{m}^2$ ) of the perivascular tissue (C). Rats were pretreated intravenously with saline (S) or BbCI (2.5  $\text{mg}\cdot\text{kg}^{-1}$ ) 30 min before the local injection of carrageenan (100  $\mu\text{g}\cdot 0.1\text{ mL}^{-1}$ ). Leucocyte–endothelial interactions at the spermatic fascia were evaluated 2 h thereafter. Data are expressed as mean  $\pm$  SEM for six animals in each group \* $P < 0.001$  versus other groups.



**Figure 4**

Photomicrographs of a venular segment of the spermatic fascia under intravital microscopy 2 h after the local injection of carrageenan. (100  $\mu\text{g}\cdot 0.1\text{ mL}^{-1}$ ). Rats were pretreated intravenously with saline or BbCI (2.5  $\text{mg}\cdot\text{kg}^{-1}$ ) 30 min before carrageenan injection. Arrows show sites of leucocyte adhesion. Magnification ( $\times 3400$ ).

neutrophils increased twofold in both groups. Monocytes and eosinophils did not change. A slight decrease in the number of lymphocytes was observed in BbCI-treated inflamed rats. In non-inflamed groups, total and differential leucocyte counts in the blood did not change either in saline- or BbCI-treated rats.

#### *Effect of BbCI on IL-1 $\beta$ , TNF- $\alpha$ and CINC-1 levels*

IL-1 $\beta$ , TNF- $\alpha$  and CINC-1 concentrations were measured in pleural exudates and serum samples in order to determine whether the release/production of these inflammatory mediators could be modulated by BbCI. Relative to saline- or BbCI-treated non-inflamed rats, equivalent increases in the levels

**Table 1**

Effect of BbCI on haemodynamic parameters

Treatment	Vessel diameter ( $\mu\text{m}$ )	Mean arterial pressure (mmHg)	Blood flow velocity ( $\text{mm}\cdot\text{s}^{-1}$ )	Venular wall shear rate ( $\text{s}^{-1}$ )
Saline + carrageenan	$19.8 \pm 1.7$	$120 \pm 6$	$2.00 \pm 0.05$	$823 \pm 71$
BbCI + carrageenan	$19.4 \pm 1.3$	$122 \pm 4$	$1.92 \pm 0.09$	$790 \pm 58$

Rats were pretreated with saline or BbCI ( $2.5 \text{ mg}\cdot\text{kg}^{-1}$ , i.v.) 30 min before the local injection of carrageenan ( $100 \mu\text{g}$ ) and evaluated 2 h thereafter. Values are presented as mean  $\pm$  SEM of six animals in each group.

**Table 2**

Effect of BbCI on carrageenan-induced 4 h pleurisy

Leucocytes ( $\times 10^6$ ) Group	Total	PMN	MN
Non-inflamed			
Saline ( $n = 8$ )	$1.65 \pm 0.16$	$0.35 \pm 0.1$	$1.30 \pm 0.1$
BbCI ( $n = 8$ )	$1.43 \pm 0.15$	$0.33 \pm 0.1$	$1.10 \pm 0.12$
Inflamed			
Saline ( $n = 15$ )	$9.11 \pm 0.48$	$7.33 \pm 0.5$	$1.78 \pm 0.15$
BbCI ( $n = 15$ )	$3.55 \pm 0.23^*$	$2.85 \pm 0.2^*$	$0.70 \pm 0.1$

Rats were pretreated with saline or BbCI ( $2.5 \text{ mg}\cdot\text{kg}^{-1}$ , i.v.) 30 min before the intrapleural injection of saline (non-inflamed) or carrageenan ( $100 \mu\text{g}$ , inflamed). Pleural exudates were collected 4 h thereafter. PMN and MN (leucocytes) in exudate samples were counted. Values are presented as mean  $\pm$  SEM.  $^*P < 0.05$  versus corresponding values in saline-treated inflamed rats.

of IL-1 $\beta$  and TNF- $\alpha$  were observed in pleural exudates obtained from saline- or BbCI-treated inflamed rats (Figure 5A,B). In contrast, levels of these cytokines in the serum did not differ between non-inflamed and inflamed rats in both saline- and BbCI-treated groups (Figure 6A,B).

CINC-1 concentrations significantly increased in both pleural exudate and serum samples from the saline-treated inflamed group, as compared to those from the non-inflamed group (saline or BbCI) as illustrated in Figures 5C and 6C respectively. In contrast, CINC-1 levels in pleural exudate and serum were significantly reduced in the inflamed rats, pretreated with BbCI.

## Discussion

Several different proteins isolated from *Bauhinia* seeds have been purified and characterized in terms of their activities on a range of functions, such as haemostasis (Oliva *et al.*, 2003), digestion (Oliva *et al.*, 1999a,b; Sumikawa *et al.*, 2006; Oliva and Sampaio, 2008) and protein–protein interaction (Macedo *et al.*, 2004). Many studies have used crude plant extracts which were neither chemically nor biochemically adequately characterized.

In the present study, we investigated the effect of the sequenced and crystallized proteinase inhibitor, BbCI, on the course of the inflammatory response. BbCI is a potent inhibitor of human neutrophil elastase ( $K_{\text{iapp}}$  5.3 nM), cathepsin G ( $K_{\text{iapp}}$  160.0 nM) and cathepsin L ( $K_{\text{iapp}}$  0.2 nM; de Oliveira *et al.*, 2001). These enzymes degrade components of endothelial cell junctions, allowing PMN migration (Owen and Campbell, 1999; Korkmaz *et al.*, 2008).

Endogenously,  $\alpha_1$ -antipeptidase,  $\alpha_2$ -macroglobulin and SLPI effectively regulate both extracellular neutrophil elastase and cathepsin G activities (Weiss, 1989; Skrzydlewska *et al.*, 2005). The proteolytic–antiproteolytic balance of plasma peptidases and their inhibitors is critically affected by reactive oxygen species (ROS) generated by activated neutrophils, as these inhibitors contain methionine in their structure and are thus susceptible to inactivation by oxidizing compounds, resulting in enhanced proteolysis and destruction of extracellular matrix (Weiss, 1989; Vogt, 1995; Skrzydlewska *et al.*, 2005). The structure of BbCI is unlike these inhibitors, as it lacks methionine and cysteine residues, and this lack suggests a potential resistance to oxidizing conditions and thus a therapeutic value for this inhibitor, which could control the activity of

**Table 3**

Effect of BbCI on blood leucocyte counts

Cells·mm <sup>-3</sup> Group	Total	Lymphocytes	Neutrophils	Monocytes	Eosinophils
Non-inflamed					
Saline (n = 8)					
0 h	12.485 ± 432	9.505 ± 360	2.495 ± 133	326 ± 31	159 ± 26
4 h	13.387 ± 325	10.257 ± 317	2.572 ± 139	370 ± 53	188 ± 25
BbCI (n = 8)					
0 h	12.745 ± 494	9.458 ± 472	2.694 ± 189	362 ± 34	231 ± 19
4 h	13.005 ± 547	9.544 ± 415	2.872 ± 120	390 ± 42	199 ± 29
Inflamed					
Saline (n = 15)					
0 h	12.222 ± 347	9.333 ± 294	2.288 ± 98	449 ± 32	152 ± 18
4 h	15.402 ± 394*	9.687 ± 325	4.989 ± 148*	538 ± 32	188 ± 18
BbCI (n = 15)					
0 h	12.458 ± 281	8.901 ± 235	2.962 ± 129	369 ± 32	226 ± 28
4 h	15.026 ± 307*	7.449 ± 224*	6.846 ± 230*	508 ± 46	223 ± 21

Rats were pretreated with saline or BbCI (2.5 mg·kg<sup>-1</sup>, i.v.) 30 min before the intrapleural injection of saline (non-inflamed) or carrageenan (100 µg, inflamed). Blood samples were collected from the cut tip of the tail of the animals before and 4 h thereafter. \**P* < 0.05 versus corresponding values before pleurisy.

these peptidases in the presence of ROS (de Oliveira *et al.*, 2001; Araujo *et al.*, 2005).

In this work, carrageenan was used to induce inflammation. It is known that the oedematogenic effect of carrageenan is triphasic (i.e. the first phase involves the release of histamine and 5-HT; the second phase is related to kinin release, especially BK; and the third phase is attributed to local production of prostaglandin E<sub>2</sub>) (Di Rosa, 1972; Morris, 2003).

The pro-inflammatory properties of kinins include vasodilation, increased microvascular permeability, release of nitric oxide and a modulating effect on the release of histamine and prostaglandin E<sub>2</sub> (Moreau *et al.*, 2005). The anti-oedematogenic effect of BbCI was seen in the second and third phases of carrageenan-induced paw oedema, which suggests a major influence of BbCI on kinin production. Indeed, a significant reduction in the levels of BK was observed in BbCI-treated inflamed rats, probably due to the inhibitory action of BbCI on kinogenase enzyme activity such as cathepsin L, which can also release BK (Desmazes *et al.*, 2001; Puzer *et al.*, 2004).


Neuhof *et al.* (2003) have shown that the pulmonary oedema in isolated rabbit lungs caused by neutrophil elastase is significantly decreased by 100 µmol·L<sup>-1</sup> BbCI. In the present study, the strong inhibitory effect of BbCI on leucocyte rolling (70%),

adhesion (95%) and migration (90%) in inflamed tissues was demonstrated. This inhibitory effect could be due to haemodynamic changes resulting from BbCI pretreatment. However, there were no differences in blood flow velocity and venular shear rate between saline-treated and BbCI-treated groups, under the inflammatory conditions we have studied here.

Similar results were observed in carrageenan-induced pleurisy. A significant decrease (60%) of neutrophil migration to the pleural cavity was demonstrated in BbCI-treated rats. This inhibitory effect apparently was not related to a defect on neutrophil mobilization from bone marrow to the circulation because equivalent increases in the number of blood neutrophils were observed in both saline- and BbCI-treated rats during the 4 h of pleurisy.


Although the levels of TNF and IL-1β in pleural exudate were not modified by BbCI, a significant decrease in CINC-1 levels occurred and may explain the effect of BbCI on neutrophil migration. This chemokine can be activated by elastase and cathepsin G (Mezyk-Kopiec *et al.*, 2005; Soehnlein *et al.*, 2009). Elastase enhances the expression of adhesion molecules such as the intercellular adhesion molecule-1 and chemokine receptors (Zimmermann *et al.*, 2005; Ishihara *et al.*, 2006). Thus, BbCI by inhibiting elastase and cathepsin G might reduce CINC-1 release and neutrophil migration.





**Figure 5**

Effect of BbCI on cytokine in the pleural exudate. Rats were pretreated intravenously with saline (S) or BbCI (2.5 mg·kg<sup>-1</sup>) 30 min before the intrapleural injection of saline (non-inflamed) or carrageenan (100 µg·0.1 mL<sup>-1</sup>, inflamed). Concentrations of IL-1β (A), TNF-α (B) and CINC-1 (C) were determined by ELISA in pleural exudate 4 h after the induction of pleurisy. Data are expressed as mean ± SEM for six animals in each group. There was a significant decrease in CINC-1 levels in the exudate from BbCI-treated inflamed rats. \**P* < 0.05 versus corresponding values in saline-treated inflamed rats.



**Figure 6**

Effect of BbCI on serum cytokine levels. Rats were pretreated intravenously with saline (S) or BbCI (2.5 mg·kg<sup>-1</sup>) 30 min before the intrapleural injection of saline (non-inflamed) or carrageenan (100 µg·0.1 mL<sup>-1</sup>, inflamed). Concentrations of IL-1β (A), TNF-α (B) and CINC-1 (C) were determined by ELISA in serum 4 h after the induction of pleurisy. Data are expressed as mean ± SEM for six animals in each group. There was a significant decrease in CINC-1 levels in the serum from BbCI-treated inflamed rats. \**P* < 0.05 versus corresponding values in saline-treated inflamed rats.

Our studies led to the conclusion that BbCI has potential anti-inflammatory actions by inhibiting the activity of proteolytic enzymes in the inflammatory site, and the release of BK and CINC-1, and thus decreasing neutrophil migration during the course of the inflammatory response. We should point out that BbCI had no effect in non-inflamed animals. Thus, this inhibitor may be relevant for investigation of other pathological processes in which neutrophil elastase, cathepsin G and cathepsin L are involved.

## Acknowledgements

We are grateful to Lucimeire A de Santana for technical assistance, and to CAPES, CNPq and FAPESP for providing financial support.

## Conflict of interest

The authors declare no conflict of interest.

## References

- Araujo AP, Hansen D, Vieira DF, Oliveira C, Santana LA, Beltramini LM *et al.* (2005). Kunitz-type *Bauhinia bauhinioides* inhibitors devoid of disulfide bridges: isolation of the cDNAs, heterologous expression and structural studies. *Biol Chem* 386: 561–568.
- Baez S (1973). A method for in-line measurement of lumen and wall of microscopic vessels *in vivo*. *Microvasc Res* 5: 299–308.
- Campbell SJ, Hughes PM, Iredale JP, Wilcockson DC, Waters S, Docagne F *et al.* (2003). CINC-1 is an acute-phase protein induced by focal brain injury causing leukocyte mobilization and liver injury. *Faseb J* 17: 1168–1170.
- Coussens LM, Werb Z (2002). Inflammation and cancer. *Nature* 420: 860–867.
- Davis MJ (1987). Determination of volumetric flow in capillary tubes using an optical Doppler velocimeter. *Microvasc Res* 34: 223–230.
- Desmazes C, Gauthier F, Lalmanach G (2001). Cathepsin L, but not cathepsin B, is a potential kininogenase. *Biol Chem* 382: 811–815.
- Di Rosa M (1972). Biological properties of carrageenan. *J Pharm Pharmacol* 24: 89–102.
- Farsky SP, Sannomiya P, Garcia-Leme J (1995). Secreted glucocorticoids regulate leukocyte–endothelial interactions in inflammation. A direct vital microscopic study. *J Leukoc Biol* 57: 379–386.
- Fortes ZB, Farsky SP, Oliveira MA, Garcia-Leme J (1991). Direct vital microscopic study of defective leukocyte–endothelial interaction in diabetes mellitus. *Diabetes* 40: 1267–1273.
- Gore RW, Bohlen HG (1977). Microvascular pressures in rat intestinal muscle and mucosal villi. *Am J Physiol* 233: H685–H693.
- Gozzo AJ, Nunes VA, Carmona AK, Nader HB, von Dietrich CP, Silveira VL *et al.* (2002). Glycosaminoglycans affect the action of human plasma kallikrein on kininogen hydrolysis and inflammation. *Int Immunopharmacol* 2: 1861–1865.
- Harada Y, Hatanaka K, Kawamura M, Saito M, Ogino M, Majima M *et al.* (1996). Role of prostaglandin H synthase-2 in prostaglandin E2 formation in rat carrageenin-induced pleurisy. *Prostaglandins* 51: 19–33.
- Ishihara K, Yamaguchi Y, Uchino S, Furuhashi T, Yamada S, Kihara S *et al.* (2006). ICAM-1 signal transduction in cells stimulated with neutrophil elastase. *Dig Dis Sci* 51: 2102–2112.
- Jung HI, Kim SI, Ha KS, Joe CO, Kang KW (1995). Isolation and characterization of guamerin, a new human leukocyte elastase inhibitor from *Hirudo nipponia*. *J Biol Chem* 270: 13879–13884.
- Kasama T, Miwa Y, Isozaki T, Odai T, Adachi M, Kunkel SL (2005). Neutrophil-derived cytokines: potential therapeutic targets in inflammation. *Curr Drug Targets Inflamm Allergy* 4: 273–279.
- Kohl J, Baelder R, Lewkowich IP, Pandey MK, Hawlisch H, Wang L *et al.* (2006). A regulatory role for the C5a anaphylatoxin in type 2 immunity in asthma. *J Clin Invest* 116: 783–796.
- Korkmaz B, Moreau T, Gauthier F (2008). Neutrophil elastase, proteinase 3 and cathepsin G: physicochemical properties, activity and physiopathological functions. *Biochimie* 90: 227–242.
- Macedo ML, de Sa CM, Freire MD, Parra JR (2004). A Kunitz-type inhibitor of coleopteran proteases, isolated from *Adenanthera pavonina* L. seeds and its effect on *Callosobruchus maculatus*. *J Agric Food Chem* 52: 2533–2540.
- Mezyk-Kopec R, Bzowska M, Bzowska M, Mickowska B, Mak P, Potempa J *et al.* (2005). Effects of elastase and cathepsin G on the levels of membrane and soluble TNFalpha. *Biol Chem* 386: 801–811.
- Moraes TJ, Chow CW, Downey GP (2003). Proteases and lung injury. *Crit Care Med* 31: S189–S194.
- Moreau ME, Garbacki N, Molinaro G, Brown NJ, Marceau F, Adam A (2005). The kallikrein–kinin system: current and future pharmacological targets. *J Pharmacol Sci* 99: 6–38.
- Morris CJ (2003). Carrageenan-induced paw edema in the rat and mouse. *Methods Mol Biol* 225: 115–121.
- Nathan C (2002). Points of control in inflammation. *Nature* 420: 846–852.

- Neuhof C, Oliva ML, Maybauer D, Maybauer M, de Oliveira C, Sampaio MU *et al.* (2003). Effect of plant Kunitz inhibitors from *Bauhinia bauhinoides* and *Bauhinia rufa* on pulmonary edema caused by activated neutrophils. *Biol Chem* 384: 939–944.
- Oliva ML, Sampaio UM (2008). *Bauhinia* Kunitz-type proteinase inhibitors: structural characteristics and biological properties. *Biol Chem* 389: 1007–1013.
- Oliva ML, Andrade SA, Batista IF, Sampaio MU, Juliano M, Fritz H *et al.* (1999a). Human plasma kallikrein and tissue kallikrein binding to a substrate based on the reactive site of a factor Xa inhibitor isolated from *Bauhinia unguolata* seeds. *Immunopharmacology* 45: 145–149.
- Oliva ML, Mendes CR, Juliano MA, Chagas JR, Rosa JC, Greene LJ *et al.* (1999b). Characterization of a tissue kallikrein inhibitor isolated from *Bauhinia bauhinoides* seeds: inhibition of the hydrolysis of kininogen related substrates. *Immunopharmacology* 45: 163–169.
- Oliva ML, Souza-Pinto JC, Batista IF, Araujo MS, Silveira VF, Auerswald EA *et al.* (2000). *Leucaena leucocephala* serine proteinase inhibitor: primary structure and action on blood coagulation, kinin release and rat paw edema. *Biochim Biophys Acta* 1477: 64–74.
- Oliva ML, Santomauro-Vaz EM, Andrade SA, Juliano MA, Pott VJ, Sampaio MU *et al.* (2001). Synthetic peptides and fluorogenic substrates related to the reactive site sequence of Kunitz-type inhibitors isolated from *Bauhinia*: interaction with human plasma kallikrein. *Biol Chem* 382: 109–113.
- Oliva ML, Andrade SA, Juliano MA, Sallai RC, Torquato RJ, Sampaio MU *et al.* (2003). Kinetic characterization of factor Xa binding using a quenched fluorescent substrate based on the reactive site of factor Xa inhibitor from *Bauhinia unguolata* seeds. *Curr Med Chem* 10: 1085–1093.
- de Oliveira C, Santana LA, Carmona AK, Cezari MH, Sampaio MU, Sampaio CA *et al.* (2001). Structure of cruzipain/cruzain inhibitors isolated from *Bauhinia bauhinoides* seeds. *Biol Chem* 382: 847–852.
- Owen CA, Campbell EJ (1995). Neutrophil proteinases and matrix degradation. The cell biology of pericellular proteolysis. *Semin Cell Biol* 6: 367–376.
- Owen CA, Campbell EJ (1999). The cell biology of leukocyte-mediated proteolysis. *J Leukoc Biol* 65: 137–150.
- Panes J, Kurose I, Rodriguez-Vaca D, Anderson DC, Miyasaka M, Tso P *et al.* (1996). Diabetes exacerbates inflammatory responses to ischemia–reperfusion. *Circulation* 93: 161–167.
- Pease JE, Williams TJ (2006). The attraction of chemokines as a target for specific anti-inflammatory therapy. *Br J Pharmacol* 147 (Suppl. 1): S212–S221.
- Pham CT (2006). Neutrophil serine proteases: specific regulators of inflammation. *Nat Rev Immunol* 6: 541–550.
- Puzer L, Cotrin SS, Alves MF, Egborge T, Araujo MS, Juliano MA *et al.* (2004). Comparative substrate specificity analysis of recombinant human cathepsin V and cathepsin L. *Arch Biochem Biophys* 430: 274–283.
- Rhodin JA (1986). Perfusion and superfusion fixation effects on rat mesentery microvascular beds. Intravital and electron microscope analyses. *J Submicrosc Cytol* 18: 453–470.
- Sallenave JM (2000). The role of secretory leukocyte proteinase inhibitor and elafin (elastase-specific inhibitor/skin-derived antileukoprotease) as alarm antiproteinases in inflammatory lung disease. *Respir Res* 1: 87–92.
- Shimamoto K, Ando T, Tanaka S, Nakahashi Y, Nishitani T, Hosoda S *et al.* (1982). An improved method for the determination of human blood kinin levels by sensitive kinin radioimmunoassay. *Endocrinol Jpn* 29: 487–494.
- Silveira VL, Limaos EA, Nunes DW (1995). Participation of the adrenal gland in the anti-inflammatory effect of polyunsaturated diets. *Mediators Inflamm* 4: 359–363.
- Skrzydewska E, Sulkowska M, Koda M, Sulkowski S (2005). Proteolytic–antiproteolytic balance and its regulation in carcinogenesis. *World J Gastroenterol* 11: 1251–1266.
- Soehnlein O, Weber C, Lindbom L (2009). Neutrophil granule proteins tune monocytic cell function. *Trends Immunol* 30: 538–546.
- Sumikawa JT, Nakahata AM, Fritz H, Mentele R, Sampaio MU, Oliva ML (2006). A Kunitz-type glycosylated elastase inhibitor with one disulfide bridge. *Planta Med* 72: 393–397.
- Travis J, Salvesen GS (1983). Human plasma proteinase inhibitors. *Annu Rev Biochem* 52: 655–709.
- Vaz A, Tozzi A (2005). Synopsis of *Bauhinia* sect. *Pauletia* (Cav.) DC. (Leguminosae: Caesalpinioideae: Cercideae) in Brazil. *Rev Bras Bot* 28: 477–491.
- Vinegar R, Truax JF, Selph JL, Voelker FA (1982). Pathway of onset, development, and decay of carrageenan pleurisy in the rat. *Fed Proc* 41: 2588–2595.
- Vogt W (1995). Oxidation of methionyl residues in proteins: tools, targets, and reversal. *Free Radic Biol Med* 18: 93–105.
- Weiss SJ (1989). Tissue destruction by neutrophils. *N Engl J Med* 320: 365–376.
- Wohlers M, Xavier RA, Oyama LM, Ribeiro EB, do Nascimento CM, Casarini DE *et al.* (2005). Effect of fish or soybean oil-rich diets on bradykinin, kallikrein, nitric oxide, leptin, corticosterone and macrophages in carrageenan stimulated rats. *Inflammation* 29: 81–89.
- Wu Y, Stabach P, Michaud M, Madri JA (2005). Neutrophils lacking platelet-endothelial cell adhesion molecule-1 exhibit loss of directionality and motility in CXCR2-mediated chemotaxis. *J Immunol* 175: 3484–3491.

Zabel BA, Zuniga L, Ohyama T, Allen SJ, Cichy J, Handel TM *et al.* (2006). Chemoattractants, extracellular proteases, and the integrated host defense response. *Exp Hematol* 34: 1021–1032.

Zimmermann F, Lautenschlager K, Heppert V, Wentzensen A, Hansch GM, Wagner C (2005). Expression of elastase on polymorphonuclear

neutrophils in vitro and in vivo: identification of CD11b as ligand for the surface-bound elastase. *Shock* 23: 216–223.

Zlotnik A, Yoshie O (2004). Chemokines: a new classification system and their role in immunity. *Immunity* 12: 121–127.