

REVIEW

Modulation of diabetic retinopathy pathophysiology by natural medicines through PPAR- γ -related pharmacology

Min K Song, Basil D Roufogalis and Tom H W Huang

Herbal Medicines Research and Education Centre, Faculty of Pharmacy, The University of Sydney, NSW, Australia

Diabetic retinopathy (DR) is one of the most common microvascular complications of diabetes and remains a major cause of preventable blindness among adults at working age. DR involves an abnormal pathology of major retinal cells, including retinal pigment epithelium, microaneurysms, inter-retinal oedema, haemorrhage, exudates (hard exudates) and intraocular neovascularization. The biochemical mechanisms associated with hyperglycaemic-induced DR are through multifactorial processes. Peroxisome proliferator-activated receptor- γ (PPAR- γ) plays an important role in the pathogenesis of DR by inhibiting diabetes-induced retinal leukostasis and leakage. Despite DR causing eventual blindness, only a few visual or ophthalmic symptoms are observed until visual loss develops. Therefore, early medical interventions and prevention are the current management strategies. Laser photocoagulation therapy is the most common treatment. However, this therapy may cause retinal damage and scarring. Herbal and traditional natural medicines may provide an alternative to prevent or delay the progression of DR. This review provides an analysis of the therapeutic potential of herbal and traditional natural medicines or their active components for the slowdown of progression of DR and their possible mechanism through the PPAR- γ pathway.

LINKED ARTICLE

This article is commented on by Costa and Ciccodicola, pp. 1–3 of this issue. To view this commentary visit <http://dx.doi.org/10.1111/j.1476-5381.2011.01443.x>

Abbreviations

15-dPGJ₂, 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂; AGEs, advanced glycation end products; BRB, blood retinal barrier; COX-2, cyclooxygenase-2; DR, diabetic retinopathy; ICAM-1, intercellular cell adhesion molecule-1; iNOS, inducible nitric oxide synthase; MMP-9, matrix metalloproteinase-9; NF- κ B, nuclear factor-kappaB; PPAR- γ , peroxisome proliferator-activated receptor- γ ; ROS, reactive oxygen species; TNF- α , tumour necrosis factor; VEGF, vascular endothelial growth factor

Introduction

Diabetic retinopathy (DR) is one of the most common microvascular complications of diabetes and one of the leading causes of blindness worldwide (Fong *et al.*, 2004a). The prevalence of DR increases with the duration of diabetes, and nearly all patients with type I diabetes and more than 60% with type II diabetes have some degree of retinopathy after 20 years (Fong *et al.*, 2003; 2004b; Williams *et al.*, 2004). Therefore, early detection and prevention are the current management strategies (Ciulla *et al.*, 2003). Chronic hyperglycaemia is

believed to be the primary pathogenic factor for inducing damage to retinal cells (Knudsen *et al.*, 2002; Hammes, 2005; Yanagi, 2008). However, the mechanisms that lead to DR are not fully understood (West *et al.*, 2006). DR is characterized by microaneurysms, inter-retinal oedema, haemorrhages, exudates (hard exudates) and intraocular pathological neovascularization (Fong *et al.*, 2004a,b). Laser photocoagulation therapy is the most common treatment modality for DR. However, this therapy may damage neural tissue resulting in the deterioration of vision (Bloomgarden, 2007). Therefore, development of new therapeutic strategies for the treatment

Correspondence

Tom Hsun-Wei Huang, Faculty of Pharmacy, The University of Sydney, NSW 2006, Australia.
E-mail: thwh@yahoo.com

Keywords

diabetic retinopathy; peroxisome proliferator-activated receptor- γ ; herbal and traditional natural medicines

Received

10 February 2011

Accepted

10 March 2011

of excessive retinal vasopermeability and angiogenic changes are the basis for further research focus (Garcia *et al.*, 2008).

The peroxisome proliferator-activated receptor- γ (PPAR- γ) is a ligand-inducible transcription factor that belongs to a large superfamily, comprising the nuclear receptors for steroids, thyroid hormones and retinoids (Malchiodi-Albedi *et al.*, 2008). PPAR- γ plays an important role in adipogenesis, glucose metabolism, angiogenesis and inflammation (Sarafidis and Bakris, 2006). Synthetic PPAR- γ ligands, the thiazolidinediones, including pioglitazone and rosiglitazone, improve insulin resistance in diabetic patients, and have become one of the most popular anti-diabetic drugs in developed countries (Tilg and Moschen, 2008; Tontonoz and Spiegelman, 2008; Zinn *et al.*, 2008). In addition, natural PPAR- γ ligands, such as 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15-dPGJ₂) have very potent anti-inflammatory effects that also modulate cellular defence against oxidative stress (Ershov and Bazan, 2000; Straus and Glass, 2001). Moreover, there is accumulating data to show that PPAR- γ activators exert anti-inflammatory, antioxidative and anti-proliferative effects in various cells including major retinal cells (Yamagishi *et al.*, 2007; Gerry and Pascual, 2008; Giaginis *et al.*, 2008; Sulistio *et al.*, 2008; Yanagi, 2008).

Plants remain an important source of material for maintaining human health with unparalleled diversity and have improved the quality of life for centuries (Li *et al.*, 2005). Moreover, medicinal plants are an abundant source of biologically active molecules that play an important role in past and modern medicine and which act as a 'stepping stone' for the discovery of novel pharmacologically active ligands (Barnett *et al.*, 2003). In recent years, numerous plant-derived ligands of PPAR- γ receptors have been identified (Huang *et al.*, 2005). However, very limited research has been undertaken on medicinal plant-derived PPAR- γ ligands in the modulation of DR-related pathophysiology. Thus, the current review is a detailed discussion summarizing the current understanding of the potential of herbal and traditional natural medicines for management and potential reversal of DR-related pathogenesis.

Diabetic retinopathy

Diabetes damages all the major cells of the retina, vascular cells (endothelial cells and pericytes; Hammes *et al.*, 1995; Mizutani *et al.*, 1996) and pigment epithelial cells (Decanini *et al.*, 2008). Basement membrane thickening, pericyte dropout and retinal capillary non-perfusion occur prior to the damage, which changes the production pattern of a number of mediators, such as growth factors, vasoactive agents, coagulation factors and adhesion molecules. These result in increased blood flow and capillary diameter, proliferation of the extracellular matrix and thickening of basal membranes, altered cell turnover (apoptosis, proliferation, hypertrophy) and procoagulant/proaggregant patterns, and finally angiogenesis with tissue remodelling. These pathological changes cause increase retinal vasopermeability and breakdown of blood retinal barrier (BRB), resulting in retinal haemorrhages, swelling, exudates and retinal detachment (Gardner *et al.*, 2000; Ciulla *et al.*, 2003; Yam and Kwok, 2007).




Figure 1

Pathophysiological cascades implicated in the development of diabetic retinopathy. Schematic representation showing the involvement of PPAR- γ and its receptor system in the progression of diabetic retinopathy (adapted from Yamagishi *et al.*, 2007). VCAM-1, vascular cell adhesion molecule-1.

The underlying pathophysiological mechanisms associated with hyperglycaemic-induced DR are through excessive formation of advanced glycation end products (AGEs) and production of excessive oxidative stress (Ciulla *et al.*, 2003; Abu El-Asrar *et al.*, 2009). Moreover, these biochemical mechanisms lead to a cascade of events, such as promotion of apoptosis, inflammation and angiogenesis, which induce damage to diabetic retina, leading to DR (Ciulla *et al.*, 2003; Abu El-Asrar *et al.*, 2009; Figure 1).

Advanced glycation end products (AGEs) in diabetic retinopathy

Advanced glycation end products are associated with modification of proteins or lipids that are generated from intermediate glycation products by non-enzymatic reaction of glucose with protein side chains (Schmidt *et al.*, 1994; Goh and Cooper, 2008). These intermediate glycation products undergo further condensation, dehydration or rearrangement, leading to eventual irreversible AGEs formation (Chu and Ali, 2008). AGEs formation occurs normally over time whereas an accelerated rate of AGE formation is accompanied by hyperglycaemia (Munch *et al.*, 1998).

The accumulated AGEs products are detected in the retinal blood vessel wall, responsible for mediating the pathological angiogenesis and hyper-permeability in retina. Several bodies of evidence suggest that the interaction between AGEs and their receptor activates nicotinamide adenine dinucleotide phosphate oxidase and enhances the formation of oxygen radicals, with subsequent activation and translocation of nuclear factor-kappaB (NF- κ B), followed by release of pro-inflammatory cytokines and growth factors (Abu El-Asrar *et al.*, 2009). Moreover, AGEs enhance apoptosis in retinal pericytes, corneal endothelial cells, neuronal cells and renal

mesangial cells through increased oxidative stress or via induced expression of pro-apoptotic cytokines (Kasper *et al.*, 2000; Denis *et al.*, 2002; Kaji *et al.*, 2003). Indeed, AGEs induce apoptosis, angiogenesis, breakdown of BRB, and leukocyte adhesion in the retina. Thus, AGEs are detrimental to the retinal vasculature and contribute to the pathogenesis of DR (Stitt, 2001; Sato *et al.*, 2006).

Oxidative stress in diabetic retinopathy

Oxidative stress appears when there is a serious imbalance between generation of reactive oxygen species (ROS) and its clearance by antioxidant defences (Ceriello *et al.*, 2001; Brownlee, 2005). Activation of receptor for advanced glycation end products results in production of oxidative stress (conversely, glycation itself is promoted by oxidative stress), and subsequent activation of NF- κ B transcription factor in microvascular endothelial cells, which is considered to be linked to endothelial dysfunction (Moore *et al.*, 2003; Vincent *et al.*, 2007). Retina, a tissue rich in polyunsaturated fatty acid, uses more oxygen and glucose oxidation than any other tissue in the body, and is very susceptible to damage (Schmidt *et al.*, 2003). Diabetic-induced oxidative stress, followed by activation of NF- κ B in the retina, are early events in the pathogenesis of DR (Kowluru *et al.*, 1996; 2001; Kowluru, 2005; Obrosova and Julius, 2005). Moreover, oxidative stress has been linked to the accelerated apoptosis of retinal capillary cells and microvascular abnormalities in DR (Allen *et al.*, 2005).

Nuclear factor-kappaB in diabetic retinopathy

Nuclear factor-kappaB is a multi-protein complex which can activate many kinds of genes involved in cellular functions. Pathogenic stimuli allow NF- κ B to enter the nucleus, and to bind to DNA recognition sites in regulatory regions of target genes (Schreck *et al.*, 1992; Baeuerle and Henkel, 1994; Boileau *et al.*, 2003). NF- κ B is required for maximal transcription of many pro-inflammatory molecules thought to be important in the generation of inflammation, including molecules (intracellular adhesion molecule 1), critical enzymes [inducible nitric oxidase synthase, cyclooxygenase-2 (COX-2)] and a number of cytokines (interleukin-1 β , tumour necrosis factor- α , IL-6) (Siebenlist *et al.*, 1994; Blackwell and Christman, 1997). The activation of NF- κ B is considered a key signalling pathway by which high glucose induces apoptosis in endothelial cells (Du *et al.*, 1999). In the retina, NF- κ B is localized in sub-retinal membranes and in microvessels (Hammes *et al.*, 1999) and its activation is considered responsible for the accelerated loss of pericytes observed in DR (Romeo *et al.*, 2002).

Inflammation in diabetic retinopathy

In recent years inflammation has been linked to vascular leakage in DR, at least in part (Adamis, 2002; Jousen *et al.*, 2002). Hyperglycaemia is a contributing risk factor for the development of vascular dysfunction and production of inflammatory markers (Haubner *et al.*, 2007; Tawfik *et al.*, 2009). Indeed, pro-inflammatory cytokines, chemokines and other inflammatory mediators play an important role in the pathogenesis of DR. These lead to persistent low-grade inflammation, the adhesion of leukocytes to the retinal vasculature

(leukostasis), breakdown of BRB and neovascularization with subsequent sub-retinal fibrosis or disciform scarring (Carmo *et al.*, 1999; Malchiodi-Albedi *et al.*, 2008). Several inflammatory molecules are involved in the pathogenesis of DR, including tumour necrotic factor (TNF- α), fibronectin, COX-2, intercellular cell adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 and matrix metalloproteinase-9 (MMP-9) (Miyamoto and Ogura, 1999; Yuuki *et al.*, 2001; Jousen *et al.*, 2004).

Angiogenesis in diabetic retinopathy

Angiogenesis is defined as the growth of new vessels from pre-existing capillaries which is a complex process comprising endothelial cell proliferation, migration, extracellular proteolysis, tube formation and vessel remodelling (Beck and D'Amore, 1997). In retina, vascular endothelial growth factor (VEGF) is the major angiogenic factor for neovascularization and vascular leakage via the mitogen-activated protein kinase pathway (Miller *et al.*, 1994; Kliffen *et al.*, 1997; De Luca *et al.*, 2008). Additional pro-angiogenic factors, including MMP-9, are also required for the process of ocular neovascularization through either synergistic effects with angiogenic factor or as a stimulant for the secretion of angiogenic factors (Hollborn *et al.*, 2007). In addition, MMP-9 expression acts as a factor in increasing vascular permeability in ocular neovascularization (Herzlich *et al.*, 2008).

Apoptosis in diabetic retinopathy

Apoptosis is programmed cell death and is characterized by chromatin condensation, fragmentation and formation of apoptotic bodies that can be triggered by various signals (Nagata, 1997; Li *et al.*, 1998; Leal *et al.*, 2009). In healthy organisms apoptosis and cell proliferation are in homeostasis to maintain a physiological balance during normal embryonic development and in the cell turnover throughout life (van den Oever *et al.*, 2010; Zeng *et al.*, 2010). In contrast, retinal microvascular cells are lost selectively via apoptosis before other histopathology is detectable in diabetes (Mizutani *et al.*, 1996; Kern *et al.*, 2000; Kowluru and Odenbach, 2004). Moreover, it has been well established that apoptosis represents a final common pathway of cell loss and hence vision loss (Doonan *et al.*, 2009). Therefore, apoptosis plays an important role in the progression and pathogenesis of DR (Allen *et al.*, 2005; Leal *et al.*, 2009). As oxidative stress is closely linked to apoptosis in diabetes, oxidative stress-induced apoptotic episodes have been demonstrated by retinal abnormalities, potential visual changes and the onset of the first vascular change (Lopes de Faria *et al.*, 2001; Kowluru, 2005). Moreover, apoptotic cell death in retinal regions is a probable stimulus for the increased expression of molecules that enhance the breakdown of the BRB and lead to vascular proliferation (Henkind, 1978; Patz, 1982). Several studies have shown that retinal pigment epithelial (RPE) cells, Glial cells (Zeng *et al.*, 2009) and retinal pericytes (Leal *et al.*, 2009; Zeng *et al.*, 2010) undergo high glucose-induced apoptosis. High glucose causes activation of several proteins involved in apoptotic cell death, including members of the caspase and Bcl-2 family (Allen *et al.*, 2005).

Pathogenesis of retinal pigment epithelium in diabetic retinopathy

The RPE cells form a monolayer between the neuroretina and the choriocapillaris which are the essential component of the outer BRB that maintain physiological and structural balance within the retina (Bok, 1993; Rizzolo, 1997). The main characteristics of RPE cells are the presence of tight junctions at the apical side of their lateral molecules, which limit access of blood components to the retina. Other important functions include regulation of transport of nutrients to the photoreceptors, phagocytosis of damaged or old rod outer segments and production of growth factors (Holtkamp *et al.*, 2001). Moreover, RPE and photoreceptors are particularly susceptible to oxidative stress because of high oxygen consumption by photoreceptors (Beatty *et al.*, 2000). In response to damage caused by the hyperglycaemic condition, RPE cells migrate and proliferate, leading to a breakdown in adhesion between the RPE and the choroidal capillaries, followed by BRB breakdown compromising blood flow within the RPE layer and leading to eventual retinal oedema (Kennedy *et al.*, 1995). These cascade episodes trigger the serum components and inflammatory cells to enter the vitreous cavity and sub-retinal space, exposing the RPE cells to a variety of cytokines, pro-inflammatory mediators, extracellular matrix proteins and growth factors, causing DR (Kimoto *et al.*, 2004). Several studies have shown that the expression of angiogenic cytokines, growth factors (e.g. VEGF) and metalloproteinases (e.g. MMP-9) are produced by RPE (Grossniklaus *et al.*, 2002). Moreover, the combined effects from chronic sustained inflammation and ROS generation promote the development of RPE damage (Winkler *et al.*, 1999; SanGiovanni and Chew, 2005; de Jong, 2006). There is a strong body of evidence on the prevalence of the variety of anti-angiogenic agents, anti-inflammatory agents, antioxidants and anti-fibrogenesis present in the retinal regions for slowing down the progression of DR (Winkler *et al.*, 1999; Zhang *et al.*, 2006; Kern, 2007; Cheng *et al.*, 2008). Moreover, the discovery of biochemical mechanisms that either protect cells or promote cellular recovery from damage is important (Geiger *et al.*, 2005).

Peroxisome proliferator-activated receptor- γ and diabetic retinopathy

Peroxisome proliferator-activated receptor- γ is heterogeneously expressed in the mammalian eye, prominently present in the retinal pigmented epithelium, photoreceptor outer segments, choriocapillaries and retina (Murata *et al.*, 2000; Sarayba *et al.*, 2005; Herzlich *et al.*, 2008). Recent studies have shown that retinal expression of PPAR- γ was suppressed in experimental models of diabetes and in endothelial cells treated with high glucose (Tawfik *et al.*, 2009). Moreover, PPAR- γ ligands are potent inhibitors of corneal angiogenesis and neovascularization (Xin *et al.*, 1999; Muranaka *et al.*, 2006). Administration of 15d-PGJ₂ inhibited VEGF-stimulated angiogenesis in rat cornea (Xin *et al.*, 1999). Similarly, choroidal neovascularization was markedly reduced by intravitreal injection of troglitazone. Laser photocoagulation-induced lesions in rat and monkey eyes showed significantly less leakage in troglitazone-treated animals (Murata *et al.*, 2000).

Troglitazone demonstrated that intravitreal injections significantly inhibited the percentage of lesions as well as leakage per lesion (Murata *et al.*, 2000). Increase in catalase mRNA has been shown when using known PPAR- γ agonists rosiglitazone and ciglitazone in rat brain microvascular endothelium cells, one of the cell types damaged during inflammatory responses induced by ROS generation (Girnun *et al.*, 2002). In neonatal mice, intravitreal injection of rosiglitazone or troglitazone inhibited development of new retinal vessels. In the same study, thiazolidinediones have been found to inhibit retinal endothelial cell proliferation, migration and tube formation in response to VEGF treatment (Touyz and Schiffrin, 2006). In addition, rosiglitazone inhibits both the retinal leukostasis and retinal leakage observed in experimental diabetic rats which leads to the aggravation of retinal leukostasis and retinal leakage in diabetic mice (Muranaka *et al.*, 2006). Moreover, rosiglitazone has been shown to delay the onset of DR (Shen *et al.*, 2008). These findings suggest that PPAR- γ is involved in the pathogenesis of DR (Figure 1).

Peroxisome proliferator-activated receptor- γ and AGEs

Peroxisome proliferator-activated receptor- γ ligands provide a significant prevention of AGEs-induced microvascular complications, including DR (Dolhofer-Bliesener *et al.*, 1996; Watson *et al.*, 2005). Indeed, a body of evidence has shown that PPAR- γ ligands inhibit the formation of AGEs (Rahbar *et al.*, 2000; Sobal *et al.*, 2005). The inhibitory action of PPAR- γ ligands on AGE formation may be ascribed to their antioxidative properties (Yamagishi *et al.*, 2007; Gerry and Pascual, 2008; Giaginis *et al.*, 2008; Sulistio *et al.*, 2008). In addition, activation of PPAR- γ by rosiglitazone inhibits AGE-induced inducible NO synthase expression, nitrite release, fibronectin and type IV collagen production (Chang *et al.*, 2004; Yu *et al.*, 2007). Moreover, rosiglitazone has been shown to inhibit extracellular matrix accumulation, fibronectin and type IV collagen in AGE-injected rats, and also inhibits the AGE-induced proliferation and NO production in cardiac fibroblasts (Yu *et al.*, 2007; Li *et al.*, 2008).

Peroxisome proliferator-activated receptor- γ in NF- κ B, inflammatory mediators and angiogenesis

Peroxisome proliferator-activated receptor- γ plays a critical role in a variety of biological processes, including inflammation and angiogenesis, mediated through the inhibition of NF- κ B (Rosen and Spiegelman, 2001; Lee *et al.*, 2006; Sung *et al.*, 2006; Kim *et al.*, 2007). In streptozotocin-induced DR, rosiglitazone was shown to inhibit both retinal leukostasis and retinal leakage by the inhibition of NF- κ B activation, with consequent suppression of ICAM-1 expression (Muranaka *et al.*, 2006). In addition, one recent body of evidence has shown that the suppression of PPAR- γ in diabetic retina is associated with the activation of NF- κ B target gene expression (Remels *et al.*, 2009; Tawfik *et al.*, 2009). The activation of PPAR- γ inhibits the pro-inflammatory pathways, including cytokine secretion (Uchimura *et al.*, 2001; Wong *et al.*, 2001) and inducible nitric oxide synthase (iNOS) expression (Petrova *et al.*, 1999; Reilly *et al.*, 2001) in a variety of cell lines. Indeed, PPAR- γ ligands have also been shown to

suppress the expression of iNOS, observed at both mRNA and protein levels (Ricote *et al.*, 1998b). The murine iNOS promoter contains 24 transcriptional factor-binding sites, including NF- κ B transcription factor (Chang *et al.*, 2004). Inhibition of ICAM-1 expression and retinal vascular leakage in experimental diabetes has been shown by rosiglitazone, and the increase in the same parameters by depletion of the gene encoding PPAR- γ (Muranaka *et al.*, 2006). PPAR- γ ligands have also been shown to inhibit the expression of VEGF receptors and the subsequent activation of downstream signalling pathway (Xin *et al.*, 1999; Murata *et al.*, 2001). Moreover, rosiglitazone has been shown to inhibit retinal neovascularization in oxygen-induced retinopathy by a mechanism downstream from VEGF-induced angiogenesis (Murata *et al.*, 2001). In addition, it has been suggested that ICAM-1 is involved in VEGF-induced leukocyte-endothelial cell interactions and subsequent BRB breakdown in the diabetic retina (Miyahara *et al.*, 2004). COX-2 has emerged as a key regulator of inflammatory angiogenesis and its expression is induced by VEGF (Wu *et al.*, 2003; Gately and Li, 2004). Moreover, PPAR- γ activation inhibits VEGF-mediated angiogenesis through the modulation of the stimulated COX-2 expression and activity (Scoditti *et al.*, 2009).

Peroxisome proliferator-activated receptor- γ and apoptosis

Apoptosis is a complex process, involving a multitude of signalling pathways that regulate the activities of pro- and anti-apoptotic members of Bcl-2 family of proteins which play an important role in various cell types (Fehlberg *et al.*, 2003; Bonne, 2005; Fuenzalida *et al.*, 2007). Moreover, oxidative stress can induce mitochondria dysfunction, followed by cytochrome c release and subsequent activation of caspases, a group of enzymes that execute apoptosis (Danial and Korsmeyer, 2004; Lin and Beal, 2006). A recent study has shown that rosiglitazone protects against oxidative stress-induced apoptosis through up-regulation of anti-apoptotic Bcl-2 family proteins (Ren *et al.*, 2009). Moreover, rosiglitazone and PPAR- γ over-expression protect against apoptosis induced by oxygen and glucose deprivation followed by re-oxygenation and up-regulation of Bcl-2 (Wu *et al.*, 2009). In contrast, down-regulation of NF- κ B activation by PPAR- γ ligands protects the cells from destruction via the apoptotic pathways (Grey *et al.*, 1999; Baker *et al.*, 2001). A screen of FDA-approved compounds identified rosiglitazone as a novel anti-apoptotic agent in retinal cells in both *in vivo* and *in vitro* (Doonan *et al.*, 2009). In addition, troglitazone has shown cytoprotective activity in apoptotic-induced ARPE-19 cells (Rodrigues *et al.*, 2010). One further study has indicated that 15d-PGJ₂ helps RPE cells to maintain mitochondrial integrity by prevention of cytochrome c release and subsequent activation of the apoptosis pathway (Garg and Chang, 2004; Chang and So, 2008).

Peroxisome proliferator-activated receptor- γ and RPE cells

A number of studies have shown that RPE might be the prime target for oxidative stress and PPAR- γ ligands modulate cellular defence against the oxidative stress (Chang *et al.*, 2008). 15-dPGJ₂ protects RPE cells from oxidative stress by elevating GSH and enhancing mitogen-activated protein kinase activa-

tion through the PPAR- γ independent pathway (Qin *et al.*, 2006). In addition, 15-dPGJ₂, independent of its PPAR γ activity, protects RPE cells from oxidative injury by raising intracellular GSH levels and extending hydrogen peroxide-induced activation of JNK and p38, suggesting the possible application of the agents in preventing ocular diseases from oxidative stress (Garg and Chang, 2003; Qin *et al.*, 2006). It has also been shown that PPAR- γ ligands inhibit VEGF-induced choroidal angiogenesis, without apparent toxicity to the adjacent retina, in a laser-induced model of choroidal neovascularization (Murata *et al.*, 2000). A recent study has shown that PPAR- γ inhibits the expression of MMP-9 in response to 15-dPGJ₂, and synthetic PPAR- γ ligands by antagonizing the activities of NF- κ B to decrease vascular permeability or neovascularization (Ricote *et al.*, 1998a,b; Herzlich *et al.*, 2008). Moreover, rosiglitazone inhibits the endothelial effects of VEGF (Murata *et al.*, 2000). Interestingly, synthetic PPAR γ agonists, AGN195037 and rosiglitazone, have been shown to be less effective in the protection of RPE cells against hydrogen peroxide-induced injury. Hence, the cytoprotective effect of this agent has not been conclusively demonstrated (Qin *et al.*, 2006; Chang *et al.*, 2008). There is growing interest in natural products with combined anti-glycation and antioxidant properties as they may have reduced toxicity. Indeed, a number of plant-derived flavonoids with antioxidant activity (quercetin, rutin and kaempferol) have been reported to inhibit glycation (Ahmed, 2005).

Herbal and traditional medicines and diabetic retinopathy

In recent years, there has been growing attention to alternative therapies and the therapeutic use of plant-origin natural products (Li *et al.*, 2005). More than 400 traditional medicinal plants and the effective constituents derived from over 800 herbal compounds are reported to possess therapeutic potential or efficacy for the treatment of diabetes (Bailey and Day, 1989; Al-Rowais, 2002; Dey *et al.*, 2002). Herbal and traditional medicines have been globally used for diabetic complications for centuries. Moreover, it may provide an alternative to prevent or delay the progression of DR (Head, 1999). Despite modern pharmacotherapeutics and advancement in an ever changing world of biotechnology, a lack of understanding still exists on the bioactivity of these herbal medicines and related active ligands towards the modulation of DR pathogenesis (Bailey and Day, 1989). This has prompted research to understand the mechanism of action of these compounds and seek new products for better management of diabetes and its complications (Ceylan-Isik *et al.*, 2008). This section summarizes the current research on various active components of herbal and traditional medicines capable of modulating DR pathogenesis (Table 1).

Traditional Chinese medicines in diabetic retinopathy

Traditional Chinese medicines (TCMs) have been used for more than 2000 years in China and other Asian countries with the fundamental aim of targeting not only the treatment of diabetes but also the prevention of its complications (Covington, 2001; Li *et al.*, 2004). The main concept on

Table 1

Modern research on natural medicines capable of modulating diabetic retinopathy-related pathogenesis

Herbal medicines	Active component	Functions	References
Goji berry (<i>Lycium barbarum</i> L.)	<i>Lycium barbarum</i> polysaccharides (LBP)	Beneficial effect for maintaining macular pigment density in age-related macular degeneration by increase fasting plasma zeaxanthin levels in human clinical trial. Effective in glaucoma and modulating immune system in retinal ganglion cells in a rat hypertension model LBP antagonizes glutamate excitotoxicity in rat cortical neurons and increase Bcl-2 level in lens epithelial cells of the whole lens exposed to hydrogen peroxide	(Cheng <i>et al.</i> , 2005) (Chiu <i>et al.</i> , 2009) (Ho <i>et al.</i> , 2009) (Wang <i>et al.</i> , 2002)
Danshen dripping pills	<i>Salviae miltiorrhiae</i>	Improve visual acuity, control micro-haemorrhage Improve ischaemia of the retina and the visual field in early diabetic retinopathy; reduce the number of microaneurysms	(Qi <i>et al.</i> , 2007) (Deng <i>et al.</i> , 2005)
<i>Ginkgo biloba</i>	Ginkgo flavone glycosides	Beneficial outcome for the treatment of diabetic retinopathy when <i>Ginkgo dipyrldolum</i> injection combined with laser photocoagulation therapy in human clinical trials Clinical improvement of diabetic dyschromatopsia in early diabetic retinopathy in a double-blind trial	(Wei <i>et al.</i> , 2005)
<i>Anisodus tanguticus</i>	Anisodamine	Increase neural tissue oxygen tension in retinal hypoxia Increase blood flow and oxygen delivery to the retina-choroid and iris-ciliary body of the eye, and lipid peroxidation in the retinal cells	(Linsenmeier <i>et al.</i> , 1989; Zhang <i>et al.</i> , 1990)
<i>Stephania tetrandra</i> S. Moore (STMS) (Han Fang Ji)	Tetrandrine	Direct effect on retinal capillary of posterior ocular region and suppression of neovascularization of the retinal capillary in diabetic rats	(Liang <i>et al.</i> , 2002)
<i>Radix Puerariae</i>	Not known	The ethanol extract of <i>Radix Puerariae</i> inhibits glycosylation of rat lens protein Discovery of a new key chemical constituent proven to be useful in DR by inhibiting aldose reductase	(Hirakura <i>et al.</i> , 1989) (Duan <i>et al.</i> , 2000)
<i>Gymnema sylvestre</i>	A polyol, conduritol A	Beneficial effect on cataract by inhibiting lens aldose reductase	(Miyatake <i>et al.</i> , 1994)
<i>Astragalus radix</i>	Astragalosides	Potentially useful for the prevention of clinical diabetic complications such as diabetic retinopathy by inhibiting AGE product Controls the balance of sodium, potassium and calcium ions for maintenance of lens transparency. This finding may lead the possibility of prophylactic use in eye diseases such as diabetic cataract	(Motomura <i>et al.</i> , 2009) (Kamei <i>et al.</i> , 1987)
Hachimi-jio-gan (Trrhmannia Eight Formula)	Not known	Beneficial effect for abnormal accumulation of sorbitol and fructose in the diabetic lens Restore disintegration of the inner nuclear layer cells with reduction in rough endoplasmic reticulum and swelling of mitochondria in the diabetic retina Cataract-suppressing effect by inhibiting lens aldose reductase	(Miyatake <i>et al.</i> , 1994) (Preet <i>et al.</i> , 2006)
<i>Trigonella foenum graecum</i> (Fenugreek)	Saponins, Flavonoids, 4-hydroxyisoleucine	In a clinical study, 12 adult diabetic patients were treated with 600 mg of anthocyanosides daily for 2 months, followed by significant decrease in the biosynthesis of connective tissue to protect against diabetic retinopathy In 31 patients suffering from various types of retinopathy, treatment with <i>V. myrtillus</i> extract was associated with a reduced tendency towards retinal haemorrhage	(Boniface and Robert, 1996) (Perossini <i>et al.</i> , 1988)
<i>Vaccinium myrtillus</i> (bilberry)	Anthocyanidins	Protects high glucose-induced ARPE-19 cells by reducing low-grade inflammation mediators, interleukin-6 and interleukin-8, COX-2, VEGF and connexin 43 down-regulations Control retinal haemorrhage absorption and its recurrence in retinopathic eyes	(Loso <i>et al.</i> , 2010) (Srivastava <i>et al.</i> , 2009)
Resveratrol (<i>trans</i> -3, 5, 4'-trihydroxystilbene)	Not known		(Sharma <i>et al.</i> , 1992)
<i>Saptamrita lauha</i>	Flavanoid-like compounds but not specified		

mechanisms of DR in TCMs are *yin* vacuity with dryness and heat leading to accumulation of phlegm and blood stasis, and causing not nourishing the eyes and static blood obstructing the network vessels of the eyes (Flaws *et al.*, 2002). Many clinical studies have shown the benefit of TCMs in reducing symptoms, improving visual acuity and visual field and improving the outcome under ophthalmoscopy (Ma *et al.*, 2004; Song *et al.*, 2006). Moreover, a growing body of evidence has shown the benefit of combining TCMs with Western treatment strategies (Donnelly *et al.*, 2006). Indeed, a combination of laser therapy with TCMs has been explored for DR in China (Gong, 2007). Traditionally, 'blood-vitalizing herbs' are used to treat microvascular-related DR (Han and Xu, 1988). Extracts used in TCMs provide therapeutic effects that are dependent on its related chemical compounds (Shi *et al.*, 2008). Despite the growing expenditure of TCMs for DR management, there is still no conclusive evidence to support its efficacy and issues of safety (Yeh *et al.*, 2003; Liu *et al.*, 2004; West *et al.*, 2006). Thus, the study of active compounds from the extract and its possible applicable mechanisms are required for better understanding the effect of TCMs in DR (Su *et al.*, 2007).

One human supplementation trial showed that the *Lycium (L.) barbarum* (Goji berry) intake increases the fasting plasma zeaxanthin levels, beneficial for maintaining macular pigment density in age-related macular degeneration (Cheng *et al.*, 2005). *L. barbarum* polysaccharides, most abundantly present in Goji berry, induce an activation of microglia (brain macrophages) in retina (Chiu *et al.*, 2009). Indeed, Goji berry has been shown to be effective in glaucoma and in modulating immune response in retinal ganglion cells in a rat hypertension model (Chan *et al.*, 2007; Chiu *et al.*, 2009). Moreover, LPB has been shown to antagonize glutamate excitotoxicity in rat cortical neurons (Ho *et al.*, 2009). In addition, *L. barbarum* polysaccharides has been shown to increase anti-apoptotic protein Bcl-2 level in lens epithelial cells of the whole lens incubated in culture medium and exposed to hydrogen peroxide (Wang *et al.*, 2002).

Clinical study of DR treated with compound Danshen dripping pills (*Salviae miltiorrhiae*, Radix notoginseng and borneol) showed improvement in visual acuity, and control of micro-haemorrhage and microaneurysm of fundus. Moreover, the pills were effective in visual function recovery of DR (Qi *et al.*, 2007). Another study on the compound Danshen dripping pill was shown to improve ischaemia of the retina and the visual field in early DR, and to reduce the number of microaneurysms (Deng *et al.*, 2005).

Anisodamine is an alkaloid, originally isolated from *Anisodus tanguticus* which was found to increase neural tissue oxygen tension of the retina (Linsenmeier *et al.*, 1989). Moreover, anisodamine has been shown to increase blood flow and oxygen delivery to the retina-choroid and iris-ciliary body of the eye, and to decrease lipid peroxidation in the retinal cells (Zhang *et al.*, 1990).

The ethanol extract of *Radix (R.) Puerariae* has shown to inhibit glycosylation of rat lens protein (Duan *et al.*, 2000). Moreover, a new key chemical constituent 7-(6-O-malonyl- β -D-glucopyranosyloxy)-3-(4-hydroxyphenyl)-4H-1benzopyran-4-one was isolated from *R. Puerariae*. This new aldose reductase-inhibitory compound proved to be useful in DR (Hirakura *et al.*, 1989). The active components astragalo-

sides from crude extract of *Astragalus (A.) Radix* has also been shown to be potentially useful for the prevention of clinical diabetic complications such as DR by inhibiting AGEs production (Motomura *et al.*, 2009).

A study of therapeutic efficacy of the root extracts of *Stephania tetrandra* S. Moore for delaying the progression of retinopathy was carried out in the streptozotocin-induced diabetic rat model. It was considered to work through the activation by its active component, tetrandrine, of suppression of neovascularization of the retinal capillary (Liang *et al.*, 2002).

Traditional Chinese formula Hachimi-Jio-gan (Trhmannia Eight Formula) controls the balance of sodium, potassium and calcium ions for maintenance of lens transparency. This finding leads to the possibility of prophylactic use of Hachimi-Jio-gan in eye diseases such as diabetic cataract (Kamei *et al.*, 1987). However, further study is warranted for human application.

Ayurvedic herbal medicines and diabetic retinopathy

Ayurveda has been widely used in India and some European countries, and recognition of its modality is increasing worldwide (Saxena and Vikram, 2004). More than 800 plants are used as traditional remedies for diabetes and its complications in the ancient Indian Ayurvedic and Unani system of medicine (Alarcon-Aguilara *et al.*, 1998; Saxena and Vikram, 2004). However, only a few medicinal herbs have been evaluated by scientific investigations (Saxena and Vikram, 2004).

A polyol, conduritol A from *Gymnema sylvestre* may be responsible for the cataract-suppressing effect by inhibiting lens aldose reductase (Miyatake *et al.*, 1994).

The seeds of *Trigonella foenum graecum*, commonly known as fenugreek, have been known for hypoglycaemic, hypocholesterolemic and hyperinsulinemic effects in diabetic patients and experimental animals (Khosla *et al.*, 1995; Puri *et al.*, 2002). It has been shown that fenugreek has a beneficial effect on abnormal accumulation of sorbitol and fructose in the diabetic lens. Moreover, disintegration of the inner nuclear layer cells with reduction in rough endoplasmic reticulum and swelling of mitochondria in the diabetic retina have been restored by fenugreek. Saponins, flavanoids and 4-hydroxyisoleucine are potential plausible effectors in reversing the histopathological and biochemical abnormalities observed in DR and warrants further investigation (Preet *et al.*, 2006). Clinical study has shown that an indigenous drug, *Saptamrita lauha* may have a role in control of retinal haemorrhage absorption and its recurrence in retinopathic eyes (Sharma *et al.*, 1992).

Western herbal medicines and diabetic retinopathy

Western herbal medicine is the most widely used form of herbal medicine although Ayurvedic and Chinese herbal medicines are becoming better known (Wohlmuth *et al.*, 2002).

The fruits of *Vaccinium (V.) myrtillus* (billberry), containing 25% anthocyanidins, have been widely used in Europe to prevent and treat DR (Murray, 1997; Camire, 2000). In a clinical study, 12 adult diabetic patients were treated with 600 mg of anthocyanosides daily for 2 months. Following

Table 2

Modern research on herbal and traditional medicines affecting PPAR- γ activation and its possible mechanisms in diabetic retinopathy

PPAR- γ activity in herbal medicine	Active component	Functions in diabetic retinopathy	References
<i>Astragalus membranaceus</i> and <i>Pueraria thomsonii</i>	Biochanin A	Inhibitory effect on protein glycosylation in diabetic retinopathy	(Asgary <i>et al.</i> , 2002)
<i>Swietenia mahagony</i>	Not known	Antioxidant activity of <i>Swietenia mahagony</i> seed may provide a new avenue against diabetic retinopathy	(Rodrigues <i>et al.</i> , 2010)
Korean red ginseng	Ginsenoside 20(S) Ginsenoside Rb2 Ginsenoside Re	Prevention of high glucose-induced apoptosis through decreased in Bcl-2 expression and increased Bax expression Protective effect against oxidative stress in the eye of diabetic rats which can be effective in the prevention of diabetic microvasculopathy	(Park, 2008) (Cho <i>et al.</i> , 2006)
Dan-shao-hua-xian formula (<i>Astragali Radix</i>) (<i>Stephania tetrandra</i> S. Moore)	Astragalosides Tetrandrine	Beneficial for the prevention of diabetic retinopathy by inhibiting AGEs formation Suppression of neovascularization of retinopathy in diabetes	(Wang and Cheng, 2008) (Motomura <i>et al.</i> , 2009) (Liang <i>et al.</i> , 2002)
Turmeric	Curcumin Curcuminoids	Inhibition of VEGF level in diabetes-induced retina Antioxidation property by inhibition of Ca ²⁺ entry and protein kinase activity	(Zhou <i>et al.</i> , 2007) (Kowluru and Kanwar, 2007; Mrudula <i>et al.</i> , 2007) (Balasubramanyam <i>et al.</i> , 2003)

treatment, there was a significant decrease in the biosynthesis of connective tissue. The authors interpreted these results to mean that anthocyanosides have a protective property against DR (Boniface and Robert, 1996). In another study, treatment of 31 patients suffering from various types of retinopathy with *V. myrtillus* extract was associated with a reduced tendency towards retinal haemorrhage (Perossini *et al.*, 1988). Resveratrol (*trans*-3, 5, 4'-trihydroxystilbene) is a phytoalexin, particularly rich in the skin of red grapes, which has known benefit for cancer, DR, rheumatoid arthritis and cardiovascular disorders (Srivastava *et al.*, 2009; Losso *et al.*, 2010). In particular, resveratrol has been shown to protect high glucose-induced ARPE-19 cells by reducing low-grade inflammatory mediators, interleukin-6 and interleukin-8, COX-2, VEGF and connexin 43 down-regulation (Losso *et al.*, 2010).

One study has shown the beneficial outcome of the treatment of DR with laser photocoagulation therapy when combined with *Ginkgo* (*G.*) *dipyrifolium* injection than the laser therapy alone in human clinical trials (Wei *et al.*, 2005). In addition, the therapeutic efficiency of the *G. biloba* extract was estimated for early DR, associated with diabetic dyschromatopsia in a double-blind trial. An improvement tendency was evidenced by *G. biloba* treatment without retinal ischemia (Lanthony and Cosson, 1988).

Herbal and traditional medicines that modulate diabetic retinopathy through PPAR- γ activation

Over the last two decades, herbal plants or their active components have been major sources of bioactive agents and continue to represent a potential 'goldmine' for novel ligands

to be discovered (Egan, 2002; Shen *et al.*, 2003). However, the mechanisms of action for most of the herbal medicines associated with modulating DR have not been fully elucidated. Nevertheless, the experience obtained from their traditional use over many years should not be ignored (Elvin-Lewis, 2001). Despite numerous plant-derived PPAR- γ activators having been identified from plants in recent years (Huang *et al.*, 2009), there are no conclusive studies on DR and its mechanism of modulation through the PPAR- γ activation. The next section of this review summarizes the current studies on herbal or traditional medicine associated with PPAR- γ activation and the possible mechanisms relevant to the management of DR (Table 2).

Biochanin A is a common isoflavone from *Astragalus membranaceus* and *Pueraria thomsonii* which has been shown to induce adipocyte differentiation activation *in vitro* through or at least in part, PPAR- γ , as measured by reporter-gene activity. This finding has suggested the potential value of Biochanin A as an anti-diabetic agent (Shen *et al.*, 2006). Moreover, biochanin A has shown beneficial effect in DR by inhibition of protein glycosylation (Asgary *et al.*, 2002). *Swietenia mahagony* is native to the West Indies, and is traditionally used for the treatment of hypertension and malaria as a folk medicine in Indonesia (Nagalakshmi *et al.*, 2001). This medicinal herb has shown anti-diabetic activity with PPAR- γ transcriptional regulatory function as one of its *in vivo* mechanisms (Li *et al.*, 2005). Moreover, antioxidant activity of *Swietenia mahagony* seed may provide a new approach to diabetic-related complications (Rodrigues *et al.*, 2010).

One study has evaluated the effect of three traditional anti-diabetic herbs on the progression of diabetes. The results showed that mulberry leaf, Korean red ginseng and banaba leaf extracts significantly increased insulin sensitivity and

improved hyperglycaemia through the PPAR- γ pathway (Park *et al.*, 2005). In particular, ginsenoside 20(S), one of the ginsenoside metabolites, was found to increase PPAR- γ -transactivation activity and expression of glucose transporter 4 (GLUT 4) in 3T3-L1 adipocytes. Moreover, ginsenoside Rb2, an active component from Korean red ginseng, prevents high glucose-induced apoptosis through an increase in Bcl-2 expression and decrease in Bax expression in ARPE-19 cells (Park, 2008). Another active component, ginsenoside Re also has shown a protective effect against oxidative stress in the eye of diabetic rats, which is effective in the prevention of diabetic microvasculopathy (Cho *et al.*, 2006). Dan-shao-hua-xian, a TCM preparation, has been shown in a recent article to play a role in treating hepatic fibrosis by up-regulation of the gene expression and generation of PPAR- γ , and also by suppressing TNF- α production through down-regulation of the NF- κ B signalling pathway (Wang and Cheng, 2008). Moreover, *Stephania tetrandra* S. Moore, one of the ingredients in Dan-shao-hua-xian, has been shown to suppress neovascularization of retinopathy in diabetes through the activation of a major chemical component, tetrandrine (Liang *et al.*, 2002). In addition, one of the other ingredient of Dan-shao-hua-xian, *A. Radix*, and its active components astragalosides, have been shown to be beneficial for the prevention of DR by inhibiting AGEs formation (Motomura *et al.*, 2009). Curcumin, the phytochemical component in turmeric has been shown to interrupt the platelet-derived growth factor and epidermal growth factor signalling pathway by stimulation of PPAR- γ gene expression in rat activated hepatic stellate cell (Zhou *et al.*, 2007). Moreover, a recent review has highlighted the importance of the anti-inflammatory property of curcumin component from turmeric by suppression of TNF- α expression and release, which is potentially mediated by the up-regulation of PPAR- γ (Jacob *et al.*, 2007). The anti-inflammatory property of curcumin has also been shown to inhibit VEGF level in streptozotocin-induced diabetic rat retina which gives a potential role in DR (Kowluru and Kanwar, 2007; Mrudula *et al.*, 2007). Curcuminoids in turmeric have shown antioxidant properties by inhibition of calcium entry and protein kinase activity, which may have therapeutic implications in DR (Balasubramanyam *et al.*, 2003).

Terpenoids are one of many bioactive phytochemicals in medicinal plants, which include carotenoids, isoprenoids and isoprenols (He *et al.*, 1997). Studies have shown that terpenoids are significant in cancer and cardiovascular diseases (Takahashi *et al.*, 2002). On the other hand, typical isoprenols from herbs and fruits, including farnesol and geranylgeraniol, activate PPAR- γ in 3T3-L1 adipocytes and HepG2 hepatocytes. In addition, these herbal derivatives up-regulate the expression of some lipid metabolic target genes of PPAR- γ , including an adipocyte-specific gene, *aP2* (He *et al.*, 1997; Takahashi *et al.*, 2002). Moreover, terpenoids such as ginkgolides from *G. biloba* improve blood flow by dilating blood vessels and reducing the stickiness of platelets for eye disorders, including DR (Braquet *et al.*, 1985; Baudouin *et al.*, 1999).

Conclusion

Diabetic retinopathy remains one of the major risk factors and a leading cause of preventable blindness worldwide. The

increasing impotence of understanding the specific molecular and biochemical changes in DR leads to the requirement for development of novel therapeutic interventions. Although it is an important cause of blindness, initially DR presents few visual or ophthalmic symptoms until complete visual loss occurs (Fong *et al.*, 2004a,b). Current treatments of DR rarely improve visual function and are limited to surgical options in an advanced stage, with excessive side effects and significant financial burden. Hence, emerging treatments, possibly in combination with standard therapy, may provide superior efficacy and safety profile for the treatment or prevention of DR. Moreover, the new strategies move to a paradigm in treating the early stages of DR. The recent advancements in the knowledge of the pathogenic alterations driving ocular damage and vision loss in DR strongly focus on PPAR- γ as a valuable target to control high glucose-induced inflammation and apoptosis. PPAR- γ functions as a transcription factor and thereby controls cellular processes at the level of gene expression, through modulation by its nuclear receptor activity of selective downstream gene expression (Huang *et al.*, 2005). The complexity of PPAR- γ activation not only provides beneficial effects but also introduces risks from undesirable side effects, such as cardiovascular complications with long-term application (Roehr, 2010). PPAR- γ is an attractive and relatively unexploited target for herbal-derived medicines in DR. This review confirms that natural and traditional herbal medicines have potential as alternative or combination therapy for DR, and emphasizes their possible mechanisms through PPAR- γ activity. Despite the long history of herbal and natural traditional medicines for the management of DR, there is still no conclusive evidence on their effectiveness and safety profiles. Therefore, future studies are warranted for extensive investigation to gather proof of efficacy in various preclinical and clinical settings.

Acknowledgements

This work was supported by a grant from the National Institute of Complementary Medicine (NICM) to B. D. R.

Conflict of interest

The authors declare no conflict of interest.

References

- Abu El-Asrar AM, Al-Mezaine HS, Ola MS (2009). Pathophysiology and management of diabetic retinopathy. *Expert Review of Ophthalmology* 4: 627–647.
- Adams AP (2002). Is diabetic retinopathy an inflammatory disease? *Br J Ophthalmol* 86: 363–365.
- Ahmed N (2005). Advanced glycation endproducts–role in pathology of diabetic complications. *Diabetes Res Clin Pract* 67: 3–21.

- Al-Rowais NA (2002). Herbal medicine in the treatment of diabetes mellitus. *Saudi Med J* 23: 1327–1331.
- Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CC, Flores-Saenz JL *et al.* (1998). Study of the anti-hyperglycemic effect of plants used as antidiabetics. *J Ethnopharmacol* 61: 101–110.
- Allen DA, Yaqoob MM, Harwood SM (2005). Mechanisms of high glucose-induced apoptosis and its relationship to diabetic complications. *J Nutr Biochem* 16: 705–713.
- Asgary S, Naderi GA, Zadegan NS, Vakili R (2002). The inhibitory effects of pure flavonoids on *in vitro* protein glycosylation. *J Herb Pharmacother* 2: 47–55.
- Baeuerle PA, Henkel T (1994). Function and activation of NF-kappa B in the immune system. *Annu Rev Immunol* 12: 141–179.
- Bailey CJ, Day C (1989). Traditional plant medicines as treatments for diabetes. *Diabetes Care* 12: 553–564.
- Baker MS, Chen X, Cao XC, Kaufman DB (2001). Expression of a dominant negative inhibitor of NF-kappaB protects MIN6 beta-cells from cytokine-induced apoptosis. *J Surg Res* 97: 117–122.
- Balasubramanyam M, Koteswari AA, Kumar RS, Monickaraj SF, Maheswari JU, Mohan V *et al.* (2003). Curcumin-induced inhibition of cellular reactive oxygen species generation: novel therapeutic implications. *J Biosci* 28: 715–721.
- Barnett AH, Grant PJ, Hitman GA, Mather H, Pawa M, Robertson L *et al.* (2003). Rosiglitazone in Type 2 diabetes mellitus: an evaluation in British Indo-Asian patients. *Diabet Med* 20: 387–393.
- Baudouin C, Pisella PJ, Ettaiche M, Goldschild M, Becquet F, Gstaad P *et al.* (1999). Effects of EGB761 and superoxide dismutase in an experimental model of retinopathy generated by intravitreal production of superoxide anion radical. *Graefes Arch Clin Exp Ophthalmol* 237: 58–66.
- Beatty S, Koh H, Phil M, Henson D, Boulton M (2000). The role of oxidative stress in the pathogenesis of age-related macular degeneration. *Surv Ophthalmol* 45: 115–134.
- Beck L Jr, D'Amore PA (1997). Vascular development: cellular and molecular regulation. *FASEB J* 11: 365–373.
- Blackwell TS, Christman JW (1997). The role of nuclear factor-kappa B in cytokine gene regulation. *Am J Respir Cell Mol Biol* 17: 3–9.
- Bloomgarden ZT (2007). Screening for and managing diabetic retinopathy: current approaches. *Am J Health Syst Pharm* 64: S8–14.
- Boileau TW, Bray TM, Bomser JA (2003). Ultraviolet radiation modulates nuclear factor kappa B activation in human lens epithelial cells. *J Biochem Mol Toxicol* 17: 108–113.
- Bok D (1993). The retinal pigment epithelium: a versatile partner in vision. *J Cell Sci Suppl* 17: 189–195.
- Boniface R, Robert AM (1996). [Effect of anthocyanins on human connective tissue metabolism in the human]. *Klin Monbl Augenheilkd* 209: 368–372.
- Bonne C (2005). [PPAR gamma: a novel pharmacological target against retinal and choroidal neovascularization]. *J Fr Ophtalmol* 28: 326–330.
- Braquet P, Etienne A, Touvy C, Bourgain RH, Lefort J, Vargaftig BB *et al.* (1985). Involvement of platelet activating factor in respiratory anaphylaxis, demonstrated by PAF-acether inhibitor BN 52021. *Lancet* 1: 1501.
- Brownlee M (2005). The pathobiology of diabetic complications: a unifying mechanism. *Diabetes* 54: 1615–1625.
- Camire ME (2000). Bilberries and blueberries as functional foods and nutraceuticals. In: Oomah BD (ed.). *Herbs, Botanicals & Teas*. Technomic Publishing Company: Lancaster, PA, pp. 289–319.
- Carmo A, Cunha-Vaz JG, Carvalho AP, Lopes MC (1999). L-arginine transport in retinas from streptozotocin diabetic rats: correlation with the level of IL-1 beta and NO synthase activity. *Vision Res* 39: 3817–3823.
- Ceriello A, Mercuri F, Quagliaro L, Assaloni R, Motz E, Tonutti L *et al.* (2001). Detection of nitrotyrosine in the diabetic plasma: evidence of oxidative stress. *Diabetologia* 44: 834–838.
- Ceylan-Isik AF, Fliethman RM, Wold LE, Ren J (2008). Herbal and traditional Chinese medicine for the treatment of cardiovascular complications in diabetes mellitus. *Curr Diabetes Rev* 4: 320–328.
- Chan HC, Chang RC, Koon-Ching Ip A, Chiu K, Yuen WH, Zee SY *et al.* (2007). Neuroprotective effects of Lycium barbarum Lynn on protecting retinal ganglion cells in an ocular hypertension model of glaucoma. *Exp Neurol* 203: 269–273.
- Chang JY, Bora PS, Bora NS (2008). Prevention of Oxidative Stress-Induced Retinal Pigment Epithelial Cell Death by the PPARgamma Agonists, 15-Deoxy-Delta 12, 14-Prostaglandin J(2). *PPAR Res* 2008: 720163.
- Chang PC, Chen TH, Chang CJ, Hou CC, Chan P, Lee HM *et al.* (2004). Advanced glycosylation end products induce inducible nitric oxide synthase (iNOS) expression via a p38 MAPK-dependent pathway. *Kidney Int* 65: 1664–1675.
- Chang RC, So KF (2008). Use of anti-aging herbal medicine, Lycium barbarum, against aging-associated diseases. What do we know so far? *Cell Mol Neurobiol* 28: 643–652.
- Cheng CY, Chung WY, Szeto YT, Benzie IF (2005). Fasting plasma zeaxanthin response to Fructus barbarum L. (wolfberry; Kei Tze) in a food-based human supplementation trial. *Br J Nutr* 93: 123–130.
- Cheng HC, Ho TC, Chen SL, Lai HY, Hong KF, Tsao YP (2008). Troglitazone suppresses transforming growth factor beta-mediated fibrogenesis in retinal pigment epithelial cells. *Mol Vis* 14: 95–104.
- Chiu K, Chan HC, Yeung SC, Yuen WH, Zee SY, Chang RC *et al.* (2009). Modulation of microglia by Wolfberry on the survival of retinal ganglion cells in a rat ocular hypertension model. *J Ocul Biol Dis Infor* 2: 47–56.
- Cho WC, Chung WS, Lee SK, Leung AW, Cheng CH, Yue KK *et al.* (2006). Ginsenoside Re of Panax ginseng possesses significant antioxidant and antihyperlipidemic efficacies in streptozotocin-induced diabetic rats. *Eur J Pharmacol* 550: 173–179.
- Chu J, Ali Y (2008). Diabetic retinopathy: a review. *Drug Dev Res* 69: 1–14.
- Ciulla TA, Amador AG, Zinman B (2003). Diabetic retinopathy and diabetic macular edema: pathophysiology, screening, and novel therapies. *Diabetes Care* 26: 2653–2664.
- Covington M (2001). Traditional Chinese medicine in the treatment of diabetes. *Diabetes Spectrum* 14: 154–159.
- Danial NN, Korsmeyer SJ (2004). Cell death: critical control points. *Cell* 116: 205–219.
- De Luca A, Carotenuto A, Rachiglio A, Gallo M, Maiello MR, Aldinucci D *et al.* (2008). The role of the EGFR signaling in tumor microenvironment. *J Cell Physiol* 214: 559–567.

- Decanini A, Karunadharma PR, Nordgaard CL, Feng X, Olsen TW, Ferrington DA (2008). Human retinal pigment epithelium proteome changes in early diabetes. *Diabetologia* 51: 1051–1061.
- Deng H, Jin M, Yuan W, Xiang Z, Wang J (2005). Clinical observation treating early diabetic retinopathy with compound danshen dripping pills. *Zhongguo Zhong Yi Yan Ke Za Zhi [J Tradit Chin Ophthalmol]* 15: 72–74.
- Denis U, Lecomte M, Paget C, Ruggiero D, Wiernsperger N, Lagarde M *et al.* (2002). Advanced glycation end-products induce apoptosis of bovine retinal pericytes in culture: involvement of diacylglycerol/ceramide production and oxidative stress induction. *Free Radic Biol Med* 33: 236–247.
- Dey L, Attele AS, Yuan CS (2002). Alternative therapies for type 2 diabetes. *Altern Med Rev* 7: 45–58.
- Dolhofer-Bliesener R, Lechner B, Gerbitz KD (1996). Possible significance of advanced glycation end products in serum in end-stage renal disease and in late complications of diabetes. *Eur J Clin Chem Clin Biochem* 34: 355–361.
- Donnelly R, Wang B, Qu X (2006). Type 2 diabetes in China: partnerships in education and research to evaluate new antidiabetic treatments. *Br J Clin Pharmacol* 61: 702–705.
- Doonan F, Wallace DM, O'Driscoll C, Cotter TG (2009). Rosiglitazone acts as a neuroprotectant in retinal cells via up-regulation of sestrin-1 and SOD-2. *J Neurochem* 109: 631–643.
- Du X, Stocklauser-Farber K, Rosen P (1999). Generation of reactive oxygen intermediates, activation of NF-kappaB, and induction of apoptosis in human endothelial cells by glucose: role of nitric oxide synthase? *Free Radic Biol Med* 27: 752–763.
- Duan YJ, Wang S.Y., Wang X.Y., Liu Y., Gao G.L. (2000). Inhibitory effects of Radix Puerariae on nonenzymic glycosylation of proteins. *Chin J Diabetes* 17: 61–62, 78.
- Egan CD (2002). Addressing use of herbal medicine in the primary care setting. *J Am Acad Nurse Pract* 14: 166–171.
- Elvin-Lewis M (2001). Should we be concerned about herbal remedies. *J Ethnopharmacol* 75: 141–164.
- Ershov AV, Bazan NG (2000). Photoreceptor phagocytosis selectively activates PPARgamma expression in retinal pigment epithelial cells. *J Neurosci Res* 60: 328–337.
- Fehlberg S, Gregel CM, Goke A, Goke R (2003). Bisphenol A diglycidyl ether-induced apoptosis involves Bax/Bid-dependent mitochondrial release of apoptosis-inducing factor (AIF), cytochrome c and Smac/DIABLO. *Br J Pharmacol* 139: 495–500.
- Flaws B, Kuchinski L, Casañas R (2002). Diabetic retinopathy. In: Brearton EJ (ed.). *The Treatment of Diabetes Mellitus with Chinese Medicine*. A Division of Blue Poppy Enterprises Inc.: Boulder, CO, pp. 155–168.
- Fong DS, Aiello L, Gardner TW, King GL, Blankenship G, Cavallerano JD *et al.* (2003). Diabetic retinopathy. *Diabetes Care* 26: 226–229.
- Fong DS, Aiello L, Gardner TW, King GL, Blankenship G, Cavallerano JD *et al.* (2004a). Retinopathy in diabetes. *Diabetes Care* 27 (Suppl. 1): S84–S87.
- Fong DS, Aiello LP, Ferris FL 3rd, Klein R (2004b). Diabetic retinopathy. *Diabetes Care* 27: 2540–2553.
- Fuenzalida K, Quintanilla R, Ramos P, Piderit D, Fuentealba RA, Martinez G *et al.* (2007). Peroxisome proliferator-activated receptor gamma up-regulates the Bcl-2 anti-apoptotic protein in neurons and induces mitochondrial stabilization and protection against oxidative stress and apoptosis. *J Biol Chem* 282: 37006–37015.
- Garcia C, Aranda J, Arnold E, Thebault S, Macotela Y, Lopez-Casillas F *et al.* (2008). Vasoinhibins prevent retinal vasopermeability associated with diabetic retinopathy in rats via protein phosphatase 2A-dependent eNOS inactivation. *J Clin Invest* 118: 2291–2300.
- Gardner TW, Antonetti DA, Barber AJ, LaNoue KF, Nakamura M (2000). New insights into the pathophysiology of diabetic retinopathy: potential cell-specific therapeutic targets. *Diabetes Technol Ther* 2: 601–608.
- Garg TK, Chang JY (2003). Oxidative stress causes ERK phosphorylation and cell death in cultured retinal pigment epithelium: prevention of cell death by AG126 and 15-deoxy-delta 12, 14-PGJ2. *BMC Ophthalmol* 3: 5.
- Garg TK, Chang JY (2004). 15-deoxy-delta 12, 14-Prostaglandin J2 prevents reactive oxygen species generation and mitochondrial membrane depolarization induced by oxidative stress. *BMC Pharmacol* 4: 6.
- Gately S, Li WW (2004). Multiple roles of COX-2 in tumor angiogenesis: a target for antiangiogenic therapy. *Semin Oncol* 31: 2–11.
- Geiger RC, Waters CM, Kamp DW, Glucksberg MR (2005). KGF prevents oxygen-mediated damage in ARPE-19 cells. *Invest Ophthalmol Vis Sci* 46: 3435–3442.
- Gerry JM, Pascual G (2008). Narrowing in on cardiovascular disease: the atheroprotective role of peroxisome proliferator-activated receptor gamma. *Trends Cardiovasc Med* 18: 39–44.
- Giaginis C, Tsourouflis G, Theocharis S (2008). Peroxisome proliferator-activated receptor-gamma (PPAR-gamma) ligands: novel pharmacological agents in the treatment of ischemia reperfusion injury. *Curr Mol Med* 8: 562–579.
- Girnun GD, Domann FE, Moore SA, Robbins ME (2002). Identification of a functional peroxisome proliferator-activated receptor response element in the rat catalase promoter. *Mol Endocrinol* 16: 2793–2801.
- Goh SY, Cooper ME (2008). Clinical review: the role of advanced glycation end products in progression and complications of diabetes. *J Clin Endocrinol Metab* 93: 1143–1152.
- Gong H (2007). Ji guang guang ning lian he hu wang ming mu tang zhi liao tang niao bing shi wang mo bing bian de liao xiao guan cha. *Zhong Wai Jian Kang Wen Zhang [World Health Digest Med Period]* 4: 55–56.
- Grey ST, Arvelo MB, Hasenkamp W, Bach FH, Ferran C (1999). A20 inhibits cytokine-induced apoptosis and nuclear factor kappaB-dependent gene activation in islets. *J Exp Med* 190: 1135–1146.
- Grossniklaus HE, Ling JX, Wallace TM, Dithmar S, Lawson DH, Cohen C *et al.* (2002). Macrophage and retinal pigment epithelium expression of angiogenic cytokines in choroidal neovascularization. *Mol Vis* 8: 119–126.
- Hammes HP (2005). Pericytes and the pathogenesis of diabetic retinopathy. *Horm Metab Res* 37 (Suppl. 1): 39–43.
- Hammes HP, Strodter D, Weiss A, Bretzel RG, Federlin K, Brownlee M *et al.* (1995). Secondary intervention with aminoguanidine retards the progression of diabetic retinopathy in the rat model. *Diabetologia* 38: 656–660.

- Hammes HP, Hoerauf H, Alt A, Schleicher E, Clausen JT, Bretzel RG *et al.* (1999). N(epsilon)(carboxymethyl)lysine and the AGE receptor RAGE colocalize in age-related macular degeneration. *Invest Ophthalmol Vis Sci* 40: 1855–1859.
- Han D, Xu R (1988). Progress in the research of blood activation and hemostasis removal. *Abstr Chin Med* 2: 466–483.
- Haubner F, Lehle K, Munzel D, Schmid C, Birnbaum DE, Preuner JG *et al.* (2007). Hyperglycemia increases the levels of vascular cellular adhesion molecule-1 and monocyte-chemoattractant-protein-1 in the diabetic endothelial cell. *Biochem Biophys Res Commun* 360: 560–565.
- He L, Mo H, Hadisusilo S, Qureshi AA, Elson CE (1997). Isoprenoids suppress the growth of murine B16 melanomas *in vitro* and *in vivo*. *J Nutr* 127: 668–674.
- Head KA (1999). Natural therapies for ocular disorders, part one: diseases of the retina. *Altern Med Rev* 4: 342–359.
- Henkind P (1978). Ocular neovascularization. The Krill memorial lecture. *Am J Ophthalmol* 85: 287–301.
- Herzlich AA, Tuo J, Chan CC (2008). Peroxisome proliferator-activated receptor and age-related macular degeneration. *PPAR Res* 2008: 389507.
- Hirakura K, Nakajima K, Sato S, Mihashi H (1989). Isolation of 7-(6-O-Malonyl-β-D-glucopyranosyloxy)-3-(4-hydroxyphenyl)-4H-1benzopyran-4-one from *Pueraria lobata* Ohwi as aldose reductase inhibitors and pharmaceutical formulations. Japan Kokai Tokkyo Koho (Patent). Patent Number: JP 01146894.
- Ho YS, Yu MS, Yang XF, So KF, Yuen WH, Chang RC *et al.* (2009). Neuroprotective effects of polysaccharides from wolfberry, the fruits of *Lycium barbarum*, against homocysteine-induced toxicity in rat cortical neurons. *J Alzheimers Dis* 19: 813–827.
- Hollborn M, Stathopoulos C, Steffen A, Wiedemann P, Kohen L, Bringmann A *et al.* (2007). Positive feedback regulation between MMP-9 and VEGF in human RPE cells. *Invest Ophthalmol Vis Sci* 48: 4360–4367.
- Holtkamp M, Zschenderlein R, Bruck W, Weber JR (2001). Chronic inflammatory demyelinating polyradiculoneuropathy with histologically proven optic neuritis. *Acta Neuropathol* 101: 529–531.
- Huang TH, Kota BP, Razmovski V, Roufogalis BD (2005). Herbal or natural medicines as modulators of peroxisome proliferator-activated receptors and related nuclear receptors for therapy of metabolic syndrome. *Basic Clin Pharmacol Toxicol* 96: 3–14.
- Huang TH, Teoh AW, Lin BL, Lin DS, Roufogalis B (2009). The role of herbal PPAR modulators in the treatment of cardiometabolic syndrome. *Pharmacol Res* 60: 195–206.
- Jacob A, Wu R, Zhou M, Wang P (2007). Mechanism of the anti-inflammatory effect of curcumin: PPAR-gamma activation. *PPAR Res* 2007: 89369.
- de Jong PT (2006). Age-related macular degeneration. *N Engl J Med* 355: 1474–1485.
- Joussen AM, Poulaki V, Mitsiades N, Kirchhof B, Koizumi K, Dohmen S *et al.* (2002). Nonsteroidal anti-inflammatory drugs prevent early diabetic retinopathy via TNF-alpha suppression. *FASEB J* 16: 438–440.
- Joussen AM, Poulaki V, Le ML, Koizumi K, Esser C, Janicki H *et al.* (2004). A central role for inflammation in the pathogenesis of diabetic retinopathy. *FASEB J* 18: 1450–1452.
- Kaji Y, Amano S, Usui T, Oshika T, Yamashiro K, Ishida S *et al.* (2003). Expression and function of receptors for advanced glycation end products in bovine corneal endothelial cells. *Invest Ophthalmol Vis Sci* 44: 521–528.
- Kamei A, Hisada T, Iwata S (1987). The evaluation of therapeutic efficacy of hachimi-jio-gan (traditional Chinese medicine) to rat galactosemic cataract. *J Ocul Pharmacol* 3: 239–248.
- Kasper M, Roehlecke C, Witt M, Fehrenbach H, Hofer A, Miyata T *et al.* (2000). Induction of apoptosis by glyoxal in human embryonic lung epithelial cell line L132. *Am J Respir Cell Mol Biol* 23: 485–491.
- Kennedy CJ, Rakoczy PE, Constable IJ (1995). Lipofuscin of the retinal pigment epithelium: a review. *Eye* 9 (Pt 6): 763–771.
- Kern TS (2007). Contributions of inflammatory processes to the development of the early stages of diabetic retinopathy. *Exp Diabetes Res* 2007: 95103.
- Kern TS, Tang J, Mizutani M, Kowluru RA, Nagaraj RH, Romeo G *et al.* (2000). Response of capillary cell death to aminoguanidine predicts the development of retinopathy: comparison of diabetes and galactosemia. *Invest Ophthalmol Vis Sci* 41: 3972–3978.
- Khosla P, Gupta DD, Nagpal RK (1995). Effect of *Trigonella foenum graecum* (Fenugreek) on blood glucose in normal and diabetic rats. *Indian J Physiol Pharmacol* 39: 173–174.
- Kim EK, Kwon KB, Koo BS, Han MJ, Song MY, Song EK *et al.* (2007). Activation of peroxisome proliferator-activated receptor-gamma protects pancreatic beta-cells from cytokine-induced cytotoxicity via NF kappaB pathway. *Int J Biochem Cell Biol* 39: 1260–1275.
- Kimoto K, Nakatsuka K, Matsuo N, Yoshioka H (2004). p38 MAPK mediates the expression of type I collagen induced by TGF-beta 2 in human retinal pigment epithelial cells ARPE-19. *Invest Ophthalmol Vis Sci* 45: 2431–2437.
- Kliffen M, Sharma HS, Mooy CM, Kerkvliet S, de Jong PT (1997). Increased expression of angiogenic growth factors in age-related maculopathy. *Br J Ophthalmol* 81: 154–162.
- Knudsen ST, Bek T, Poulsen PL, Hove MN, Rehling M, Mogensen CE *et al.* (2002). Macular edema reflects generalized vascular hyperpermeability in type 2 diabetic patients with retinopathy. *Diabetes Care* 25: 2328–2334.
- Kowluru RA (2005). Diabetic retinopathy: mitochondrial dysfunction and retinal capillary cell death. *Antioxid Redox Signal* 7: 1581–1587.
- Kowluru RA, Kanwar M (2007). Effects of curcumin on retinal oxidative stress and inflammation in diabetes. *Nutr Metab (Lond)* 4: 8.
- Kowluru RA, Odenbach S (2004). Effect of long-term administration of alpha-lipoic acid on retinal capillary cell death and the development of retinopathy in diabetic rats. *Diabetes* 53: 3233–3238.
- Kowluru RA, Kern TS, Engerman RL, Armstrong D (1996). Abnormalities of retinal metabolism in diabetes or experimental galactosemia. III. Effects of antioxidants. *Diabetes* 45: 1233–1237.
- Kowluru RA, Tang J, Kern TS (2001). Abnormalities of retinal metabolism in diabetes and experimental galactosemia. VII. Effect of long-term administration of antioxidants on the development of retinopathy. *Diabetes* 50: 1938–1942.
- Lanthony P, Cosson JP (1988). [The course of color vision in early diabetic retinopathy treated with Ginkgo biloba extract. A preliminary double-blind versus placebo study]. *J Fr Ophtalmol* 11: 671–674.

- Leal EC, Aveleira CA, Castilho AF, Serra AM, Baptista FI, Hosoya K *et al.* (2009). High glucose and oxidative/nitrosative stress conditions induce apoptosis in retinal endothelial cells by a caspase-independent pathway. *Exp Eye Res* 88: 983–991.
- Lee KS, Kim SR, Park SJ, Park HS, Min KH, Jin SM *et al.* (2006). Peroxisome proliferator activated receptor-gamma modulates reactive oxygen species generation and activation of nuclear factor-kappaB and hypoxia-inducible factor 1alpha in allergic airway disease of mice. *J Allergy Clin Immunol* 118: 120–127.
- Li DD, Chen JH, Chen Q, Li GW, Chen J, Yue JM *et al.* (2005). Swietenia mahagony extract shows agonistic activity to PPAR(gamma) and gives ameliorative effects on diabetic db/db mice. *Acta Pharmacol Sin* 26: 220–222.
- Li H, Zhu H, Xu CJ, Yuan J (1998). Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell* 94: 491–501.
- Li J, Liu NF, Wei Q (2008). Effect of rosiglitazone on cardiac fibroblast proliferation, nitric oxide production and connective tissue growth factor expression induced by advanced glycation end-products. *J Int Med Res* 36: 329–335.
- Li WL, Zheng HC, Bukuru J, De Kimpe N (2004). Natural medicines used in the traditional Chinese medical system for therapy of diabetes mellitus. *J Ethnopharmacol* 92: 1–21.
- Liang XC, Hagino N, Guo SS, Tsutsumi T, Kobayashi S (2002). Therapeutic efficacy of Stephania tetrandra S. Moore for treatment of neovascularization of retinal capillary (retinopathy) in diabetes-*in vitro* study. *Phytomedicine* 9: 377–384.
- Lin MT, Beal MF (2006). Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443: 787–795.
- Linsenmeier RA, Goldstick TK, Zhang SL (1989). Chinese herbal medicine increases tissue oxygen tension. *Adv Exp Med Biol* 248: 795–801.
- Liu JP, Zhang M, Wang WY, Grimsaard S (2004). Chinese herbal medicines for type 2 diabetes mellitus. *Cochrane Database Syst Rev* (3), Art. No. CD003642.
- Lopes de Faria JM, Katsumi O, Cagliero E, Nathan D, Hirose T (2001). Neurovisual abnormalities preceding the retinopathy in patients with long-term type 1 diabetes mellitus. *Graefes Arch Clin Exp Ophthalmol* 239: 643–648.
- Losso JN, Truax RE, Richard G (2010). Trans-resveratrol inhibits hyperglycemia-induced inflammation and connexin downregulation in retinal pigment epithelial cells. *J Agric Food Chem* 58: 8246–8252.
- Ma G, Zhu Y, Yu P, Shao Y, Wang Y (2004). Clinical observation on the traditional Chinese medicine combined with laser treatment for diabetic retinopathy. *Zhejiang J Inter Trad Chinese Western Med* 14: 762–763.
- Malchiodi-Albedi F, Matteucci A, Bernardo A, Minghetti L (2008). PPAR-gamma, microglial cells, and ocular inflammation: new venues for potential therapeutic approaches. *PPAR Res* 2008: 295784.
- Miller JW, Adamis AP, Shima DT, D'Amore PA, Moulton RS, O'Reilly MS *et al.* (1994). Vascular endothelial growth factor/vascular permeability factor is temporally and spatially correlated with ocular angiogenesis in a primate model. *Am J Pathol* 145: 574–584.
- Miyahara S, Kiryu J, Yamashiro K, Miyamoto K, Hirose F, Tamura H *et al.* (2004). Simvastatin inhibits leukocyte accumulation and vascular permeability in the retinas of rats with streptozotocin-induced diabetes. *Am J Pathol* 164: 1697–1706.
- Miyamoto K, Ogura Y (1999). Pathogenetic potential of leukocytes in diabetic retinopathy. *Semin Ophthalmol* 14: 233–239.
- Miyatake K, Kensho G, GFujimoto T, Noguchi E, Shinohara M, Takenaka S *et al.* (1994). Effects of conduritol A, a polyol from *Glynnema sylvestre*, on the development of diabetic cataracts in streptozotocin-treated rats and on aldose reductase. *Biosci Biotechnol Biochem* 58: 756–757.
- Mizutani M, Kern TS, Lorenzi M (1996). Accelerated death of retinal microvascular cells in human and experimental diabetic retinopathy. *J Clin Invest* 97: 2883–2890.
- Moore TC, Moore JE, Kaji Y, Frizzell N, Usui T, Poulaki V *et al.* (2003). The role of advanced glycation end products in retinal microvascular leukostasis. *Invest Ophthalmol Vis Sci* 44: 4457–4464.
- Motomura K, Fujiwara Y, Kiyota N, Tsurushima K, Takeya M, Nohara T *et al.* (2009). Astragalosides isolated from the root of astragalus radix inhibit the formation of advanced glycation end products. *J Agric Food Chem* 57: 7666–7672.
- Mrudula T, Suryanarayana P, Srinivas PN, Reddy GB (2007). Effect of curcumin on hyperglycemia-induced vascular endothelial growth factor expression in streptozotocin-induced diabetic rat retina. *Biochem Biophys Res Commun* 361: 528–532.
- Munch G, Schinzel R, Loske C, Wong A, Durany N, Li JJ *et al.* (1998). Alzheimer's disease-synergistic effects of glucose deficit, oxidative stress and advanced glycation endproducts. *J Neural Transm* 105: 439–461.
- Muranaka K, Yanagi Y, Tamaki Y, Usui T, Kubota N, Iriyama A *et al.* (2006). Effects of peroxisome proliferator-activated receptor gamma and its ligand on blood-retinal barrier in a streptozotocin-induced diabetic model. *Invest Ophthalmol Vis Sci* 47: 4547–4552.
- Murata T, He S, Hangai M, Ishibashi T, Xi XP, Kim S *et al.* (2000). Peroxisome proliferator-activated receptor-gamma ligands inhibit choroidal neovascularization. *Invest Ophthalmol Vis Sci* 41: 2309–2317.
- Murata T, Hata Y, Ishibashi T, Kim S, Hsueh WA, Law RE *et al.* (2001). Response of experimental retinal neovascularization to thiazolidinediones. *Arch Ophthalmol* 119: 709–717.
- Murray M (1997). Bilberry (*Vaccinium myrtillus*). *Am J Nat Med* 4: 18–22.
- Nagalakshmi MA, Thangadurai D, Rao DM, Pullaiah T (2001). Phytochemical and antimicrobial study of *Chukrasia tabularis* leaves. *Fitoterapia* 72: 62–64.
- Nagata S (1997). Apoptosis by death factor. *Cell* 88: 355–365.
- Obrosova IG, Julius UA (2005). Role for poly(ADP-ribose) polymerase activation in diabetic nephropathy, neuropathy and retinopathy. *Curr Vasc Pharmacol* 3: 267–283.
- van den Oever IA, Raterman HG, Nurmohamed MT, Simsek S (2010). Endothelial dysfunction, inflammation, and apoptosis in diabetes mellitus. *Mediators Inflamm* 2010: 792393.
- Park MY, Lee KS, Sung MK (2005). Effects of dietary mulberry, Korean red ginseng, and banaba on glucose homeostasis in relation to PPAR-alpha, PPAR-gamma, and LPL mRNA expressions. *Life Sci* 77: 3344–3354.
- Park SH (2008). Ginsenosides Rb2 protects high glucose-induced apoptosis in ARPE cells. *Lab Anim Res* 24: 199–123.
- Patz A (1982). Clinical and experimental studies on retinal neovascularization. XXXIX Edward Jackson Memorial Lecture. *Am J Ophthalmol* 94: 715–743.

- Perossini MG, Chiellini S Guidi G, Siravo D (1988). Diabetic and hypertensive retinopathy therapy with *Vaccinium myrtillus* anthocyanosides. *Ann Ottalmol Clin Ocul* 113: 1173–1190.
- Petrova TV, Akama KT, Van Eldik LJ (1999). Cyclopentenone prostaglandins suppress activation of microglia: down-regulation of inducible nitric-oxide synthase by 15-deoxy-Delta12,14-prostaglandin J2. *Proc Natl Acad Sci U S A* 96: 4668–4673.
- Preet A, Siddiqui MR, Taha A, Badhai J, Hussain ME, Yadava PK *et al.* (2006). Long-term effect of *Trigonella foenum graecum* and its combination with sodium orthovanadate in preventing histopathological and biochemical abnormalities in diabetic rat ocular tissues. *Mol Cell Biochem* 289: 137–147.
- Puri D, Prabhu KM, Murthy PS (2002). Mechanism of action of a hypoglycemic principle isolated from fenugreek seeds. *Indian J Physiol Pharmacol* 46: 457–462.
- Qi CX, Tan XH, Li QG, Wang XL (2007). [Clinical study of diabetic retinopathy treated by compound danshen dripping pills]. *Zhong Yao Cai* 30: 375–377.
- Qin S, McLaughlin AP, De Vries GW (2006). Protection of RPE cells from oxidative injury by 15-deoxy-delta12,14-prostaglandin J2 by augmenting GSH and activating MAPK. *Invest Ophthalmol Vis Sci* 47: 5098–5105.
- Rahbar S, Natarajan R, Yerneni K, Scott S, Gonzales N, Nadler JL *et al.* (2000). Evidence that pioglitazone, metformin and pentoxifylline are inhibitors of glycation. *Clin Chim Acta* 301: 65–77.
- Reilly CM, Oates JC, Sudian J, Crosby MB, Halushka PV, Gilkeson GS *et al.* (2001). Prostaglandin J(2) inhibition of mesangial cell iNOS expression. *Clin Immunol* 98: 337–345.
- Remels AH, Langen RC, Gosker HR, Russell AP, Spaapen F, Voncken JW *et al.* (2009). PPARgamma inhibits NF-kappaB-dependent transcriptional activation in skeletal muscle. *Am J Physiol Endocrinol Metab* 297: E174–E183.
- Ren Y, Sun C, Sun Y, Tan H, Wu Y, Cui B *et al.* (2009). PPAR gamma protects cardiomyocytes against oxidative stress and apoptosis via Bcl-2 upregulation. *Vascu Pharmacol* 51: 169–174.
- Ricote M, Huang J, Fajas L, Li A, Welch J, Najib J *et al.* (1998a). Expression of the peroxisome proliferator-activated receptor gamma (PPARgamma) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proc Natl Acad Sci U S A* 95: 7614–7619.
- Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK (1998b). The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature* 391: 79–82.
- Rizzolo LJ (1997). Polarity and the development of the outer blood-retinal barrier. *Histol Histopathol* 12: 1057–1067.
- Rodrigues GA, Maurier-Mahe F, Shurland DL, McLaughlin AP, Luhrs K, Throo E *et al.* (2010). Differential effects of PPARgamma ligands on oxidative stress-induced death of retinal pigmented epithelial cells. *Invest Ophthalmol Vis Sci* 52: 890–903.
- Roehr B (2010). FDA committee urges tight restrictions on rosiglitazone. *BMJ* 341: c3862.
- Romeo G, Liu WH, Asnaghi V, Kern TS, Lorenzi M (2002). Activation of nuclear factor-kappaB induced by diabetes and high glucose regulates a proapoptotic program in retinal pericytes. *Diabetes* 51: 2241–2248.
- Rosen ED, Spiegelman BM (2001). PPARgamma: a nuclear regulator of metabolism, differentiation, and cell growth. *J Biol Chem* 276: 37731–37734.
- SanGiovanni JP, Chew EY (2005). The role of omega-3 long-chain polyunsaturated fatty acids in health and disease of the retina. *Prog Retin Eye Res* 24: 87–138.
- Sarafidis PA, Bakris GL (2006). Protection of the kidney by thiazolidinediones: an assessment from bench to bedside. *Kidney Int* 70: 1223–1233.
- Sarayba MA, Li L, Tungsiripat T, Liu NH, Sweet PM, Patel AJ *et al.* (2005). Inhibition of corneal neovascularization by a peroxisome proliferator-activated receptor-gamma ligand. *Exp Eye Res* 80: 435–442.
- Sato T, Iwaki M, Shimogaito N, Wu X, Yamagishi S, Takeuchi M (2006). TAGE (toxic AGEs) theory in diabetic complications. *Curr Mol Med* 6: 351–358.
- Saxena A, Vikram NK (2004). Role of selected Indian plants in management of type 2 diabetes: a review. *J Altern Complement Med* 10: 369–378.
- Schmidt AM, Hori O, Brett J, Yan SD, Wautier JL, Stern D *et al.* (1994). Cellular receptors for advanced glycation end products. Implications for induction of oxidant stress and cellular dysfunction in the pathogenesis of vascular lesions. *Arterioscler Thromb* 14: 1521–1528.
- Schmidt M, Giessl A, Laufs T, Hankeln T, Wolfrum U, Burmester T *et al.* (2003). How does the eye breathe? Evidence for neuroglobin-mediated oxygen supply in the mammalian retina. *J Biol Chem* 278: 1932–1935.
- Schreck R, Albermann K, Baeuerle PA (1992). Nuclear factor kappa B: an oxidative stress-responsive transcription factor of eukaryotic cells (a review). *Free Radic Res Commun* 17: 221–237.
- Scoditti E, Massaro M, Carluccio MA, Distanto A, Storelli C, De Caterina R *et al.* (2009). PPARgamma agonists inhibit angiogenesis by suppressing PKCalpha- and CREB-mediated COX-2 expression in the human endothelium. *Cardiovasc Res* 86: 302–310.
- Sharma KR, Bhatia RP, Kumar V (1992). Role of the indigenous drug saptamrita lauha in hemorrhagic retinopathies. *Ann Ophthalmol* 24: 5–8.
- Shen J, Xu X, Cheng F, Liu H, Luo X, Chen K *et al.* (2003). Virtual screening on natural products for discovering active compounds and target information. *Curr Med Chem* 10: 2327–2342.
- Shen LQ, Child A, Weber GM, Folkman J, Aiello LP (2008). Rosiglitazone and delayed onset of proliferative diabetic retinopathy. *Arch Ophthalmol* 126: 793–799.
- Shen P, Liu MH, Ng TY, Chan YH, Yong EL (2006). Differential effects of isoflavones, from *Astragalus membranaceus* and *Pueraria thomsonii*, on the activation of PPARalpha, PPARgamma, and adipocyte differentiation *in vitro*. *J Nutr* 136: 899–905.
- Shi L, Zhang W, Zhou YY, Zhang YN, Li JY, Hu LH *et al.* (2008). Corosolic acid stimulates glucose uptake via enhancing insulin receptor phosphorylation. *Eur J Pharmacol* 584: 21–29.
- Siebenlist U, Franzoso G, Brown K (1994). Structure, regulation and function of NF-kappa B. *Annu Rev Cell Biol* 10: 405–455.
- Sobal G, Menzel EJ, Sinzinger H (2005). Troglitazone inhibits long-term glycation and oxidation of low-density lipoprotein. *J Cardiovasc Pharmacol* 46: 672–680.
- Song Y, Lv P, Tong J, Yan X (2006). Gegensu yan zhou xue wei zhu she zhi liao er xing tang niao bing xing shi wang mo bing 50 li. *Shaanxi J Trad Chin Med* 27: 472–474.

- Srivastava RK, Unterman TG, Shankar S (2009). FOXO transcription factors and VEGF neutralizing antibody enhance antiangiogenic effects of resveratrol. *Mol Cell Biochem* 337: 201–212.
- Stitt AW (2001). Advanced glycation: an important pathological event in diabetic and age related ocular disease. *Br J Ophthalmol* 85: 746–753.
- Straus DS, Glass CK (2001). Cyclopentenone prostaglandins: new insights on biological activities and cellular targets. *Med Res Rev* 21: 185–210.
- Su W, Dai DZ, Liu HR, Na T, Dai Y (2007). Upregulated endothelin system in diabetic vascular dysfunction and early retinopathy is reversed by CPU0213 and total triterpene acids from *Fructus Corni*. *Clin Exp Pharmacol Physiol* 34: 1228–1233.
- Sulistio MS, Zion A, Thukral N, Chilton R (2008). PPARgamma agonists and coronary atherosclerosis. *Curr Atheroscler Rep* 10: 134–141.
- Sung B, Park S, Yu BP, Chung HY (2006). Amelioration of age-related inflammation and oxidative stress by PPARgamma activator: suppression of NF-kappaB by 2,4-thiazolidinedione. *Exp Gerontol* 41: 590–599.
- Takahashi N, Kawada T, Goto T, Yamamoto T, Taimatsu A, Matsui N *et al.* (2002). Dual action of isoprenols from herbal medicines on both PPARgamma and PPARalpha in 3T3-L1 adipocytes and HepG2 hepatocytes. *FEBS Lett* 514: 315–322.
- Tawfik A, Sanders T, Kahook K, Akeel S, Elmarakby A, Al-Shabrawey M *et al.* (2009). Suppression of retinal peroxisome proliferator-activated receptor gamma in experimental diabetes and oxygen-induced retinopathy: role of NADPH oxidase. *Invest Ophthalmol Vis Sci* 50: 878–884.
- Tilg H, Moschen AR (2008). Inflammatory mechanisms in the regulation of insulin resistance. *Mol Med* 14: 222–231.
- Tontonoz P, Spiegelman BM (2008). Fat and beyond: the diverse biology of PPARgamma. *Annu Rev Biochem* 77: 289–312.
- Touyz RM, Schiffrin EL (2006). Peroxisome proliferator-activated receptors in vascular biology-molecular mechanisms and clinical implications. *Vascul Pharmacol* 45: 19–28.
- Uchimura K, Nakamura M, Enjoji M, Irie T, Sugimoto R, Muta T *et al.* (2001). Activation of retinoic X receptor and peroxisome proliferator-activated receptor-gamma inhibits nitric oxide and tumor necrosis factor-alpha production in rat Kupffer cells. *Hepatology* 33: 91–99.
- Vincent AM, Perrone L, Sullivan KA, Backus C, Sastry AM, Lastoskie C *et al.* (2007). Receptor for advanced glycation end products activation injures primary sensory neurons via oxidative stress. *Endocrinology* 148: 548–558.
- Wang HY, Cheng ML (2008). Effects of Dan-shao-hua-xian on expression of PPAR-gamma and NF-kappa B in rat liver fibrosis. *Hepatobiliary Pancreat Dis Int* 7: 179–184.
- Wang Y, Zhao H, Sheng X, Gambino PE, Costello B, Bojanowski K *et al.* (2002). Protective effect of *Fructus Lycii* polysaccharides against time and hyperthermia-induced damage in cultured seminiferous epithelium. *J Ethnopharmacol* 82: 169–175.
- Watson GS, Cholerton BA, Reger MA, Baker LD, Plymate SR, Asthana S *et al.* (2005). Preserved cognition in patients with early Alzheimer disease and amnesic mild cognitive impairment during treatment with rosiglitazone: a preliminary study. *Am J Geriatr Psychiatry* 13: 950–958.
- Wei G, Liu M, Wen Z, Zhou R (2005). Yin xing da mo zhu she ye lian he shi wang mo ji guang guang ning shu zhi liao tang niao bing shi wang mo bing bian jin qi liao xiao guan cha. *Zhongguo Zong He Lin Chuang [Clin Med China]* 21: 833–835.
- West AL, Oren GA, Moroi SE (2006). Evidence for the use of nutritional supplements and herbal medicines in common eye diseases. *Am J Ophthalmol* 141: 157–166.
- Williams R, Airey M, Baxter H, Forrester J, Kennedy-Martin T, Girach A *et al.* (2004). Epidemiology of diabetic retinopathy and macular oedema: a systematic review. *Eye* 18: 963–983.
- Winkler BS, Boulton ME, Gottsch JD, Sternberg P (1999). Oxidative damage and age-related macular degeneration. *Mol Vis* 5: 32.
- Wohlmuth H, Oliver C, Nathan PJ (2002). A review of the status of Western herbal medicine in Australia. *J Herb Pharmacother* 2: 33–46.
- Wong A, Dukic-Stefanovic S, Gasic-Milenkovic J, Schinzel R, Wiesinger H, Riederer P *et al.* (2001). Anti-inflammatory antioxidants attenuate the expression of inducible nitric oxide synthase mediated by advanced glycation endproducts in murine microglia. *Eur J Neurosci* 14: 1961–1967.
- Wu G, Mannam AP, Wu J, Kirbis S, Shie JL, Chen C *et al.* (2003). Hypoxia induces myocyte-dependent COX-2 regulation in endothelial cells: role of VEGF. *Am J Physiol Heart Circ Physiol* 285: H2420–H2429.
- Wu JS, Lin TN, Wu KK (2009). Rosiglitazone and PPAR-gamma overexpression protect mitochondrial membrane potential and prevent apoptosis by upregulating anti-apoptotic Bcl-2 family proteins. *J Cell Physiol* 220: 58–71.
- Xin X, Yang S, Kowalski J, Gerritsen ME (1999). Peroxisome proliferator-activated receptor gamma ligands are potent inhibitors of angiogenesis *in vitro* and *in vivo*. *J Biol Chem* 274: 9116–9121.
- Yam JC, Kwok AK (2007). Update on the treatment of diabetic retinopathy. *Hong Kong Med J* 13: 46–60.
- Yamagishi S, Nakamura K, Matsui T (2007). Potential utility of telmisartan, an angiotensin II type 1 receptor blocker with peroxisome proliferator-activated receptor-gamma (PPAR-gamma)-modulating activity for the treatment of cardiometabolic disorders. *Curr Mol Med* 7: 463–469.
- Yanagi Y (2008). Role of Peroxisome Proliferator Activator Receptor gamma on Blood Retinal Barrier Breakdown. *PPAR Res* 2008: 679237.
- Yeh GY, Eisenberg DM, Kaptchuk TJ, Phillips RS (2003). Systematic review of herbs and dietary supplements for glycemic control in diabetes. *Diabetes Care* 26: 1277–1294.
- Yu X, Li C, Li X, Cai L (2007). Rosiglitazone prevents advanced glycation end products-induced renal toxicity likely through suppression of plasminogen activator inhibitor-1. *Toxicol Sci* 96: 346–356.
- Yuuki T, Kanda T, Kimura Y, Kotajima N, Tamura J, Kobayashi I *et al.* (2001). Inflammatory cytokines in vitreous fluid and serum of patients with diabetic vitreoretinopathy. *J Diabetes Complications* 15: 257–259.
- Zeng K, Xu H, Mi M, Zhang Q, Zhang Y, Chen K *et al.* (2009). Dietary taurine supplementation prevents glial alterations in retina of diabetic rats. *Neurochem Res* 34: 244–254.
- Zeng K, Xu H, Mi M, Chen K, Zhu J, Yi L *et al.* (2010). Effects of taurine on glial cells apoptosis and taurine transporter expression in retina under diabetic conditions. *Neurochem Res* 35: 1566–1574.

Zhang SL, Lax D, Li Y, Stejskal E, Lucas RV Jr, Einzig S *et al.* (1990). Anisodamine increases blood flow to the retina-choroid and protects retinal and pancreatic cells against lipid peroxidation. *J Ethnopharmacol* 30: 121–134.

Zhang SX, Wang JJ, Gao G, Parke K, Ma JX *et al.* (2006). Pigment epithelium-derived factor downregulates vascular endothelial growth factor (VEGF) expression and inhibits VEGF-VEGF receptor 2 binding in diabetic retinopathy. *J Mol Endocrinol* 37: 1–12.

Zhou Y, Zheng S, Lin J, Zhang QJ, Chen A (2007). The interruption of the PDGF and EGF signaling pathways by curcumin stimulates gene expression of PPARgamma in rat activated hepatic stellate cell *in vitro*. *Lab Invest* 87: 488–498.

Zinn A, Felson S, Fisher E, Schwartzbard A (2008). Reassessing the cardiovascular risks and benefits of thiazolidinediones. *Clin Cardiol* 31: 397–403.