

Themed Section: Secretin Family (Class B) G Protein-Coupled Receptors –
from Molecular to Clinical Perspectives

RESEARCH PAPER

The third extracellular loop of the human calcitonin receptor-like receptor is crucial for the activation of adrenomedullin signalling

Kenji Kuwasako¹, Debbie L Hay², Sayaka Nagata³, Tomomi Hikosaka³,
Kazuo Kitamura³ and Johji Kato¹

¹Frontier Science Research Center, University of Miyazaki, Miyazaki, Japan, ²School of Biological
Sciences, University of Auckland, Auckland, New Zealand, and ³Division of Circulation and
Body Fluid Regulation, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

Correspondence

Kenji Kuwasako, Frontier Science
Research Center, University of
Miyazaki, 5200 Kihara, Kiyotake,
Miyazaki, Miyazaki 889-1692,
Japan. E-mail:
kuwasako@fc.miyazaki-u.ac.jp

Keywords

adrenomedullin receptor;
adrenomedullin; calcitonin
receptor-like receptor;
extracellular loop; chimera;
receptor activation; receptor
activity-modifying protein

Received

11 March 2011

Revised

25 October 2011

Accepted

11 November 2011

BACKGROUND AND PURPOSE

The extracellular loops (ECLs) in Family A GPCRs are important for ligand binding and receptor activation, but little is known about the function of Family B GPCR ECLs, especially ECL3. Calcitonin receptor-like receptor (CLR), a Family B GPCR, functions as a calcitonin gene-related peptide (CGRP) and an adrenomedullin (AM) receptor in association with three receptor activity-modifying proteins (RAMPs). Here, we examined the function of the ECL3 of human CLR within the CGRP and AM receptors.

EXPERIMENTAL APPROACH

A CLR ECL3 chimera, in which the ECL3 of CLR was substituted with that of VPAC2 (a Family B GPCR that is unable to interact with RAMPs), and CLR ECL3 point mutants were constructed and transiently transfected into HEK-293 cells along with each RAMP. Cell-surface expression of each receptor complex was then measured by flow cytometry; [¹²⁵I]-CGRP and [¹²⁵I]-AM binding and intracellular cAMP accumulation were also measured.

KEY RESULTS

Co-expression of the CLR ECL3 chimera with RAMP2 or RAMP3 led to significant reductions in the induction of cAMP signalling by AM, but CGRP signalling was barely affected, despite normal cell-surface expression of the receptors and normal [¹²⁵I]-AM binding. The chimera had significantly decreased AM, but not CGRP, responses in the presence of RAMP1. Not all CLR ECL3 mutants supported these findings.

CONCLUSIONS AND IMPLICATIONS

The human CLR ECL3 is crucial for AM-induced cAMP responses via three CLR/RAMP heterodimers, and activation of these heterodimers probably relies on AM-induced conformational changes. This study provides a clue to the molecular basis of the activation of RAMP-based Family B GPCRs.

LINKED ARTICLES

This article is part of a themed section on Secretin Family (Class B) G Protein-Coupled Receptors. To view the other articles in this section visit <http://dx.doi.org/10.1111/bph.2012.166.issue-1>

Abbreviations

AM, adrenomedullin; CGRP, calcitonin gene-related peptide; CLR, calcitonin receptor-like receptor; ECL, extracellular loop; RAMP, receptor activity-modifying protein; TM, transmembrane domain

Introduction

Among the Family B GPCRs, calcitonin receptor-like receptor (CLR) is the first known partner of all three receptor activity-modifying proteins (RAMPs) (McLatchie *et al.*, 1998). When acting as chaperones, RAMPs transport CLR to the cell surface, where CLR/RAMP1 forms the functional calcitonin gene-related peptide (CGRP) receptor. CLR/RAMP2 and -3 both form adrenomedullin (AM) receptors, although CLR/RAMP2 (AM₁ receptor) is more specific for AM than CLR/RAMP3 (AM₂ receptor) (Poyner *et al.*, 2002; Hay *et al.*, 2003; Muff *et al.*, 2003). CLR/RAMP1 can also induce a similarly strong response to higher concentrations of AM (Kuwasako *et al.*, 2004a). AM, like CGRP, is a potent vasodilator that also exerts strong protective effects against multi-organ damage (Gibbons *et al.*, 2007; Kuwasako *et al.*, 2011b).

Recent crystal structural analysis revealed that activation of Family A GPCRs is accompanied by movements among the seven transmembrane domains (TMs) (Wess *et al.*, 2008; Topiol and Sabio, 2009; Simpson *et al.*, 2011; Standfuss *et al.*, 2011). A similar phenomenon is believed to also occur in Family B GPCRs (Conner *et al.*, 2007a; Chugunov *et al.*, 2010), but no crystal structure of a whole receptor is currently available. It is known from the crystal structures of Family A GPCRs that their extracellular loops (ECLs) are orientated to interact with each other and with the TMs. In addition to serving as linkers between TMs, the ECLs of Family A GPCRs are known to be important determinants of ligand binding and receptor activation (Lawson and Wheatley, 2004; Hawtin *et al.*, 2006; Peeters *et al.*, 2011). The orientation of ECL2 in the majority of Family A GPCRs is restricted by a conserved disulfide bond between ECL2 and the top of TM3 (Conner *et al.*, 2007b). Similarly, ECLs 1 and 2 in Family B GPCRs form a conserved disulfide bond for receptor stabilization (Kuwasako *et al.*, 2003). The Family B GPCR ECLs are important for peptide ligand binding and receptor activation, but little is known about the function of ECL3, and there have been no reports on the function of TMs 1, 2, 4, 5 and 7 of CLR (Walker *et al.*, 2010; Wheatley *et al.*, 2012).

As with Family A GPCRs, the functions of the various regions of Family B GPCRs have been investigated using a chimeric receptor strategy, in which the target sequence of one GPCR was replaced with the corresponding sequence from a different GPCR (Nielsen *et al.*, 2000; Van Rampelbergh *et al.*, 2000; Unson *et al.*, 2002; Runge *et al.*, 2003; Koller *et al.*, 2004; Salvatore *et al.*, 2006; Kuwasako *et al.*, 2009). To determine which structural regions of human (h)CLR govern CLR/RAMP trafficking and function, we recently generated a set of nine chimeras (CH-1 to CH-9) in which regions of CLR were replaced with corresponding sequences from vasoactive intestinal peptide (VIP)/pituitary adenylate cyclase-activating polypeptide type 2 receptor (VPAC2; another Family B GPCR), which does not interact with RAMPs (Christopoulos *et al.*, 2003). Using these chimeras, we first determined the CLR regions (between TM1 and TM5) responsible for the trafficking interactions with each RAMP (Kuwasako *et al.*, 2009). Our preliminary experiments showed that CH-7, which contained ECL3 and TM7 from VPAC2, impaired the production of cAMP by AM via the AM₁ and AM₂ receptors but affected CGRP responses via the CGRP receptor to a lesser extent.

To investigate the function of the CLR ECL3 within the CGRP and AM receptors, we generated CH-I and CH-II, in which the ECL3 and TM7 of CLR were each substituted with the corresponding sequences from VPAC2. Our findings suggest that the CLR ECL3 plays a key role in AM-induced cAMP signalling via the three CLR/RAMP heterodimers, and the activation of these heterodimers probably relies on AM-induced conformational changes.

Methods

Expression constructs

Double V5 epitope-tagged human CLR (V5-CLR) and all three V5-RAMPs were prepared as described previously (Kuwasako *et al.*, 2009) and cloned into the mammalian expression vector pCAGGS/Neo (Kuwasako *et al.*, 2000). Corresponding untagged constructs served as controls.

Chimera construction

We previously used seven restriction sites to construct nine human CLR chimeras (CH-1 to CH-9), in which CLR domains were sequentially substituted with corresponding sequences from the human VPAC2 receptor (Kuwasako *et al.*, 2009). Of those chimeras, CH-7, in which the ECL3 and TM7 of CLR were exchanged with those of VPAC2, was used in the present study (Figure 1). As shown in Figure 1B, we also constructed two new CLR/VPAC2 chimeras, CH-I and CH-II, in which either CLR ECL3 or TM7 was replaced with the corresponding sequence from VPAC2 using three restriction sites *Eco*T14 I (*Sty* I), *Eco*T22 I (*Ava* III) and *Avi* II (*Mst* I). Human CLR naturally possesses an *Eco*T14 I site, and the other two sites were introduced without altering the amino acid sequence of the receptor. Human VPAC2 shares only 30% amino acid sequence identity with human CLR and contains none of the three restriction sites. Therefore, corresponding VPAC2 fragments containing the necessary restriction sites were prepared by PCR using a primer set containing the sites 5'-*Eco*T14 I-VPAC2 ECL3-*Eco*T22 I-3' and 5'-*Eco*T22 I-VPAC2 TM7-*Avi* II-3'. The separate CLR and VPAC2 fragments were then ligated into the same pCAGGS/Neo expression vector. The resulting chimeric constructs were all sequenced using an Applied Biosystems 310 Genetic Analyzer (Foster City, CA, USA).

Site-directed mutagenesis

Single and triple amino acid substitutions were carried out using a QuikChange kit (Stratagene Corporation, La Jolla, CA, USA) according to the manufacturer's instructions, with pIRES1-V5-CLR serving as the template. pIRES1-V5-CLR was constructed by subcloning the coding sequence of human V5-CLR into pIRES/Neo (Clontech, Palo Alto, CA). For each mutation, two complementary 30- to 40-mer oligonucleotides (sense and antisense) were designed with the mutation in the middle. The resulting mutants were all sequenced using an Applied Biosystems 310 Genetic Analyzer.

Cell culture and DNA transfection

HEK-293 cells were maintained in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% heat-inactivated




Figure 1

(A) Amino acid sequence alignment of ECL3 and TM7 of human CLR with the corresponding domains of the human VPAC2 receptor. Alignment of CLR ECL3 and TM7 is based on a previous report (Bailey and Hay, 2007). A homology search for both receptors was performed using DNASIS (Hitachi Solutions, Ltd, Tokyo, Japan). Vertical dashes indicate conserved amino acids. (B) Alignment of the three CLR/VPAC2 chimeras. Chimeras were constructed using three restriction sites: *Eco*T14 I, *Eco*T22 I and *Avi* II (see Methods section). CH-7 was generated by replacing CLR ECL3 and TM7 with the corresponding sequence from VPAC2 using 5' *Eco*T14 I and 3' *Avi* II sites (Kuwasako *et al.*, 2009). Likewise, CH-I and CH-II were constructed by substituting CLR ECL3 (using 5' *Eco*T14 I and 3' *Eco*T22 I sites) or CLR TM7 (using 5' *Eco*T22 I and 3' *Avi* II sites), respectively.

fetal bovine serum (FBS), 100 U·mL⁻¹ penicillin G, 100 µg·mL⁻¹ streptomycin and 0.25 µg·mL⁻¹ amphotericin B at 37°C under a humidified atmosphere of 95% air/5% CO₂. Transient transfection of the cells was accomplished using Lipofectamine™ with Plus™ reagent (Invitrogen Corporation, Carlsbad, CA, USA) as previously described (Kuwasako *et al.*, 2011a). Briefly, the cells were seeded into 12-well plates (for flow cytometric analysis) or 24-well plates (for binding and cAMP assays) and, upon reaching 70–80% confluence, were transfected with empty vector (pCAGGS/Neo or pIRES/Neo) (Mock) or V5-tagged wild-type (WT), pCAGGS-chimeric or pIRES-mutant constructs; V5-CLR was included in each transfection set. DNA complex with transfection reagents was formed by incubating the cells for 4 h in OptiMEM 1 medium containing plasmid DNAs (0.2 µg per well for 24-well plates; 0.4 µg per well for 12-well plates), Plus reagent (2 µL per well for 24-well plates; 2.5 µL per well for 12-well plates) and Lipofectamine reagent (2 µL per well for 24-well plates; 2.5 µL per well for 12-well plates). All experiments were performed 36–48 h after transfection.

Flow cytometry

Flow cytometry was used to assess the cell-surface expression levels of V5-tagged receptor proteins. Following transient co-transfection of the indicated cDNAs for WT or mutant V5-CLR and RAMP or those for WT or mutant CLR and V5-RAMP into HEK-293 cells in 12-well plates, the cells were washed once with ice-cold PBS and then non-enzymatically harvested with ice-cold FACS buffer (Kuwasako *et al.*, 2000). After centrifugation at 200× *g* for 4 min at 4°C, the cells were resuspended in FACS buffer and labelled with fluorescein isothiocyanate (FITC) conjugated mouse anti-V5 monoclonal antibody (anti-V5-FITC antibody) (Invitrogen, diluted 1:1000

in FACS buffer) for 2 h at 4°C in the dark. The cells were then washed twice with ice-cold FACS buffer, resuspended to a density of 2 × 10⁵ cells per tube in FACS buffer containing 5 µg·mL⁻¹ propidium iodide and subjected to flow cytometry in an EPICS XL flow cytometer (Beckman Coulter, Fullerton, CA, USA). The cell-surface expression frequency of each V5-tagged receptor (% of cells) was analysed using EXPO 2 software (Beckman Coulter) (Kuwasako *et al.*, 2000). FITC fluorescence was excited at 488 nm, and emission was monitored at 530 nm. Viability was assessed on the basis of the exclusion of propidium iodide.

Radioligand binding

[¹²⁵I]-[His¹⁰]-αCGRP (specific activity 2.2 µCi·pmol⁻¹) was purchased from PerkinElmer (Yokohama, Japan), and [¹²⁵I]-AM (specific activity 2 µCi·pmol⁻¹) was produced in our laboratory as previously described (Kitamura *et al.*, 1994). To assess whole-cell radioligand binding, transfected HEK-293 cells in 24-well plates were washed once with pre-warmed PBS and incubated for 20 min at 37°C with 0.1% BSA/PBS to reduce non-specific binding of αCGRP and AM, after which, the remaining adherent cells were washed with ice-cold PBS. The cells were then incubated with 40 000 c.p.m. of [¹²⁵I]-αCGRP or 100 000 c.p.m. of [¹²⁵I]-AM for 4 h at 4°C in the absence (for total binding) or presence of different concentrations of unlabelled αCGRP or AM as appropriate (1 µM was used to define non-specific binding) in modified Krebs-Ringers-HEPES medium (Kuwasako *et al.*, 2000). After washing once with ice-cold PBS, the cells were solubilized with 0.5 mL of 0.5 M NaOH, and the associated cellular radioactivity was measured in a γ-counter. Specific binding was defined as the difference between the total binding and non-specific binding.

Measurement of intracellular cAMP

cAMP assays were carried out as described previously (Kuwasako *et al.*, 2011a). Transfectants in 24-well plates were incubated for 15 min at 37°C in Hanks' buffer containing 20 mM HEPES, 0.2% BSA, 0.5 mM 3-isobutyl-1-methylxanthine (IBMX) and the indicated concentrations of AM. The reactions were terminated by the addition of lysis buffer (GE Healthcare, Tokyo, Japan), after which, the cAMP content was determined using a commercial enzyme immunoassay kit according to the manufacturer's instructions (GE Healthcare) for a non-acetylation protocol.

Data analysis and statistics

At least five independent replicates were performed for each of the aforementioned experiments, and the results are expressed as mean $\pm$ SEM. Data were analysed using Prism 5.02 (GraphPad Software Inc., San Diego, CA, USA). The binding and cAMP data were fitted to obtain pIC_{50} and pEC_{50} values, respectively. A four-parameter logistic equation was used for curve fitting. For each fit, the Hill slope was compared to unity by *F*-test. When the Hill slope was not different from one, the data were re-fitted to a three-parameter logistic equation (Hill slope of one). In cases where the Hill slope differed from one, further analysis was performed to ascertain whether Hill slopes differed within a particular data set by *F*-test. Hill slopes that deviate from unity are reported within the results narrative. Differences among multiple groups were evaluated using one-way ANOVA as appropriate, where $P < 0.05$ was considered significant. *Post hoc* testing was via Dunnett's test for comparison with control.

Reagents and other materials

All ^{125}I -labelled and unlabelled ligands used were of human origin. α CGRP was purchased from the Peptide Institute (Osaka, Japan), and AM was kindly donated by Shionogi & Co. (Osaka, Japan). Peptides were dissolved in sterile, distilled H₂O to a concentration of 1 mM and stored as aliquots at -80°C in siliconized microcentrifuge tubes. IBMX and BSA were from Sigma-Aldrich Corporation (St Louis, MO, USA). DMEM and FBS were from Invitrogen. All other reagents were of analytical grade and obtained from various commercial suppliers.

Results

Cell-surface expression of CLR/VPAC2 chimeras co-expressed with RAMPs

We initially evaluated the effects of transient co-transfection of WT-RAMPs on the cell-surface expression of V5-CH-7, V5-CH-I and V5-CH-II in HEK-293 cells (Figure 2A), which do not endogenously express any functional CGRP or AM receptors (Kuwasako *et al.*, 2004b; 2008). In cells transfected with the empty vector (Mock), surface binding of an anti-V5-FITC antibody was within the 2% limit of resolution characteristic of flow cytometry. When expressed alone, FITC-labelled V5-CLR was detected in ~30% of cells. This phenomenon has also been seen in COS-7 cells, which are frequently used in transfection studies and also express no functional RAMPs




Figure 2

(A) Flow cytometric analysis of the cell-surface expression of WT or chimeric V5-CLR (CH-7, CH-I and CH-II) following transfection into HEK-293 cells, with or without WT-RAMP1, -2 or -3. Cell-surface expression of each FITC-labelled receptor protein was estimated by flow cytometry. Data are shown as the means $\pm$ SEM of five separate experiments. * $P < 0.05$ vs. corresponding Vector/V5-CLR or -CH, # $P < 0.05$ vs. corresponding V5-CLR/WT-RAMP. (B) Flow cytometric analysis of the cell-surface expression of V5-RAMP1, -2 or -3 following transfection into HEK-293 cells, with WT or chimeric CLR (CH-7, CH-I and CH-II). Cell-surface expression of each construct was analysed by flow cytometry. Data are shown as the means $\pm$ SEM of six separate experiments. * $P < 0.05$ vs. corresponding Vector/V5-RAMP, # $P < 0.05$ vs. corresponding WT-CLR/V5-RAMP.

(Ittner *et al.*, 2004; Koller *et al.*, 2004; Kuwasako *et al.*, 2009). In the present study, the frequency of surface WT V5-CLR and chimeric V5-CLR expression was significantly increased by the co-transfection of each RAMP. In the absence of exogenous RAMPs, two chimeras, CH-7 and CH-I, appeared at the cell surface at levels nearly the same as that seen with Mock. We compared the expression of each chimera to WT, when expressed with each RAMP. With RAMP1, each chimera caused a small but significant decrease in cell-surface delivery, compared to WT. For RAMP2 and RAMP3, only CH-I delivery was slightly reduced.

We also assessed the changes in the frequency of cell-surface expression of V5-RAMPs that occurred after co-transfection of untagged WT and chimeric CLR (Figure 2B). In contrast to V5-RAMP1 and V5-RAMP2, V5-RAMP3 appeared at the surface of ~28% of cells when the protein was expressed alone, probably because of its self-




Figure 3

Specific binding of [^{125}I]-CGRP (A) and [^{125}I]-AM (B) to heterodimeric receptors composed of WT-RAMP and WT or chimeric CLR (CH-7, CH-I and CH-II). HEK-293 cells were transiently transfected with a WT-RAMP and WT-CLR or one of its three chimeras. Data are shown as the means $\pm$ SEM of five to eight experiments in (A) and five to nine experiments in (B) and are expressed as percentages of CLR/RAMP1 (A) or CLR/RAMP2 (B). * $P < 0.05$ vs. corresponding WT-CLR/RAMP (A, B).

transport to the cell surface (Flahaut *et al.*, 2002) or its interaction with other endogenous GPCRs (Hay *et al.*, 2006; Sexton *et al.*, 2009). CH-7, CH-I and CH-II all markedly increased the frequency of the surface expression of V5-RAMP1 and V5-RAMP2 to a level comparable to that seen with CLR. Significant increases in V5-RAMP3 translocation were also observed when it was co-expressed with the WT or chimeric CLR constructs, but the magnitude of the changes was much smaller than was seen with V5-RAMP1 and V5-RAMP2. The expression frequency of RAMP1 was




Figure 4

Competitive binding of [^{125}I]-CGRP to WT or chimeric CLR/RAMP1 (A) and binding of [^{125}I]-AM to WT or chimeric CLR/RAMP2 (B) or CLR/RAMP3 (C). HEK-293 cells were transiently co-transfected with each WT-RAMP and WT or chimeric CLR (CH-7, CH-I and CH-II). Data were normalized to the maximum specific binding in each experiment. Data are shown as the means $\pm$ SEM of five experiments in (A, C) and six experiments in (B). (B) Binding of [^{125}I]-CGRP (A) or [^{125}I]-AM (B, C); B_0 , total binding in the presence of competing unlabelled CGRP (A) or AM (B, C); N, nonspecific binding (measured in the presence of 10^{-6} M CGRP (A) or 10^{-6} M AM (B, C)). pIC_{50} values are given in Table 1.

unchanged in the presence of each chimera when compared to WT. A significant decrease in RAMP2 and RAMP3 expression was observed in the presence of CH-I. CH-II only reduced the expression of RAMP3.

[¹²⁵I]-CGRP and [¹²⁵I]-AM binding to cell-surface CLR/VPAC2 chimeras

We next evaluated the binding of [¹²⁵I]-CGRP and [¹²⁵I]-AM to cells expressing each of the WT and chimeric receptors (Figures 3 and 4). Cells transfected with Mock showed only low levels of specific radioligand binding (Figure 3), but marked increases in [¹²⁵I]-CGRP binding were observed in cells expressing RAMP1 with WT-CLR, CH-I or CH-II (Figure 3A). The specific binding of [¹²⁵I]-CGRP to CH-7/RAMP1 was also increased but was much lower than the binding to CH-I/RAMP1 and CH-II/RAMP1. Cells co-expressing RAMP2 or RAMP3 with WT-CLR or one of the chimeras bound little [¹²⁵I]-CGRP. In contrast, marked increases in specific [¹²⁵I]-AM binding were seen with all four CLR proteins co-expressed with RAMP2 or RAMP3 (Figure 3B). There was no significant difference between the specific [¹²⁵I]-AM binding to CH-I/RAMP2 and CLR/RAMP2. However, the specific binding of [¹²⁵I]-AM to CH-7/RAMP2 was significantly lower than to CLR/RAMP2, while the specific binding of [¹²⁵I]-AM to CH-II/RAMP2 was significantly higher than to CLR/RAMP2. CLR/RAMP3 exhibited a lower level of specific [¹²⁵I]-AM binding than CLR/RAMP2, which is consistent with earlier observations (Kuwasako *et al.*, 2006). Interestingly, the specific binding of [¹²⁵I]-AM to CH-II/RAMP3 was about twofold higher than was seen with the other three RAMP3/CLR proteins. Figure 4 shows [¹²⁵I]-CGRP and [¹²⁵I]-AM competition curves for the WT and chimeric receptors. The pIC₅₀ values derived from the curves are given in Table 1. CH-7/RAMP1 (Hill slope = 0.68) and CH-II/RAMP1 exhibited significantly higher CGRP affinity than CLR/RAMP1 (Hill slope = 0.64), whereas the affinity of CGRP for CH-I/RAMP1 (Hill

slope = 0.69) was significantly lower than that for CLR/RAMP1. Although some of the Hill slopes for these curves deviated from unity, they were not different from each other. CH-7/RAMP2 (Hill slope = 0.70) exhibited significantly higher AM affinity than CLR/RAMP2, but the affinity of AM for CH-I/RAMP2 and CH-II/RAMP2 (Hill slope = 1.29) was similar to that for CLR/RAMP2. In the case of this RAMP2 data set, the Hill slopes were different between chimeras and WT. The affinity of AM was significantly increased at CH-7/RAMP3, CH-I/RAMP3 (Hill slope = 0.81) and CH-II/RAMP3 relative to CLR/RAMP3.

CGRP- and AM-induced cAMP production via CLR/VPAC2 chimeras

The three chimeric receptors were further characterized by measuring agonist-induced cAMP accumulation (Figure 5). Their pEC₅₀ and E_{max} values are given in Table 2. As previously reported (Kuwasako *et al.*, 2009), neither CGRP nor AM elicited cAMP production in HEK-293 cells transfected with Mock (data not shown).

When co-expressed with RAMP1, there were no significant decreases in the pEC₅₀ and E_{max} values for CGRP with CH-I or CH-II compared with WT-CLR (Figure 5A, C and D). That is, CGRP was able to activate all three RAMP1-associated receptors with similar potency and efficacy. In contrast, CGRP acting via CH-7/RAMP1 showed an approximately 60-fold reduction in potency with no significant reduction in efficacy (Figure 5B). Recombinant CLR/RAMP1 can also respond fully to higher concentrations of AM (Kuwasako *et al.*, 2004a). In cells expressing CLR/RAMP1, the potency of AM was about 60-fold lower than that of CGRP (Figure 5A). Although the potency of CGRP and AM for CH-II/RAMP1 was not significantly different from that for CLR/RAMP1 (Table 2, Figure 5D), CH-7/RAMP1 significantly reduced the potency of each agonist (Figure 5B). Surprisingly, in cells expressing CH-I/RAMP1, the potency of AM was about 460-fold lower than that of CGRP, despite the fact that CH-I/RAMP1 mediated CGRP-induced cAMP responses normally (Figure 5C). For AM potency, only the Hill slope (0.63) of CH-II/RAMP1 differed from 1.

In cells expressing CLR/RAMP2 (AM₁ receptor), the potency of AM was about 210-fold higher than that of CGRP (Figure 5E), while in cells expressing CLR/RAMP3 (AM₂ receptor), the potency of AM was about 100-fold higher (Figure 5I). Cells expressing RAMP2 with CH-7 or CH-I showed marked reductions in AM potency and efficacy (Figure 5F and G). The potency of AM with CH-7/RAMP2 and CH-I/RAMP2 was reduced by about 60-fold and 100-fold, respectively, and the efficacy was reduced by about 80% and 50% compared with the WT AM₁ receptor (Figure 5F and G). Most notably, CH-I/RAMP2 exhibited normal [¹²⁵I]-AM binding (Figure 3B and Table 1). On the other hand, the potency of AM for CH-7/RAMP3 and CH-I/RAMP3 was decreased by about 15-fold and 20-fold, respectively, and the efficacy was decreased by about 40% and 30% compared with the WT AM₂ receptor (Figure 5J and K). In contrast, the efficacy of AM with CH-II/RAMP2 and CH-II/RAMP3 was not significantly different from that for WT AM₁ and AM₂ receptors, respectively (Figure 5H and L), although the AM potency for CH-II/RAMP2 and the CGRP potency for CH-II/RAMP3 were significantly reduced relative to each WT AM

Table 1

pIC₅₀ values for CGRP in competition with [¹²⁵I]-CGRP binding to RAMP1/WT or chimeric CLR and for AM in competition with [¹²⁵I]-AM binding to RAMP2/WT or chimeric CLR and RAMP3/WT or chimeric CLR

Receptor complex	pIC ₅₀ ± SEM	n
RAMP1/WT-CLR	8.43 ± 0.07	5
RAMP1/CH-7	8.91 ± 0.10*	5
RAMP1/CH-I	8.01 ± 0.06*	5
RAMP1/CH-II	8.76 ± 0.08*	5
RAMP2/WT-CLR	7.72 ± 0.04	6
RAMP2/CH-7	8.09 ± 0.05*	6
RAMP2/CH-I	7.64 ± 0.04	6
RAMP2/CH-II	7.68 ± 0.02	6
RAMP3/WT-CLR	7.56 ± 0.06	5
RAMP3/CH-7	8.13 ± 0.04*	5
RAMP3/CH-I	7.80 ± 0.05*	5
RAMP3/CH-II	7.81 ± 0.04*	5

*P < 0.05 vs. the corresponding RAMP/WT-CLR one-way ANOVA followed by Dunnett's multiple comparison test.

RAMP, receptor activity-modifying protein; WT, wild-type; CLR, calcitonin receptor-like receptor; CGRP, calcitonin gene-related peptide; AM, adrenomedullin; CH, chimera.




Figure 5

Agonist-induced cAMP production elicited via heterodimeric receptors comprised of WT-RAMP and WT or chimeric CLR (CH-7, CH-I and CH-II). HEK-293 cells were co-transfected with WT-RAMP1 plus WT-CLR (A), CH-7 (B), CH-I (C) or CH-II (D); WT-RAMP2 plus WT-CLR (E), CH-7 (F), CH-I (G) or CH-II (H); or WT-RAMP3 plus WT-CLR (I), CH-7 (J), CH-I (K) or CH-II (L). All transfected cells (A–L) were simultaneously exposed to the indicated concentrations of α CGRP (open symbols) or AM (solid symbols) 48 h after transfection. Data are shown as the means $\pm$ SEM of five separate experiments. pEC₅₀ values are given in Table 2.

receptor. Among these RAMP2- and RAMP3-based receptors, only the Hill slopes of CH-I/RAMP2 (0.49 for AM) and CH-I/RAMP3 (0.62 for CGRP) were different from 1.

Further investigation of the function of CLR ECL3 in AM receptors

The data summarized earlier indicate that among the three CLR/VPAC2 chimeras tested, co-transfection of CH-7 with RAMP1, RAMP2 or RAMP3 leads to marked decreases in CGRP and AM responses. CH-I had marked effects on responses to AM in the presence of each RAMP, despite appropriate [¹²⁵I]-AM binding and surface delivery. In contrast, CH-I had little effect on CGRP potency or efficacy. CH-II/RAMP1 maintained full CGRP receptor function,

whereas CH-II/RAMP2 and CH-II/RAMP3 exhibited small but significant reductions in CGRP/AM potency. These findings indicate that the CLR ECL3 is crucial for AM signalling via the CGRP, AM₁ and AM₂ receptors, but it is less critical for CGRP signalling via these three receptors. We therefore constructed four mutants (Mut-A, Mut-B, Mut-C and Mut-D) in which CLR ECL3 residues with no homology to VPAC2 were sequentially replaced with alanine (Ala) (Figure 6); the effects of these mutants were characterized following transient transfection.

As with WT-CLR, all four CLR mutants were increased at the cell surface, even when expressed alone (Figure 7A). The cell-surface expression of these four mutants was significantly increased by co-transfection with WT-RAMP1, WT-RAMP2 or WT-RAMP3 to such an extent that their expression frequen-

Table 2

Agonist-induced cAMP response in cells co-expressing RAMP with WT or chimeric CLR

Receptor complex	CGRP pEC ₅₀	E _{max} (pmol)	AM pEC ₅₀	E _{max} (pmol)
RAMP1/WT-CLR	9.70 ± 0.13	361.5 ± 11.1	7.92 ± 0.13	400.7 ± 16.8
RAMP1/CH-7	7.93 ± 0.18*	325.7 ± 18.7	6.29 ± 0.23*	154.3 ± 17.0*
RAMP1/CH-I	9.50 ± 0.22	345.7 ± 18.2	6.84 ± 0.13*	308.3 ± 16.3*
RAMP1/CH-II	9.47 ± 0.13	450.3 ± 14.0*	7.52 ± 0.16	526.6 ± 32.3*
RAMP2/WT-CLR	6.63 ± 0.16	244.9 ± 16.0	8.95 ± 0.22	274.7 ± 18.2
RAMP2/CH-7	5.85 ± 0.13*	57.7 ± 4.4*	7.19 ± 0.26*	51.6 ± 4.9*
RAMP2/CH-I	6.57 ± 0.10	174.6 ± 7.7*	6.93 ± 0.24*	129.5 ± 13.7*
RAMP2/CH-II	6.18 ± 0.16	128.9 ± 11.1*	8.03 ± 0.26*	253.7 ± 21.1
RAMP3/WT-CLR	7.13 ± 0.08	305.5 ± 9.4	9.12 ± 0.10	306.3 ± 8.8
RAMP3/CH-7	5.62 ± 0.10*	281.5 ± 18.5	7.94 ± 0.18*	187.9 ± 10.7*
RAMP3/CH-I	6.46 ± 0.21*	338.7 ± 36.0	7.78 ± 0.16*	218.9 ± 11.0*
RAMP3/CH-II	6.46 ± 0.15*	283.2 ± 18.7	8.56 ± 0.16*	346.6 ± 14.4

* $P < 0.05$ vs. the corresponding RAMP/WT-CLR, one-way ANOVA followed by Dunnett's multiple comparison test.

Data are displayed as the means $\pm$ SEM of five independent experiments.

E_{max} values are the maximum asymptote of concentration-effect curves and correspond to the amount of cAMP.

RAMP, receptor activity-modifying protein; WT, wild-type; CLR, calcitonin receptor-like receptor; CGRP, calcitonin gene-related peptide; AM, adrenomedullin; CH, chimera.




Figure 6

Amino acid sequence alignment of the human CLR chimera CH-7 and alanine substitutions in the region spanning amino acids 395–408. Note that the region contains CLR ECL3 (amino acids 395–406). Bold letters indicate amino acid residues (Pro, Ile and Tyr) conserved between CLR and VPAC2. Residues that are not conserved were sequentially replaced with alanine (A): mutant (Mut)-A (R395A), Mut-B (E397A + G398A + K399A), Mut-C (E402A + E403A + V404A) and Mut-D (D406A + Y407A + I408A).

cies were not significantly different from those of the corresponding V5-CLR/RAMP (Figure 7A). All four untagged mutants markedly increased the translocation of V5-RAMP1, V5-RAMP2 or V5-RAMP3 to a level comparable to that seen with WT-CLR (Figure 7B).

Figure 7C and D shows the specific binding of [125 I]-CGRP and [125 I]-AM to the WT and mutant receptors for CGRP or AM. The specific [125 I]-CGRP binding to Mut-A/RAMP1 was significantly greater than to the WT CGRP receptor, whereas the remaining three mutant receptors

exhibited specific [125 I]-CGRP binding comparable to the WT CGRP receptor (Figure 7C). Likewise, the specific binding of [125 I]-AM to Mut-A/RAMP2 was significantly greater than to the WT AM₁ receptor. The specific binding of [125 I]-AM to the remaining three mutant receptors was similar to the binding to the WT AM₁ receptor (Figure 7D). There were also no significant differences in specific [125 I]-AM binding to the WT and mutant AM₂ receptors (Figure 7D). [125 I]-CGRP and [125 I]-AM competition curves for WT and mutant receptors are shown in Figure 8A–C. The pIC₅₀ values

**Figure 7**

Effect of CLR ECL3 point mutations on cell-surface expression and specific binding of agonists. (A) Cell-surface expression of V5-tagged WT or mutant CLR. WT-CLR or one of four CLR mutants (Mut-A, -B, -C and -D) was transiently transfected into HEK-293 cells, with or without WT-RAMP. The cell-surface expression of each FITC-labelled protein was estimated by flow cytometry. Data are means $\pm$ SEM of five to eight separate experiments. * P < 0.05 vs. corresponding Vector/V5-CLR or -Mut, # P < 0.05 vs. corresponding V5-CLR/WT-RAMP. (B) Changes in the cell-surface expression of V5-RAMP induced by co-transfection with WT or mutant CLR (Mut-A, -B, -C and -D). The indicated constructs were transfected into HEK-293 cells, after which the cell-surface expression of each V5-tagged protein was analysed by flow cytometry. Data are shown as the means $\pm$ SEM of five to nine separate experiments. * P < 0.05 vs. corresponding Vector/V5-RAMP, # P < 0.05 vs. corresponding WT-CLR/V5-RAMP. (C and D) Specific binding of [125 I]-CGRP (C) and [125 I]-AM (D) to heterodimeric receptors composed of WT RAMP and WT or mutant CLR (Mut-A, -B, -C, and -D). HEK-293 cells were transfected with WT-RAMP2 or -3 plus WT-CLR or one of the indicated mutants. Data are shown as the means $\pm$ SEM of five experiments in (C) and five to seven experiments in (D) and are expressed as percentages of CLR/RAMP1 (C) or CLR/RAMP2 (D). * P < 0.05 vs. corresponding CLR/RAMP.

derived from the curves are given in Table 3. When co-expressed with RAMP1, Mut-A (Hill slope = 0.65), Mut-C (Hill slope = 0.69) and Mut-D (Hill slope = 0.57) significantly reduced CGRP affinity, although Mut-B (Hill slope = 0.73) behaved much like WT-CLR (Hill slope = 0.73). These Hill slopes were not different from each other. There were no

significant differences in the affinity of AM for the WT and mutant AM₁ receptors. Their Hill slopes were not different from each other, although the four Hill slopes (0.75 for CLR/RAMP2; 0.85 for Mut-A/RAMP2; 0.80 for Mut-B/RAMP2; 0.77 for Mut-C/RAMP2) differed from 1. On the other hand, Mut-A/RAMP3 significantly increased AM affin-




Figure 8

Effect of CLR ECL3 point mutations on agonist binding affinity and induced cAMP production. (A–C) Competitive binding of [125 I]-CGRP to WT or mutant CLR/RAMP1 (A) and binding of [125 I]-AM to WT or mutant CLR/RAMP2 (B) or CLR/RAMP3 (C). HEK-293 cells were co-transfected with WT-RAMP and WT or mutant CLR (Mut-A, -B, -C and -D). Data are shown as the means $\pm$ SEM of five experiments in (A–C). [125 I]-CGRP (A) or [125 I]-AM binding (B, C); B_0 , total binding in the presence of competing unlabelled CGRP (A) or AM (B, C); N, nonspecific binding (measured in the presence of 10^{-6} M CGRP (A) or 10^{-6} M AM (B, C)). pIC_{50} values are given in Table 3. (D–F) cAMP production elicited by WT or mutant CLR (Mut-A, -B, -C and -D) expressed with WT-RAMP. WT-CLR or one of the four CLR mutants was transiently transfected into HEK-293 cells along with RAMP1 (D), RAMP2 (E) or RAMP3 (F). All transfectants were then simultaneously exposed to the indicated concentrations of CGRP or AM. Data are shown as the means $\pm$ SEM of six experiments in (D) and seven experiments in (E, F). pEC_{50} values are given in Table 3.

ity, but Mut-D/RAMP3 significantly reduced AM affinity. The remaining two mutant receptors exhibited AM affinity comparable to WT AM_2 receptors. Their Hill slopes were different from each other (0.63 for CLR/RAMP3; 0.65 for Mut-A/RAMP3; 0.64 for Mut-B/RAMP3; 0.68 for Mut-C/RAMP3) because Mut-D/RAMP3 had a steeper slope of 1.04, compared to the others.

The functionality of the four CLR mutants co-expressed with RAMP1, RAMP2 or RAMP3 was evaluated by measuring CGRP- or AM-induced cAMP accumulation (Table 3, Figure 8D–F). The potency and efficacy of CGRP for the four mutant CGRP receptors were not significantly different from those for the WT CGRP receptor. Likewise, all four CLR mutants did not significantly change AM potency

and efficacy in the presence of RAMP2 or RAMP3. Among all the WT and mutant receptors, only the Hill slope (1.9 for AM) of the WT AM_1 receptor was different from 1.

Discussion and conclusions

In the present study, we found that exchanging the CLR ECL3 (amino acids 395–406, including the receptor signal peptide) with the corresponding VPAC2 sequence markedly reduced AM-induced cAMP production via the AM_1 and AM_2 receptors, but CGRP-induced cAMP production via the CGRP receptor was unaffected. A very recent study showed the

Table 3

pIC₅₀, pEC₅₀ and E_{max} data for WT and mutant receptors with CGRP or AM

Receptor complex	pIC ₅₀ ± SEM (n)	pEC ₅₀ ± SEM (n)	E _{max} ± SEM (pmol) (n)
RAMP1/WT-CLR	8.51 ± 0.04 (5)	8.90 ± 0.09 (6)	429.1 ± 11.1 (6)
RAMP1/Mut-A	8.19 ± 0.05 (5)*	8.95 ± 0.12 (6)	410.4 ± 15.0 (6)
RAMP1/Mut-B	8.34 ± 0.06 (5)	8.92 ± 0.09 (6)	409.9 ± 10.8 (6)
RAMP1/Mut-C	8.07 ± 0.06 (5)*	8.83 ± 0.11 (6)	438.3 ± 14.8 (6)
RAMP1/Mut-D	7.80 ± 0.08 (5)*	8.54 ± 0.11 (6)	409.0 ± 15.0 (6)
RAMP2/WT-CLR	8.24 ± 0.06 (5)	8.95 ± 0.04 (7)	200.8 ± 4.4 (7)
RAMP2/Mut-A	8.55 ± 0.04 (5)	8.96 ± 0.15 (7)	193.7 ± 8.3 (7)
RAMP2/Mut-B	8.30 ± 0.04 (5)	8.87 ± 0.12 (7)	210.5 ± 7.4 (7)
RAMP2/Mut-C	8.23 ± 0.05 (5)	8.66 ± 0.14 (7)	228.0 ± 10.0 (7)
RAMP2/Mut-D	8.08 ± 0.03 (5)	8.55 ± 0.12 (7)	250.5 ± 9.9 (7)*
RAMP3/WT-CLR	8.13 ± 0.07 (5)	8.85 ± 0.14 (7)	256.2 ± 10.6 (7)
RAMP3/Mut-A	8.58 ± 0.07 (5)*	8.99 ± 0.15 (7)	232.6 ± 10.3 (7)
RAMP3/Mut-B	8.05 ± 0.07 (5)	8.93 ± 0.13 (7)	249.3 ± 9.8 (7)
RAMP3/Mut-C	8.07 ± 0.07 (5)	8.83 ± 0.16 (7)	246.5 ± 12.1 (7)
RAMP3/Mut-D	7.73 ± 0.05 (5)*	8.59 ± 0.11 (7)	274.6 ± 9.2 (7)

*P < 0.05 vs. the corresponding RAMP/WT-CLR, one-way ANOVA followed by Dunnett's multiple comparison test.

For RAMP1-based receptors: competition between CGRP and [¹²⁵I]-CGRP and CGRP responses.

For RAMP2- or RAMP3-based receptors: competition between AM and [¹²⁵I]-AM and AM responses.

AM, adrenomedullin; RAMP, receptor activity-modifying protein; WT, wild-type; CLR, calcitonin receptor-like receptor; Mut, mutant.

effect of single point mutations in the CLR ECL3 on CGRP receptor function in the presence of hRAMP1 (Barwell *et al.*, 2011). Among all the mutant receptors, only I400A/RAMP1 showed a significant reduction (~7-fold) in CGRP potency (Barwell *et al.*, 2011), although this contribution appears to be minor. In addition to I400, P396 and Y405 in CLR are also conserved in hVPAC2. All the remaining mutants, including P396A and Y405A, maintained normal CGRP responses (Barwell *et al.*, 2011). Of our four mutants, only Mut-D (D406A/Y407A/I408A) exhibited a small but significant reduction in CGRP response. Taken together, these findings suggest that the CLR ECL3 is required for appropriate activation of the two AM receptors but is less involved in CGRP receptor activation. To our knowledge, this is the first report showing the critical involvement of ECL3 in the activation of a Family B GPCR, although there is substantial evidence of the importance of ECL3 in Family A GPCR signalling (Lawson and Wheatley, 2004; Claus *et al.*, 2005; Klco *et al.*, 2006; Kleinau *et al.*, 2008; Peeters *et al.*, 2011). In these cases, reduced signalling is due largely to a loss of agonist binding. The general view is that agonist binding to GPCRs triggers a conformational change within the TM receptor core, leading to proper G-protein activation (Wess *et al.*, 2008). Our results showed that the two ECL3 chimeric receptors CH-I/RAMP2 and CH-I/RAMP3 behaved like WT receptors with respect to their specific binding and the binding affinity of [¹²⁵I]-AM. Therefore, it is likely that the CLR ECL3 is mainly involved in AM-induced conformational changes of the AM₁ and AM₂ receptors to induce signal activation.

Among the four mutants targeting residues within the CLR ECL3, only Mut-D showed small but significant decreases

in AM potency, without affecting AM efficacy in the presence of RAMP2 or RAMP3. These results suggest that the nine amino acid residues tested, none of which are conserved in hVPAC2, are not critically involved in the activation of the two AM receptors. There is a small possibility that the remaining three residues (P396, I400 and Y405) are crucially involved in AM receptor activation because they are all conserved in hVPAC2. The hVPAC2 ECL3 shares little sequence identity with that of CLR and lacks two residues present in the CLR ECL3 (Figure 1A). Therefore, it is likely that these two differences between the ECL3 structures lead to marked reductions in AM responses via CH-I/RAMP2 or CH-I/RAMP3. Our results suggest that the entire structure of the CLR ECL3 plays a key role in AM-mediated maximal activation, which is also supported by the findings that the CLR ECL3 is highly conserved among animals despite being a non-TM region (Figure 9A).

It is well known that the receptor/G-protein complex has a higher affinity for agonists than does the free or uncoupled receptor (Zhao *et al.*, 1998). In this study, there were no significant changes in AM affinity for CH-I/RAMP2 or CH-I/RAMP3. In addition, neither of the chimeric receptors decreased their specific [¹²⁵I]-AM binding. These results suggest that the primary role for the ECL3 of CLR may be to control G_s activation, rather than G_s coupling, in the AM₁ and AM₂ receptors. For the AM₁ receptor, the CLR helix 8 (Figure 1B) has been suggested to be a key determinant of G_s-mediated signalling (Kuwasako *et al.*, 2010). Therefore, the entire ECL3 of CLR seems to contribute indirectly to the agonist-induced receptor conformational change for G_s activation.

Notably, CH-I/RAMP1 showed a 12-fold reduction in AM potency without affecting CGRP potency. In addition, CGRP




Figure 9

(A) Comparison of CLR ECL3 sequences from 11 mammals. Non-conserved amino acids are in bold. (B) Comparison of ECL3 sequences among the three human Family B GPCRs that interact with the three RAMPs: CLR, calcitonin receptor (CTR) and VPAC1 receptor. A grey asterisk indicates amino acid residues conserved in CLR. Amino acids conserved among CLR, CTR and VPAC1 are boxed.

potency for CH-I/RAMP2 and CH-I/RAMP3 was affected to a much lesser extent. These results suggest that the CLR ECL3 participates in AM signalling, but not CGRP signalling, via all three WT CLR/RAMP heterodimers. Previous studies showed that the CLR ectodomain (ECD) contains a sequence that contributes to AM binding in the presence of RAMP2; the CLR sequence was not involved in CGRP binding in the presence of RAMP1 (Koller *et al.*, 2002; 2004). The interaction between the CLR ECD and AM is supported by a 'two-domain model', which predicts that in many Family B GPCRs, agonist specificity is primarily associated with the ECD (for the C-terminus of the agonist) and with secondary recognition by a TM domain (for the N-terminus of the agonist) (Siu and Stevens, 2010). So far, there have been few reports showing that RAMPs possess critical sites for the binding of CGRP and AM (Qi and Hay, 2010; Kuwasako *et al.*, 2011b). Rather, the three RAMPs may indirectly alter the binding affinity of CGRP and AM to the ECD and TM of CLR. Taken together, it is possible that CGRP and AM may change the conformation of CLR differently within the same receptor complex. To validate our hypothesis, further work to elucidate the entire crystal structure of the three CLR/RAMP heterodimers in the presence and absence of CGRP/AM binding would be valuable.

In Family A GPCRs, ECL3 has been shown to interact with other extracellular regions (e.g. ECD, ECL2) through hydro-

gen bonds, disulphide bonds, etc. (Peeters *et al.*, 2011). Like Family B GPCRs, RAMPs possess a large ECD. However, none of the CLR ECL3 residues that are not conserved in VPAC2 were found to be significantly involved in interactions with RAMPs because our four CLR ECL3 mutants only slightly affected the surface delivery of RAMPs (Figure 7A and B). Among the CLR ECL3 residues, only P396 is conserved, not only in VPAC2 but also in calcitonin receptor and VPAC1 (Figure 9B). Although the three RAMPs can also interact strongly with calcitonin receptor and VPAC1 (Hay *et al.*, 2006; Sexton *et al.*, 2009), the cell-surface expression of P396A/RAMP1 was not reduced compared with the WT CGRP receptor (Barwell *et al.*, 2011). Therefore, it seems unlikely that P396A interacts with other RAMPs.

In contrast to ECL3, the replacement of CLR TM7 with the corresponding VPAC2 sequence had less of an effect on signalling via the CGRP, AM₁ or AM₂ receptor. The TM7 regions of CLR and VPAC2 are about 40% identical (Figure 1A), which raises the possibility that the nine conserved residues participate in TM7 functions shared by CLR and VPAC2. It was previously reported that, in the human VPAC1 receptor, the binding of Asp⁵ of VIP to Arg¹⁸⁸ in TM2 alters the interaction network between Arg¹⁸⁸ and Asn²²⁹ in TM3 and Gln³⁸⁰ in TM7, leading to G-protein activation (Chugunov *et al.*, 2010). Notably, these three residues are fully conserved among many Family B GPCRs, suggesting that there may be a common mechanism underlying Family B GPCR activation. Consistent with this idea, Glu³⁸⁰ in VPAC1 is strictly conserved in CLR and VPAC2. Additional experiments are needed to confirm whether this activation mechanism also applies to both CLR and VPAC2.

In summary, the human CLR ECL3 is crucial for AM-induced cAMP responses via the three CLR/RAMP heterodimers but is less involved in their CGRP responses. This CLR region probably participates in AM-induced conformational changes of these heterodimers and thereby induces their activation. This study provides a clue to the molecular basis of the activation of RAMP-based Family B GPCRs.

Acknowledgements

This study was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports Science and Technology, Japan and by research grants from the Takeda Science Foundation, the Mochida Memorial Foundation for Medical and Pharmaceutical Research and the Suzuken Memorial Foundation.

Conflict of interest

The authors state that they have no conflict of interest.

References

Bailey RJ, Hay DL (2007). Agonist-dependent consequences of proline to alanine substitution in the transmembrane helices of the calcitonin receptor. *Br J Pharmacol* 151: 678–687.

- Barwell J, Conner A, Poyner DR (2011). Extracellular loops 1 and 3 and their associated transmembrane regions of the calcitonin receptor-like receptor are needed for CGRP receptor function. *Biochim Biophys Acta* 1813: 1906–1916.
- Christopoulos A, Christopoulos G, Morfis M, Udawela M, Laburthe M, Couvineau A *et al.* (2003). Novel receptor partners and function of receptor activity-modifying proteins. *J Biol Chem* 278: 3293–3297.
- Chugunov AO, Simms J, Poyner DR, Dehouck Y, Rومان M, Gilis D *et al.* (2010). Evidence that interaction between conserved residues in transmembrane helices 2, 3, and 7 are crucial for human VPAC1 receptor activation. *Mol Pharmacol* 78: 394–401.
- Claus M, Jaeschke H, Kleinau G, Neumann S, Krause G, Paschke R (2005). A hydrophobic cluster in the center of the third extracellular loop is important for thyrotropin receptor signaling. *Endocrinology* 146: 5197–5203.
- Conner AC, Simms J, Barwell J, Wheatley M, Poyner DR (2007a). Ligand binding and activation of the CGRP receptor. *Biochem Soc Trans* 35 (Pt 4): 729–732.
- Conner M, Hawtin SR, Simms J, Wootten D, Lawson Z, Conner AC *et al.* (2007b). Systematic analysis of the entire second extracellular loop of the V(1a) vasopressin receptor: key residues, conserved throughout a G-protein-coupled receptor family, identified. *J Biol Chem* 282: 17405–17412.
- Flahaut M, Rossier BC, Firsov D (2002). Respective roles of calcitonin receptor-like receptor (CRLR) and receptor activity-modifying proteins (RAMP) in cell surface expression of CRLR/RAMP heterodimeric receptors. *J Biol Chem* 277: 14731–14737.
- Gibbons C, Dackor R, Dunworth W, Fritz-Six K, Caron KM (2007). Receptor activity-modifying proteins: RAMPing up adrenomedullin signaling. *Mol Endocrinol* 21: 783–796.
- Hawtin SR, Simms J, Conner M, Lawson Z, Parslow RA, Trim J *et al.* (2006). Charged extracellular residues, conserved throughout a G-protein-coupled receptor family, are required for ligand binding, receptor activation, and cell-surface expression. *J Biol Chem* 281: 38478–38488.
- Hay DL, Howitt SG, Conner AC, Schindler M, Smith DM, Poyner DR (2003). CL/RAMP2 and CL/RAMP3 produce pharmacologically distinct adrenomedullin receptors: a comparison of effects of adrenomedullin22-52, CGRP8-37 and BIBN4096BS. *Br J Pharmacol* 140: 477–486.
- Hay DL, Poyner DR, Sexton PM (2006). GPCR modulation by RAMPs. *Pharmacol Ther* 109: 173–197.
- Ittner LM, Luessi F, Koller D, Born W, Fischer JA, Muff R (2004). Aspartate(69) of the calcitonin-like receptor is required for its functional expression together with receptor-activity-modifying proteins 1 and -2. *Biochem Biophys Res Commun* 319: 1203–1209.
- Kitamura K, Ichiki Y, Tanaka M, Kawamoto M, Emura J, Sakakibara S *et al.* (1994). Immunoreactive adrenomedullin in human plasma. *FEBS Lett* 341: 288–290.
- Klco JM, Nikiforovich GV, Baranski TJ (2006). Genetic analysis of the first and third extracellular loops of the C5a receptor reveals an essential WXXG motif in the first loop. *J Biol Chem* 281: 12010–12019.
- Kleinau G, Jaeschke H, Mueller S, Raaka BM, Neumann S, Paschke R *et al.* (2008). Evidence for cooperative signal triggering at the extracellular loops of the TSH receptor. *FASEB J* 22: 2798–2808.
- Koller D, Born W, Leuthauser K, Fluhmann B, McKinney RA, Fischer JA *et al.* (2002). The extreme N-terminus of the calcitonin-like receptor contributes to the selective interaction with adrenomedullin or calcitonin gene-related peptide. *FEBS Lett* 531: 464–468.
- Koller D, Ittner LM, Muff R, Husmann K, Fischer JA, Born W (2004). Selective inactivation of adrenomedullin over calcitonin gene-related peptide receptor function by the deletion of amino acids 14–20 of the mouse calcitonin-like receptor. *J Biol Chem* 279: 20387–20391.
- Kuwasako K, Shimekake Y, Masuda M, Nakahara K, Yoshida T, Kitaura M *et al.* (2000). Visualization of the calcitonin receptor-like receptor and its receptor activity-modifying proteins during internalization and recycling. *J Biol Chem* 275: 29602–29609.
- Kuwasako K, Kitamura K, Uemura T, Nagoshi Y, Kato J, Eto T (2003). The function of extracellular cysteines in the human adrenomedullin receptor. *Hypertens Res* 26 (Suppl): S25–S31.
- Kuwasako K, Cao YN, Nagoshi Y, Kitamura K, Eto T (2004a). Adrenomedullin receptors: pharmacological features and possible pathophysiological roles. *Peptides* 25: 2003–2012.
- Kuwasako K, Cao YN, Nagoshi Y, Tsuruda T, Kitamura K, Eto T (2004b). Characterization of the human calcitonin gene-related peptide receptor subtypes associated with receptor activity-modifying proteins. *Mol Pharmacol* 65: 207–213.
- Kuwasako K, Cao YN, Chu CP, Iwatsubo S, Eto T, Kitamura K (2006). Functions of the cytoplasmic tails of the human receptor activity-modifying protein components of calcitonin gene-related peptide and adrenomedullin receptors. *J Biol Chem* 281: 7205–7213.
- Kuwasako K, Kitamura K, Nagata S, Kato J (2008). Functions of the extracellular histidine residues of receptor activity-modifying proteins vary within adrenomedullin receptors. *Biochem Biophys Res Commun* 377: 109–113.
- Kuwasako K, Kitamura K, Nagata S, Kato J (2009). Flow cytometric analysis of the calcitonin receptor-like receptor domains responsible for cell-surface translocation of receptor activity-modifying proteins. *Biochem Biophys Res Commun* 384: 249–254.
- Kuwasako K, Kitamura K, Nagata S, Hikosaka T, Kato J (2010). Function of the cytoplasmic tail of human calcitonin receptor-like receptor in complex with receptor activity-modifying protein 2. *Biochem Biophys Res Commun* 392: 380–385.
- Kuwasako K, Kitamura K, Nagata S, Hikosaka T, Kato J (2011a). Structure-function analysis of helix 8 of human calcitonin receptor-like receptor within the adrenomedullin 1 receptor. *Peptides* 32: 144–149.
- Kuwasako K, Kitamura K, Nagata S, Hikosaka T, Takei Y, Kato J (2011b). Shared and separate functions of the RAMP-based adrenomedullin receptors. *Peptides* 32: 1540–1550.
- Lawson Z, Wheatley M (2004). The third extracellular loop of G-protein-coupled receptors: more than just a linker between two important transmembrane helices. *Biochem Soc Trans* 32 (Pt 6): 1048–1050.
- McLatchie LM, Fraser NJ, Main MJ, Wise A, Brown J, Thompson N *et al.* (1998). RAMPs regulate the transport and ligand specificity of the calcitonin-receptor-like receptor. *Nature* 393: 333–339.
- Muff R, Born W, Fischer JA (2003). Adrenomedullin selectivity of calcitonin-like receptor/receptor activity modifying proteins. *Hypertens Res* 26 (Suppl): S3–S8.

- Nielsen SM, Nielsen LZ, Hjorth SA, Perrin MH, Vale WW (2000). Constitutive activation of tethered-peptide/corticotropin-releasing factor receptor chimeras. *Proc Natl Acad Sci U S A* 97: 10277–10281.
- Peeters MC, van Westen GJ, Li Q, Ijzerman AP (2011). Importance of the extracellular loops in G protein-coupled receptors for ligand recognition and receptor activation. *Trends Pharmacol Sci* 32: 35–42.
- Poyner DR, Sexton PM, Marshall I, Smith DM, Quirion R, Born W *et al.* (2002). International Union of Pharmacology. XXXII. The mammalian calcitonin gene-related peptides, adrenomedullin, amylin, and calcitonin receptors. *Pharmacol Rev* 54: 233–246.
- Qi T, Hay DL (2010). Structure–function relationships of the N-terminus of receptor activity-modifying proteins. *Br J Pharmacol* 159: 1059–1068.
- Runge S, Wulff BS, Madsen K, Brauner-Osborne H, Knudsen LB (2003). Different domains of the glucagon and glucagon-like peptide-1 receptors provide the critical determinants of ligand selectivity. *Br J Pharmacol* 138: 787–794.
- Salvatore CA, Mallee JJ, Bell IM, Zartman CB, Williams TM, Koblan KS *et al.* (2006). Identification and pharmacological characterization of domains involved in binding of CGRP receptor antagonists to the calcitonin-like receptor. *Biochemistry* 45: 1881–1887.
- Sexton PM, Poyner DR, Simms J, Christopoulos A, Hay DL (2009). Modulating receptor function through RAMPs: can they represent drug targets in themselves? *Drug Discov Today* 14: 413–419.
- Simpson LM, Wall ID, Blaney FE, Reynolds CA (2011). Modeling GPCR active state conformations: the beta(2)-adrenergic receptor. *Proteins* 79: 1441–1457.
- Siu FY, Stevens RC (2010). RAMP-ing up class-B GPCR ECD structural coverage. *Structure* 18: 1067–1068.
- Standfuss J, Edwards PC, D'Antona A, Fransen M, Xie G, Oprian DD *et al.* (2011). The structural basis of agonist-induced activation in constitutively active rhodopsin. *Nature* 471: 656–660.
- Topiol S, Sabio M (2009). X-ray structure breakthroughs in the GPCR transmembrane region. *Biochem Pharmacol* 78: 11–20.
- Unson CG, Wu CR, Jiang Y, Yoo B, Cheung C, Sakmar TP *et al.* (2002). Roles of specific extracellular domains of the glucagon receptor in ligand binding and signaling. *Biochemistry* 41: 11795–11803.
- Van Rampelbergh J, Juarranz MG, Perret J, Bondue A, Solano RM, Delporte C *et al.* (2000). Characterization of a novel VPAC(1) selective agonist and identification of the receptor domains implicated in the carboxyl-terminal peptide recognition. *Br J Pharmacol* 130: 819–826.
- Walker CS, Conner AC, Poyner DR, Hay DL (2010). Regulation of signal transduction by calcitonin gene-related peptide receptors. *Trends Pharmacol Sci* 31: 476–483.
- Wess J, Han SJ, Kim SK, Jacobson KA, Li JH (2008). Conformational changes involved in G-protein-coupled-receptor activation. *Trends Pharmacol Sci* 29: 616–625.
- Wheatley M, Wootten D, Conner MT, Simms J, Kendrick R, Logan RT *et al.* (2012). Lifting the lid on G-protein-coupled receptors: the role of extracellular loops. *Br J Pharmacol* 165: 1688–1703.
- Zhao MM, Gaivin RJ, Perez DM (1998). The third extracellular loop of the beta2-adrenergic receptor can modulate receptor/G protein affinity. *Mol Pharmacol* 53: 524–529.