

REVIEW

Lysophosphatidic acid in atherosclerotic diseases

Andreas Schober¹ and Wolfgang Siess²

¹*Institute for Molecular Cardiovascular Research, RWTH Aachen University, Aachen, Germany,*
and ²*Institute for Prevention of Cardiovascular Diseases, University of Munich, Munich,*
Germany

Correspondence

Dr Wolfgang Siess, Institute for Prevention of Cardiovascular Diseases, University of Munich, Pettenkoferstr. 9, 80336 Munich, Germany. E-mail: wsiess@med.uni-muenchen.de

Keywords

Lysophosphatidic acid; LPA receptor; endothelial cells; smooth muscle cells; macrophage; platelet; neointima; hyperlipidaemia; chemokine; atherothrombosis

Received

1 March 2012

Revised

20 April 2012

Accepted

30 April 2012

Lysophosphatidic acid (LPA) is a potent bioactive phospholipid. As many other biological active lipids, LPA is an autacoid: it is formed locally on demand, and it acts locally near its site of synthesis. LPA has a plethora of biological activities on blood cells (platelets, monocytes) and cells of the vessel wall (endothelial cells, smooth muscle cells, macrophages) that are all key players in atherosclerotic and atherothrombotic processes. The specific cellular actions of LPA are determined by its multifaceted molecular structures, the expression of multiple G-protein coupled LPA receptors at the cell surface and their diverse coupling to intracellular signalling pathways. Numerous studies have now shown that LPA has thrombogenic and atherogenic actions. Here, we aim to provide a comprehensive, yet concise, thoughtful and critical review of this exciting research area and to pinpoint potential pharmacological targets for inhibiting thrombogenic and atherogenic activities of LPA. We hope that the review will serve to accelerate knowledge of basic and clinical science, and to foster drug development in the field of LPA and atherosclerotic/atherothrombotic diseases.

Abbreviations

APT1, acyl-protein thioesterase 1; ATX, autotaxin; CCL2, chemokine ligand 2; cPA, cyclic phosphatidic acid; CREB, cAMP response element-binding; Edg, endothelial differentiation gene; Egr, early growth response gene; ESI, electrospray ionization; ICAM-1, intracellular adhesion molecule-1; LC, liquid chromatography; LCAT, lecithin-cholesterol acyltransferase; LDL, low-density lipoprotein; LPA, lysophosphatidic acid; LPC, lysophosphatidylcholine; LPL, lysophospholipids; LPP, lipid phosphate phosphatase; Lp-PLA₂, lipoprotein-associated PLA₂; LPS, lysophosphatidylserine; LYPLA-I, lysophospholipase A-I; lysoPLD, lysophospholipase D; MCP-1, monocyte chemotactic protein-1; MLC, myosin light chain; MM-LDL, minimally modified LDL; mox-LDL, mildly oxidized LDL; MS, mass spectrometry; NVAf, nonvalvular atrial fibrillation; PA, phosphatidic acid; PAF, platelet-activating factor; PAK, p21-activated kinase; PC, phosphatidylcholine; PI3K, phosphoinositide 3-kinase; PLA, phospholipase A; ox-LDL, oxidized LDL; SBI, silent brain infarction; S1P, sphingosine 1-phosphate; SMD, somatomedin domain; SPCs, smooth muscle progenitor cells; sPLA₂, secretory PLA₂; SRF, serum response factor; TF, tissue factor; VCAM-1, vascular cell adhesion molecule-1; VSMC, vascular smooth muscle cells

Introduction

Atherosclerosis is a slowly progressing, multifocal, chronic arterial disease that is characterized by inflammatory and regenerative processes, which lead to matrix remodelling and large lipid deposits. The retention of low-density lipoproteins (LDL) and activation of endothelial cells initiate atherosclerotic lesion formation in the inner layer (intima) of medium-

and large-sized arteries, such as coronary arteries and cerebral arteries, predominantly at predilection sites, where the laminar blood flow is disturbed (Libby *et al.*, 2011; Weber and Noels, 2011). In early lesions, monocytes are recruited to the arterial wall, where they engulf lipids and transform into foam cells. In the advanced stages of lesion formation, a fibroatheroma develops in the intima that is characterized by the extracellular accumulation of lipids and a fibrous cap

consisting of vascular smooth muscle cells (VSMCs) and extracellular matrix proteins. Although patients with atherosclerosis may not develop clinical symptoms, the narrowing of the arterial lumen may compromise the oxygen supply and result in ischaemia of the tissues that are supplied by the affected artery, such as the myocardium or the brain. Furthermore, acute thrombotic occlusion of the arterial lumen can occur following erosion or rupture of atherosclerotic plaques, which leads to platelet activation and fibrin formation. These acute atherothrombotic events are potentially life-threatening sequelae of atherosclerosis as they may lead to myocardial infarction or stroke (Fuster *et al.*, 2005; Thim *et al.*, 2008). Percutaneous interventions, such as coronary stent implantation, can reopen the occluded or narrowed arteries and limit ischaemic symptoms, and are often life-saving. However, such interventions can result in a re-narrowing of the target vessel (restenosis) due to the accumulation of SMCs in the neointima.

Lysophosphatidic acid (LPA) is a phospholipid that mediates a plethora of activities in blood cells and cells of the vessel wall (Tigyi, 2001; Siess, 2002; Siess and Tigyi, 2004; Smyth *et al.*, 2008). Similar to other lipid mediators such as prostaglandins and leukotrienes, LPA acts as an autacoid. After local formation in response to danger signals such as those initiated by vascular injury or inflammation LPA rapidly activates cells in the immediate vicinity (Tokumura, 1995). The action of LPA is mediated by its binding to and activation of surface GPCRs, including the Edg family of LPA receptors LPA₁₋₃, LPA₄ (GPR23), LPA₅ (GPR92), and LPA₆ (P2Y5), the latter three belonging to the purinoceptor cluster (Choi *et al.*, 2010; Chun *et al.*, 2010; Tigyi, 2010). Currently putative LPA GPCRs are GPR87, P2Y10 and GPR35; hence, there are in total nine LPA receptors (Choi *et al.*, 2010; Chun *et al.*, 2010; Tigyi, 2010). In addition to GPCRs, some of the effects of LPA appear to be mediated by the activation of the nuclear receptor PPAR γ , specifically in macrophages and VSMCs (McIntyre *et al.*, 2003; Zhang *et al.*, 2004).

Our review is focused on the role of LPA and its receptors in atherosclerosis and cardiovascular diseases. Since the seminal observations that LPA is generated during mild oxidation of LDL and accumulates in the lipid core of human atherosclerotic plaques linking, for the first time, LPA to atherosclerosis (Siess *et al.*, 1999), the area has evolved dramatically. We aim to cover this research field in the present review. General aspects of LPA receptor functions and their coupling to intracellular signalling pathways (Choi *et al.*, 2010; Tigyi, 2010; Yanagida and Ishii, 2011), of intra- and extracellular LPA formation and metabolism (Nakanaga *et al.*, 2010; Samadi *et al.*, 2011; van Meeteren and Moolenaar, 2007), and of the role of LPA in cardiovascular physiology (Smyth *et al.*, 2008) have been excellently reviewed previously.

LPA generation in vascular diseases

Although the absolute concentration of circulating LPA varies considerably in different assays (Smyth *et al.*, 2008), it is clear that LPA is detectable in plasma in the low μ M range. The generation of circulating LPA essentially requires the action of autotaxin (ATX), a secreted lysophospholipase D, which removes the polar head group from lysophospholipids (LPL),

such as lysophosphatidylcholine (LPC) (Umez-Goto *et al.*, 2002; Tokumura *et al.*, 2002c; Nakanaga *et al.*, 2010). Accordingly, plasma LPA levels of healthy subjects strongly correlate with the serum ATX activity (Hosogaya *et al.*, 2008) and are reduced by 50% in mice heterozygous for ATX (Tanaka *et al.*, 2006; van Meeteren *et al.*, 2006). Moreover, depletion of ATX completely prevents LPA production in serum (Tanaka *et al.*, 2006; Tsuda *et al.*, 2006), and ATX overexpression in transgenic mice increases LPA plasma levels (Pamuklar *et al.*, 2009). Inhibition studies demonstrated that continual ATX activity is required to maintain a steady state concentration of LPA in the circulation (Albers *et al.*, 2010; Gierse *et al.*, 2010; Gupta *et al.*, 2011). A boronic acid-based inhibitor in mice resulted in a rapid decline of LPA within minutes and the orally available ATX antagonist PF-8380 dose-dependently diminished the plasma LPA concentration by maximally 70% within 3 h (Albers *et al.*, 2010; Gierse *et al.*, 2010).

Adipocyte-derived ATX generates almost half of the LPA in plasma (Dusaulcy *et al.*, 2011). In obesity, insulin resistance in adipocytes has been linked to increased expression of ATX, implicating a potential role of LPA in the sequelae of the metabolic syndrome (Boucher *et al.*, 2005). Furthermore, hyperlipidaemia enhances the activity of circulating ATX in rabbits, which leads to enhanced generation of LPA from LPC in serum, and increases plasma LPA and adipose tissue expression of ATX in mice fed a high fat diet (Tokumura *et al.*, 2002a; Dusaulcy *et al.*, 2011).

Elevated circulating LPA levels may have pro-atherogenic effects, because systemic treatment with unsaturated LPAs has been shown to enhance atherosclerosis (Zhou *et al.*, 2011). In humans, plasma LPA levels correlate with LPC, ATX and LDL levels, indicating that hyperlipidaemia may contribute to the generation of LPA in the circulation (Dohi *et al.*, 2012). Moreover, circulating LPC is increased by hyperlipidaemia (Portman *et al.*, 1970; McConnell and Hoefner, 2006; Matsumoto *et al.*, 2007; Schmitz and Ruebsaamen, 2010). Plasma LPA, which is mainly bound to albumin is generated by endothelial lipase and by lecithin-cholesterol acyltransferase (LCAT) secreted from the liver (Wiesner *et al.*, 2009; Schmitz and Ruebsaamen, 2010). Prolonged (10–50 h) incubation of plasma at 37°C leads to a large increase in LPA from 0.5 to 15 μ M that is due to LPC generation by LCAT and not by lipoprotein-associated (Lp-) PLA₂ activity (Aoki *et al.*, 2002).

Progression of atherosclerosis is associated with conditions of chronic inflammation and oxidant stress. Under these conditions, circulating LPC can be generated by group IIA secretory PLA₂ (sPLA₂), which is an acute-phase reactant, and by Lp-PLA₂ (identical to platelet-activating factor (PAF) acetylhydrolase) (Stafforini, 2009; Rosenson, 2010). sPLA₂ acts on lipoproteins and microvesicles (Fourcade *et al.*, 1995), whereas Lp-PLA₂ specifically catalyses the removal of the acyl group at the *sn*-2 position of oxidatively truncated phospholipids present in oxidized LDL (ox-LDL) (Steinbrecher *et al.*, 1984). LDL particles oxidized to different degrees have been detected at increased levels in the circulation of patients with acute coronary syndromes (Holvoet *et al.*, 1998). Furthermore, electronegative LDL particles, which are pro-inflammatory by releasing chemokines from endothelial cells, are increased and physicochemically heterogeneous in familial hypercholesterolaemic patients (Sanchez-Quesada *et al.*, 2002; 2003). Moreover, circulating levels and enzymatic

activity of Lp-PLA₂ and sPLA₂ have been found to predict cardiovascular events (Packard *et al.*, 2000; Koenig and Khuseynova, 2009; Thompson *et al.*, 2010). Lp-PLA₂ is primarily bound to electronegative LDL in the circulation, and the levels of Lp-PLA₂ positively correlate with the concentration of LDL (Packard *et al.*, 2000; Benitez *et al.*, 2003; Albert *et al.*, 2005). Of note, the Lp-PLA₂ inhibitor darapladib, which has currently been tested in two large cardiovascular outcome trials, limits atherosclerosis in mice (Charo and Taub, 2011; Wang *et al.*, 2011). The sPLA₂ inhibitor varespladib reduces atherosclerosis in mice and a phase III cardiovascular outcome trial on the effects of varespladib is ongoing (Charo and Taub, 2011).

Although it remains to be determined whether pro-atherogenic effects of Lp-PLA₂ and sPLA₂ are related to increased LPA production, a very recent study showing a correlation between levels of Lp-PLA₂, LPCs, LPA and pro-inflammatory cytokines in human plaques supports such a link (Goncalves *et al.*, 2012). Therefore, it can also be assumed that LPC levels elevated in hyperlipidaemia may lead to increased ATX-dependent LPA generation in the circulation. An interesting question in this regard is whether mice deficient in sPLA₂ or Lp-PLA₂ show decreased cellular LPA production or decreased LPA plasma levels. However, apart from one study that shows normal LPA generation of activated washed platelet suspensions prepared from mouse strains that are deficient in sPLA₂ expression (such as the C57BL/6 and C3H strains) (le Balle *et al.*, 1999), nothing is known about this topic.

LPA accumulates in human and mouse atherosclerotic lesions (Siess *et al.*, 1999; Bot *et al.*, 2010). LPA in atherosclerotic lesions is most likely to be derived from ATX-mediated hydrolysis of LPC. Although the expression of ATX in atherosclerotic lesions has not been studied, it has been shown that ATX can be secreted from arterial endothelial cells and that it is up-regulated in the arterial wall following vascular injury (Panchatcharam *et al.*, 2008; Zhou *et al.*, 2011). Structural analysis has suggested that ATX is capable of directing the LPA produced to the cognate GPCRs via binding of its somatomedin domains (SMD) to β 3-integrins, thus contributing to localized LPA formation (Fulkerson *et al.*, 2011; Hausmann *et al.*, 2011; Nishimasu *et al.*, 2011). However, it remains to be determined whether local LPA production and delivery by ATX via binding to β 3-integrins, which are highly expressed in various cell types of atherosclerotic lesions, contributes to LPA formation in the lesion (Hoshiga *et al.*, 1995).

In atherosclerotic lesions, the ATX substrate LPC is generated mainly by LDL oxidation. Polyunsaturated fatty acids in the *sn*-2 position of phosphatidylcholine (PC) in the outer layer of the LDL particle undergo oxidative fragmentation to oxidized short-chain fatty acids (Siess, 2006). The oxidized PC molecules are then specifically hydrolyzed by Lp-PLA₂, which produces LPC (Steinbrecher *et al.*, 1984). LPC in the vessel wall accumulates during hyperlipidaemia, and LPC accumulation in a mouse atherosclerotic model precedes the progressive accumulation of LPA in atherosclerotic tissue (Portman *et al.*, 1970; Bot *et al.*, 2010).

In addition, LPA in atherosclerotic lesions might derive indirectly from microparticles or microvesicles that are shed from activated and apoptotic cells (e.g. macrophages and VSMCs) and accumulate in inflamed atherosclerotic lesions.

Membrane phospholipids of microvesicles, but not intact cells have been reported to be accessible to hydrolysis by sPLA₂, and the concentration of lysophospholipids was found to be increased in microvesicles isolated from inflammatory fluids (Fourcade *et al.*, 1995). Thus, this enzyme could also generate lysophospholipids from microparticles for ATX-dependent LPA formation in inflamed atherosclerotic lesions, although the role of microparticles in the production of lesional LPA remains to be determined.

Previous reports have described increased formation of LPA, preferentially of the alkyl-LPA species, during mild oxidation of LDL (Siess *et al.*, 1999; Zhang *et al.*, 2004), suggesting that a pathway of LPA formation during LDL oxidation may exist, which may be ATX-independent. However, the possibility that ATX, similar to other plasma proteins, might also be associated with LDL (Hoofnagle and Heinecke, 2009) cannot be excluded, although direct evidence is lacking.

Platelet activation importantly contributes to LPA formation in blood. Platelet depletion reduces the serum LPA levels by 50% in rats (Aoki *et al.*, 2002). During blood clotting, the LPA concentration rises from low micromolar plasma concentrations (0.1–1 μ M) up to 10 μ M in serum predominantly by an increase of acyl-LPA species containing polyunsaturated fatty acids (mainly 18:2 and 20:4 fatty acids) (Siess, 2002; Tigyi, 2010). Activation of isolated platelets induces the formation of intracellular LPA via a pathway that involves the PLC-catalysed hydrolysis of phosphoinositides leading to the formation of diacylglycerol that is rapidly phosphorylated by diglyceride kinase to phosphatidic acid (PA) (Siess, 2002). Subsequently, degradation by platelet PLA₂ and PLA₁ enzymes specific for PA leads to the formation of 1-acyl-LPA and 2-acyl-LPA, respectively (Billah *et al.*, 1981; Gaits *et al.*, 1997). Intracellularly produced LPA is not released into the extracellular medium (Watson *et al.*, 1985). LPA in the medium of activated platelets can only be detected in the presence of albumin (Eichholtz *et al.*, 1993). Recent studies indicate that the small amounts of extracellular LPA that can be detected upon activation of isolated platelets are due to the activity of platelet-bound ATX (Pamuklar *et al.*, 2009; Fulkerson *et al.*, 2011). Thus, it seems likely that extracellular LPA detected upon activation of washed platelets is formed by platelet-bound ATX from LPC, and albumin by binding platelet-released LPC plays an essential role in this process.

In contrast to activation of platelets in buffer, platelet activation in the presence of plasma leads to a large increase in extracellular LPA. This occurs by a multistep process involving the generation of LPL via intracellular PLAs and secreted PLA₁ and/or PLA₂ enzymes (Aoki *et al.*, 2002; Sano *et al.*, 2002). Very recent studies demonstrating the binding of ATX to β 3-integrins of activated platelets (Fulkerson *et al.*, 2011) and the identification of a new PLA enzyme secreted from platelets (Bolen *et al.*, 2011) unravel how platelet activation may lead to the large increase in certain molecular LPA species (18:2-acyl-LPA and 20:4 acyl-LPA) in serum. The platelet PLA enzyme that may feed plasma ATX with LPLs upon platelet activation is acyl-protein thioesterase 1 (APT1), also known as lysophospholipase A-I (LYPLA-I) (Bolen *et al.*, 2011). APT1 is released from activated platelets, acts like PLA₁ and generates a pool of *sn*-2-esterified lysophospholipids containing mainly C18:2 and C20:4 (Bolen *et al.*, 2011). The thermodynamically unstable *sn*-2-esterified

lysophospholipids undergo acyl migration resulting in *sn*-1-esterified lysophospholipids, which are the preferred substrates of ATX (Bolen *et al.*, 2011). Subsequently, ATX bound to activated β 3-integrins on the activated platelet membrane via its two SMD produces LPA species containing mainly 18:2 and 20:4 fatty acids (Bolen *et al.*, 2011; Fulkerson *et al.*, 2011). This platelet-dependent LPA production may be crucial for vascular repair, for example, after erosion of atherosclerotic plaques or after stent implantation, where activated platelets rapidly adhere to the denuded vascular surface.

Extracellular LPA is degraded by the ecto-activities of lipid phosphate phosphatases (LPPs) (Morris *et al.*, 2009; Samadi *et al.*, 2011). LPPs hydrolyze a large variety of bioactive lipid phosphates and pyrophosphates; their active site resides on the outer surface of plasma membranes. Three different mammalian LPP isoforms are known. Whereas i.v. injected LPA has a half-life of only 3 min in mice, an extended half-life of circulating LPA up to 12 min and increased levels of plasma LPA have been found in LPP1-deficient mice (Tomsig *et al.*, 2009). It is therefore likely that a balance between ATX and LPP activities controls the concentration of extracellular LPA and thus LPA receptor activation in a dynamic manner (Samadi *et al.*, 2011).

In addition to LPA, cyclic phosphatidic acid (1-acyl-*sn*-glycerol-2,3-cyclic phosphate; cPA) probably formed by ATX, has been identified in human serum (Kobayashi *et al.*, 1999; Tsuda *et al.*, 2006; Fujiwara, 2008; Shan *et al.*, 2008b). cPA is an LPA analogue in which the *sn*-2 hydroxyl group has formed a ring structure with the *sn*-3 phosphate. Although the function of extracellular cPA on vascular cells and blood cells is largely unknown (Fujiwara, 2008; Tigyi, 2010), intracellular cPA, formed by PLD activation, functions as inhibitor of PPAR γ and might thereby modify atherosclerotic processes (Tsukahara *et al.*, 2010).

Measurement of plasma LPA and circulating LPA levels in cardio- and cerebrovascular clinical studies

The two primary methods that are used to measure LPA in plasma are liquid chromatography-electrospray ionization tandem MS (LC-ESI-tandem MS) and enzymatic assays. Two enzymatic methods have been described: (i) a radioenzymatic assay uses recombinant LPA acid acyltransferase and a radioactive fatty acid to measure the formation of radiolabeled PA (Saulnier-Blache *et al.*, 2000); and (b) an enzymatic cycling assay uses lysophospholipase, which generates glycerol 3-phosphate, followed by enzymatic cycling using glycerol 3-phosphate oxidase and glycerol 3-phosphate dehydrogenase. The amplified concentrations of hydrogen peroxide are then measured colorimetrically (Kishimoto *et al.*, 2003). Both of these methods yielded LPA plasma levels (0.07–0.1 μ M) that were 10-fold lower than the levels measured by the LC/MS/MS method (0.7–1 μ M) (Baker *et al.*, 2001; Liebisch and Scherer, 2012).

The determination of plasma LPA is not trivial, and the conditions of sample preparation are important for accurate measurement. The majority of LPA in the circulation is bound to albumin (Tigyi *et al.*, 1991; Tigyi and Miledi, 1992), and

acidic extraction methods are required. The addition of strong acids may lead to conversion of plasma LPC to LPA in plasma samples, leading to an artificial increase in LPA (Shan *et al.*, 2008a; Liebisch and Scherer, 2012). This is important because plasma LPC levels are approximately 100–1000 times higher than that of LPA. To circumvent this problem, the butanolic extraction procedure (pH 4) described by Baker *et al.* is often the method of choice for LPA extraction (Baker *et al.*, 2001; Liebisch and Scherer, 2012). Furthermore, direct mass spectrometric analysis of crude lipid extracts should be avoided when determining the levels of plasma LPA. In-source fragmentation of LPC and lysophosphatidylserine (LPS) to LPA has been reported after direct flow injection during plasma LPA analysis (Zhao and Xu, 2009; Liebisch and Scherer, 2012). LPC can lose its choline group at the ion source before the parent ions are detected, giving rise to signals that are indistinguishable from endogenous LPA.

Activated platelets also contribute significantly to LPA generation in the blood, which is an additional factor that makes it difficult to accurately determine the circulating LPA levels. Platelets are activated easily after venipuncture, and it is impossible to immediately inhibit platelet activation before the blood comes into contact with the anticoagulant and additional platelet inhibitors that are perhaps present in the blood collection cuvette. Similar problems have previously been encountered when determining the concentration of circulating thromboxane and other substances that are released from activated platelets (FitzGerald *et al.*, 1987). However, platelet activation during the subsequent handling of the blood such as during the centrifugation of blood to obtain plasma can be controlled. The lowest plasma concentrations of LPA (approximately 0.1 μ M, measured by the enzymatic cycling assay) (Kishimoto *et al.*, 2003) and LPC (approximately 190 μ M) were found when blood was drawn into 7.5 mM EDTA plus a mixture of 10% (v v⁻¹) citrate, theophylline, adenosine and dipyridamole (Nakamura *et al.*, 2007). The latter three substances are platelet inhibitors. In blood samples that were collected in this manner, the LPA concentration was significantly higher in women (0.1 μ M) than in men (0.077 μ M), and a positive correlation between the plasma LPA concentration and serum lysophospholipase D (lysoPLD) activity was found. However, the LPA concentration could be correlated with the plasma LPC concentration only in men (Hosogaya *et al.*, 2008).

In most of the clinical studies that report LPA plasma levels, no precautions have been taken to control for the artificial contribution of LPA production from activated platelets *in vitro* during blood handling and processing. For future studies, the inclusion of a pharmacological ATX inhibitor would be ideal to circumvent this problem. After i.v. injection of a boronic acid-based inhibitor of ATX into mice, a rapid decrease in the total plasma LPA concentration was observed; this finding was interpreted to indicate that there is a dynamic turnover of LPA in the circulation (Albers *et al.*, 2010). Another explanation for these results could be that the injected ATX inhibitor attenuated the *in vitro* formation of LPA during blood handling.

In a recent cross-sectional study of consecutive patients, it was reported that patients with acute coronary syndrome have significantly increased plasma LPA levels (0.54 μ M) compared with patients with stable angina pectoris (0.36 μ M)

or angiographically normal coronary arteries (0.41 μM) (Dohi *et al.*, 2012). In this study, LPA was measured enzymatically as previously described (Kishimoto *et al.*, 2003); however, special precautions to inhibit LPA formation during blood handling *in vitro* (Nakamura *et al.*, 2007) were not taken. The blood was collected into EDTA-containing tubes (Dohi *et al.*, 2012), and the LPA plasma levels of the patients with angiographically normal coronary arteries (approximately 0.4 μM) were approximately four times higher than the LPA plasma levels that have been reported previously in healthy people. The interpretation of this clinical study is therefore difficult as patients with unstable angina pectoris might exhibit higher LPA plasma levels because their platelets are more easily activated *in vitro*.

Another clinical study investigated the influence of acetylsalicylate on plasma LPA levels in patients with ischaemic cerebral vascular diseases (Li *et al.*, 2008). Elevated LPA levels were found in patients with ischaemic cerebrovascular disease (3.11 μM) compared with healthy controls (1.77 μM). Daily administration of aspirin (100 mg) for 1 month significantly lowered the LPA levels in the patients from 4.06 to 2.41 μM . The authors concluded that their findings support a close association between increased plasma LPA levels and platelet activation. Again, it is difficult to interpret whether the effect of aspirin on the plasma LPA levels occurred *in vivo* or *in vitro*.

A separate clinical study investigated, whether there is a relationship between LPA levels and the prevalence of silent brain infarction (SBI) in patients with non-valvular atrial fibrillation (NVAF) (Li *et al.*, 2010). The plasma LPA levels in the NVAF patients with SBI were significantly higher than those in the control patients ($P < 0.01$) or the NVAF patients without SBI. The authors suggested that LPA might be a novel marker for estimating the status of platelet activation and the risk for SBI onset in NVAF patients.

The question when interpreting all of these studies is whether the plasma LPA that is measured truly reflects the circulating LPA concentration or whether it was formed *in vitro* from platelets that were activated during blood handling. In future studies, the addition of a pharmacological ATX inhibitor to the blood collection tube is recommended to minimize LPA formation during blood handling. Moreover, the measurement of total plasma LPA levels may not be sufficient to predict the cardiovascular risk, because the thrombogenicity and atherogenicity of LPA crucially depends on the type of bondage of the fatty acyl chain to the glycerol backbone (ester or ether) and the saturation of the fatty acid; alkyl-LPA species are more potent platelet activators than the corresponding acyl-LPA species (Simon *et al.*, 1982; Tokumura *et al.*, 2002d; Rother *et al.*, 2003), and only unsaturated acyl-LPA species are atherogenic (Yoshida *et al.*, 2003; Zhang *et al.*, 2004; Zhou *et al.*, 2011). In line with these findings, elevated levels of LPA during pregnancy, which is not associated with increased cardiovascular risk, is due to a rise predominantly of saturated LPA, such as LPA16:0 (Tokumura *et al.*, 2002b). Furthermore, increased circulating LPA has been found in patients with chronic hepatitis C and liver fibrosis and in experimentally induced liver fibrosis (Watanabe *et al.*, 2007a,b). In contrast to hepatitis C infection, which is associated with increased cardiovascular risk, liver cirrhosis does not lead to accelerated atherosclerosis (Petta *et al.*, 2011;

Purnak *et al.*, 2011; Adinolfi *et al.*, 2012). Unfortunately, it is currently unknown which LPA species are elevated in the different forms of liver diseases.

Further studies on the circulating levels of the various LPA species may help to clarify the role of elevated circulating LPA levels in atherogenesis. These could also give a hint for the origin of LPA, because circulating LPA might derive from activated platelets (which produce mainly 18:2-acyl-LPA and 20:4 acyl-LPA), microparticles originating from damaged or apoptotic cells (their LPA species are unknown) or other cell types (adipocytes). However, in general, the relevance of dosing LPA in plasma could be questioned, because LPA is mainly generated locally in the circulation and acts locally.

Effects of LPA on blood and vascular cells

LPA has multiple effects on blood cells and cells of the vessel wall (Siess, 2002; Smyth *et al.*, 2008). In platelets, it induces directly shape change, and it stimulates platelet aggregation and secretion only in synergy with other platelet stimuli (Rother *et al.*, 2003; Haseruck *et al.*, 2004). In human monocytes, LPA increases cytosolic Ca^{2+} (Fueller *et al.*, 2003), and in macrophages, it stimulates cell survival and ox-LDL uptake (Koh *et al.*, 1998; McIntyre *et al.*, 2003). In endothelial cells, LPA stimulates cell migration (Panetti *et al.*, 2000; 2004; Ptaszynska *et al.*, 2010), chemokine secretion (Lin *et al.*, 2006; Zhou *et al.*, 2011), adhesion molecule expression (Rizza *et al.*, 1999; Zhou *et al.*, 2011), actin stress fibre formation and cell contraction (Siess *et al.*, 1999; Hirakawa *et al.*, 2004). In confluent endothelial cells *in vitro* or in endothelium *in vivo*, LPA can either increase (van Nieuw Amerongen *et al.*, 2000; Hirase *et al.*, 2001; Sarker *et al.*, 2010) or decrease (Alexander *et al.*, 1998; Minnear *et al.*, 2001) endothelial permeability. LPA also co-operates with VEGF to stimulate angiogenesis (Tanaka *et al.*, 2006; van Meeteren *et al.*, 2006; Ptaszynska *et al.*, 2010). In VSMCs, LPA stimulates cell contraction, leading to increased vascular tone, cell migration, and proliferation [for ref. see (Siess, 2002; Smyth *et al.*, 2008)].

The specific cellular effects of LPA depend on the species of the LPA molecule, and the origin and the vascular environment of the target cells. There are multiple molecular species of LPA that have been described in biological fluids and that are present in plasma and the lipid-rich core of atherosclerotic plaques (Baker *et al.*, 2001; Rother *et al.*, 2003). The fatty acid is mainly attached at the *sn*-1 position of the glycerol backbone via an ester (1-acyl-LPA) or ether bond (1-alkyl-LPA, 1-alkenyl LPA). The fatty acids, which are of predominantly 16-, 18- and 20-carbon lengths, can be saturated, mono- or poly-unsaturated. Additionally, 2-acyl regioisomers of LPA have been detected; however, *sn*-2-esterified lysophospholipids are unstable and undergo acyl migration to yield *sn*-1-esterified lysophospholipids (Bolen *et al.*, 2011).

The types of LPA molecules that are present determine, qualitatively or quantitatively, the biological response of the cell. For example, acyl-LPA that contain unsaturated but not saturated fatty acid causes the phenotypic dedifferentiation of cultured VSMCs and elicits neointima formation in a non-injury model of the rat carotid artery (Hayashi *et al.*, 2001;

Yoshida *et al.*, 2003; Zhang *et al.*, 2004; Subramanian *et al.*, 2010). Furthermore, LPA (20:4), but not LPA (18:0), triggers the adhesion of monocytes to the vessel wall and enhances the progression of atherosclerosis (Zhou *et al.*, 2011). In platelets, alkyl-LPA (16:0) and acyl-LPA (20:4) are approximately 20- and 7-fold more potent than acyl-LPA (16:0), respectively (Tokumura *et al.*, 2002d; Rother *et al.*, 2003; Haseruck *et al.*, 2004). In cells that express individual recombinant LPA receptors, most of the receptors show an agonist preference for mono- or poly-unsaturated acyl-LPA over saturated acyl-LPA (Tigyi, 2010). However, whether endogenous LPA receptors, which are expressed at much lower levels in mammalian cells, behave similarly is not known.

The type of LPA response can also vary between species. For example, human and cat platelets, but not rodent (rat and mouse) platelets, are activated by LPA (Schumacher *et al.*, 1979; Tokumura *et al.*, 1981). Interestingly, mouse platelets are inhibited by LPA (Pamuklar *et al.*, 2009). Additionally, the type of vascular bed is important in considering the effect of LPA on endothelial cells because the endothelium shows a pronounced heterogeneity. There are varied biomechanical and biochemical inputs along the vasculature that alter the properties of the endothelium, and therefore its response to the same pathophysiological stimuli will be different (Aird, 2005). Thus, *in vivo* studies on the action of LPA on endothelial cells will be particularly important as these allow the study of endothelial cells in healthy and atherosclerotic arterial vessels *in situ* (Zhou *et al.*, 2011). In contrast, *in vitro* culture of endothelial cells cannot preserve their vascular bed-specific phenotype, and cell culture also leads to a loss of glycocalyx on the cell surface, which is an important endothelial interface with circulating blood (Chappell *et al.*, 2009).

An important consideration when investigating the activity of circulating LPA is that plasma albumin might limit the activity of LPA. Albumin binds LPA with a stoichiometry of 3 mol of LPA to 1 mol of albumin (Tigyi and Miledi, 1992; Thumser *et al.*, 1994), and albumin dose-dependently inhibits LPA-induced platelet shape change and aggregation (Tokumura *et al.*, 2002d; Haseruck *et al.*, 2004; Khandoga *et al.*, 2008). This finding explains the observation that approximately 1000-fold higher LPA concentrations (0.5–20 μM) are required to stimulate platelets in plasma and blood, as compared with the stimulation of washed platelet suspensions. It is therefore unlikely that the low LPA concentrations found in plasma (total LPA: 0.1–1 μM) affect circulating platelets, and these low levels may also not affect other blood cells or endothelial cells as well.

For many of the effects of LPA on blood cells and cells of the vessel wall, the LPA receptors that are responsible have not been identified. The availability of mice that are deficient for individual LPA receptors (Choi *et al.*, 2010) has advanced our knowledge of the involvement of specific LPA receptors in atherosclerotic processes (Panchatcharam *et al.*, 2008). Often, *in vitro* studies of LPA receptor expression in individual cells have to rely on quantitative PCR analysis of mRNA transcripts because the possibly low endogenous expression of LPA receptor proteins is difficult to determine by immunoblot with specific LPA-receptor antibodies. Furthermore, the fact that one cell type often expresses many different LPA receptors, and that a single LPA receptor can couple to differ-

ent G-proteins in the same cell (Choi *et al.*, 2010), thereby stimulating a complex LPA signalling network, makes it difficult to dissect the function of individual LPA receptors. However, studies using the siRNA technology to down-regulate individual LPA receptors in vascular cells, blood cells or isolated arteries are helpful to assign a specific LPA receptor to its cellular or vascular effect (Subramanian *et al.*, 2010; Khandoga *et al.*, 2011; Zhou *et al.*, 2011).

LPA in acute atherothrombosis

Myocardial infarction and ischaemic stroke are leading causes of morbidity and mortality in humans. The trigger for approximately 70% of myocardial infarctions is plaque rupture, and the erosion of vulnerable atherosclerotic plaques accounts for the other 30%. These events lead to the exposure of thrombogenic plaque material to circulating blood (Fernandez-Ortiz *et al.*, 1994; van Zanten *et al.*, 1994; Toschi *et al.*, 1997; Kolodgie *et al.*, 2004). Subsequent platelet activation and fibrin formation can lead to the development of an occluding thrombus with possibly fatal consequences for the patient. LPA may play different roles in thrombus formation after erosion and plaque rupture. After plaque erosion, the main thrombogenic stimulus is possibly subendothelial versican-hyaluronan matrix (Kolodgie *et al.*, 2004), whereas after plaque rupture, collagen of the disrupted cap and possibly LPA of the plaque core are the platelet stimuli (van Zanten *et al.*, 1994; Siess *et al.*, 1999; Corti and Badimon, 2002; Rother *et al.*, 2003; Penz *et al.*, 2005; Nakanaga *et al.*, 2010). The lipid core contains alkyl-LPA and acyl-LPA species with high platelet-activating potency (Rother *et al.*, 2003) (Figure 1). On the other hand, LPA locally formed by ATX bound to activated platelets that cover eroded plaques might, in concert with other platelet-derived mediators such as sphingosine 1-phosphate (S1P) and VEGF, activate neighbouring endothelial cells to migrate and proliferate, and thereby help in healing the endothelial defect (Panetti *et al.*, 2000; 2004; Ptaszynska *et al.*, 2010).

LPA that is produced by activated platelets after plaque erosion or rupture may play a role as a positive feedback mediator of platelet activation. However, in a previous study, no inhibitory effects of LPA receptor antagonists were found on aggregation of washed platelets stimulated by collagen or thrombin, indicating that the small amounts of LPA that were generated by stimulated, washed platelets do not mediate or support stimulus-induced platelet aggregation (Haseruck *et al.*, 2004). Whether LPA plays a role as a positive feedback mediator of platelet activation in stimulated blood where ATX-dependent LPA formation rises drastically is not known.

Thrombin activation of isolated platelets leads to the activation of the integrin $\alpha\text{IIb}\beta 3$ and binding of extracellular ATX to the $\beta 3$ integrin through its tandem somatomedin B (SMB) domains (Fulkerson *et al.*, 2011; Hausmann *et al.*, 2011). Echistatin, an arginine-glycine-aspartic acid (RGD)-containing peptide, and a monoclonal antibody against integrin $\alpha\text{IIb}\beta 3$ (10E5) reduced platelet adhesion to ATX and inhibited ATX-mediated LPA formation by thrombin-activated, isolated platelets (Fulkerson *et al.*, 2011). However, the RGD sequence of the tandem SMB domains is not involved in the binding of ATX to the $\beta 3$ integrin of activated




Figure 1

LPA-induced platelet signalling during platelet shape change. Activation of the LPA_5 receptor coupled to the heterotrimeric G_{13} protein stimulates Rho and Rho-kinase. The subsequent bifurcating pathway directed to either the myosin-binding subunit of MLC phosphatase or the LIM-kinase 1 leads to enhanced phosphorylation of MLC and stimulation of phospho-cofilin turnover, respectively. Phosphorylated myosin develops actin-activated ATPase activity, interacts with F-actin, and assembles into filaments, whereas cofilin regulates actin dynamics by enhancing both actin-polymerization and actin filament severing. These cytoskeleton changes underlie the folding of the surface membrane, the formation of pseudopods and the contractile wave centralizing the secretory granules during platelet shape change.

platelets, as demonstrated by an ATX-RGE mutant that showed a similar binding to activated platelets as that of wild-type ATX (Hausmann *et al.*, 2011). Although it has not been tested directly, it is likely that LPA formation in blood also depends on binding of ATX to the β_3 integrins of activated platelets. In blood, ATX, which is present at a concentration of approximately 100 nM in plasma (Nakamura *et al.*, 2008), has to compete with the much higher plasma concentration of fibrinogen (5–8 μ M) for binding to the β_3 integrin on activated platelets.

Healthy people display individual heterogeneity of the LPA-induced platelet aggregation response when measured in washed platelets, PRP or blood (Haseruck *et al.*, 2004; Pamuklar *et al.*, 2008). However, all of the blood donors showed a similar shape change response when tested in blood (Haseruck *et al.*, 2004). LPA-induced platelet aggregation was completely dependent on ADP-mediated activation of $P2Y_1$ and $P2Y_{12}$ receptors in whole blood (Haseruck *et al.*, 2004). Thus, LPA does not directly induce human platelet aggregation in blood, as LPA-induced platelet aggregation requires the presence of extracellular ADP. In stirred blood, ADP may be either secreted from platelet-dense granules or released from red cells. In a recent study using isolated platelets, the individual heterogeneity of LPA-induced platelet aggregation was hypothesized to be due to LPA-induced activation of an inhibitory pathway in the non-responders (Pamuklar *et al.*, 2008).

Low nM concentrations of LPA directly induce shape change in washed platelets (Siess *et al.*, 1999; Rother *et al.*, 2003). LPA binds to GPCRs on the platelet surface, and the signal that is emitted by the activated platelet receptor is transduced by the heterotrimeric G_{13} protein to activate Rho and Rho-kinase (Bauer *et al.*, 1999; Klages *et al.*, 1999; Gratacap *et al.*, 2001; Moers *et al.*, 2003). Rho-kinase phosphorylates the 130-kD myosin-binding subunit of myosin phosphatase, thereby decreasing the activity of myosin phosphatase and increasing myosin light chain (MLC) phosphorylation (Kimura *et al.*, 1996; Bauer *et al.*, 1999; Retzer and Essler, 2000). Rho-kinase also phosphorylates and activates LIM-kinase 1, which increases the turnover of phospho-cofilin (Pandey *et al.*, 2006; 2007). These biochemical pathways converge in the remodelling of actin–myosin structures that underlie platelet shape change (Figure 1). Cytosolic Ca^{2+} increase, Rac activation and p21-activated kinase (PAK) are not involved in LPA-induced shape change (Maschberger *et al.*, 2000; Pandey *et al.*, 2007). High concentrations of LPA (10 μ M) induce an increase in cytosolic Ca^{2+} in washed platelets that is primarily due to the stimulation of Ca^{2+} entry. This leads to cofilin dephosphorylation and secretion; the latter response requires, in addition, integrin $\alpha IIb\beta_3$ outside-in signalling (Maschberger *et al.*, 2000; Pandey *et al.*, 2007). LPA does not activate the heterotrimeric G-protein G_i in platelets, yet it shows a strong synergism in the induction of platelet aggregation with platelet stimuli such as adrenaline and ADP that activate G_i (Rother *et al.*, 2003; Haseruck *et al.*, 2004).

Until recently, the identity of the LPA receptor that mediates platelet shape change remained obscure. Human platelets express mRNA for the Edg receptors LPA_{1-3} , and the purinergic cluster LPA_4 (GPR23), LPA_5 (GPR92), LPA_6 ($P2Y_5$), and the putative receptors GPR87 and $P2Y_{10}$ (Amisten *et al.*, 2008; Khandoga *et al.*, 2008; Pamuklar *et al.*, 2008). The receptors that are most abundantly expressed at the mRNA level are LPA_4 and LPA_5 (Amisten *et al.*, 2008; Khandoga *et al.*, 2008). A previous study suggested that LPA_1 and LPA_3 play a role in inducing platelet shape change based on the ability of the LPA receptor subtype-specific antagonist diacylglycerol pyrophosphate to inhibit LPA-induced platelet shape change. However, the inhibitory effect of this compound increased with the time of pre-incubation (Rother *et al.*, 2003), which is not typical for a receptor antagonist. Two recent studies described the involvement of LPA_4 and LPA_5 in mediating LPA-induced platelet shape change (Khandoga *et al.*, 2008; Williams *et al.*, 2009). However, firm evidence that these receptors were the functional platelet LPA receptors was lacking. The LPA response of platelets did not match the pharmacological properties of the LPA_4 and LPA_5 receptors in heterologous expression systems (Khandoga *et al.*, 2008), and the pharmacological receptor agonists and antagonists that were used were not selective for LPA_5 (Williams *et al.*, 2009).

More recently, it was demonstrated that LPA-induced shape change in two human megakaryocytic cell lines was inhibited by siRNA against LPA_5 , but not by knock-down of each of the other receptors $LPA_{1-4,6,7}$ (Khandoga *et al.*, 2011). The rank order of activation by LPA species in these cells, with alkyl-LPA 18:1 and alkyl-LPA 16:0 being the most potent, was similar to that of human platelets, supporting the hypothesis that LPA_5 is the functional LPA receptor-mediating platelet shape change (Figure 1). Importantly, siRNA against LPA_5 also




Figure 2


Hypothetical role of plaque LPA and the platelet LPA_5 receptor in acute atherothrombosis after plaque rupture. LPA in the lipid core of atherosclerotic plaques may act as a cofactor with platelet-adhesive matrix proteins such as collagen type I and III in platelet activation (van Zanten *et al.*, 1994; Penz *et al.*, 2005; Schulz *et al.*, 2008). These matrix proteins are over-expressed in plaques as compared with healthy arterial intima. LPA induces, through binding to the LPA_5 receptor, shape change of circulating platelets and has a synergistic effect with ADP at inducing platelet aggregation and thrombus formation. ADP is secreted from dense granules of platelets adhering to the collagenous matrix of the ruptured cap. GPVI, glycoprotein VI.

inhibited shape change of the megakaryocytic cells that was induced by the lipid-rich core of human atherosclerotic plaques (Khandoga *et al.*, 2011). Thus, LPA_5 may be a novel target for anti-thrombotic therapy for patients with ischaemic cardio- and cerebrovascular disease.

Upon plaque rupture, LPA exposed by the lipid core might, by binding to LPA_5 , induce platelet shape change and stimulate, in synergy with ADP, platelet aggregation and thrombus formation (Haseruck *et al.*, 2004; Rother *et al.*, 2003) (Figure 2). ADP can derive from damaged red cells (Born and Wehmeier, 1979) and from activated platelets adhering to the collagenous matrix of the ruptured cap and secreting their granule contents (Penz *et al.*, 2005; Reininger *et al.*, 2010) (Figure 2). Thus, plaque LPA may act as a cofactor in acute atherothrombosis. However, the importance of LPA in the lipid-rich core for platelet activation in whole blood remains to be demonstrated. Specific LPA_5 receptor antagonists will be helpful to answer this question. Indirect evidence supports a role for plaque LPA in plaque-induced platelet activation and thrombus formation. Low (sub- μ M) concentrations of alkyl-LPA species, that may be reached locally after plaque rupture, activate platelets in blood (Haseruck *et al.*, 2004), and aspirin is equally ineffective at inhibiting LPA-triggered platelet aggregation in blood (Haseruck *et al.*, 2004) and plaque-induced platelet thrombus formation under arterial flow conditions (Penz *et al.*, 2007).

LPA in atherosclerosis

Hypercholesterolaemia promotes the continuous recruitment of circulating monocytes to the arterial wall, which drives the progression of atherosclerotic plaques (Swirski, 2006). LDL enters the vessel wall and is retained in the subendothelial space, where LDL is oxidatively modified. However, the type and degree of LDL modifications are diverse, resulting in distinct biological activities. Oxidized LDL is generally divided into two main categories: minimally modified or mildly oxidized LDL (MM-LDL or mox-LDL) and extensively oxidized LDL (ox-LDL) (Stocker and Keaney, 2004; Levitan *et al.*, 2010). Although both are chemically modified, MM-LDL differs from oxLDL, because it still binds to the LDL receptor and is not recognized by most scavenger receptors (Stocker and Keaney, 2004; Levitan *et al.*, 2010). In contrast to native LDL, MM-LDL can induce the adhesion of monocytes to endothelial cells by up-regulating the CXC chemokine, CXCL1, on the endothelial surface (Berliner *et al.*, 1990; Schwartz *et al.*, 1994). Oxidized LDL or components of LDL that are released during the oxidation, like oxidized phospholipids, trigger the inflammatory response by activating endothelial cells (Hansson and Hermansson, 2011; Weber and Noels, 2011). Activated endothelial cells express adhesion molecules and chemokines that direct the adhesion and transmigration of circulating monocytes. These monocytes



LPA promotes the accumulation of macrophages in atherosclerotic lesions. The moxLDL leads to increased formation of LPC, which is converted by endothelial-derived ATX into LPA. LPA triggers the release of the chemokine CXCL1 from endothelial cells through the activation of LPA₁ and LPA₃. CXCL1 is immobilized on the endothelial surface and induces the adhesion of monocytes to the vessel wall via its receptor CXCR2 on monocytes. These monocytes migrate into the subendothelial space and transform into macrophages, which are the primary cells in early atherosclerotic plaques.

In murine carotid arteries, atherogenic monocyte adhesion under flow conditions is primarily mediated by the CXC chemokine CXCL1 (human GRO- α /murine KC), which is in contrast to CCL2 immobilized on the endothelial surface (Weber *et al.*, 1999; Huo *et al.*, 2001). Whereas most chemokines are transcriptionally up-regulated, CXCL1 is stored in intracellular vesicles of endothelial cells (Oyenebraten, 2004; Zhou *et al.*, 2011). The modified LDL-induced secretion of endothelial CXCL1 is mediated by an unsaturated LPA species and requires the activity of ATX (Zhou *et al.*, 2011). This secretagogue effect of LPA is mediated by the LPA_{1/3}-induced activation of Rho-associated coiled-coil containing protein kinase. However, activation of NF- κ B by unsaturated LPA

Apart from the recruitment of monocytes, several cellular functions of monocytes and macrophages, including the activation status, uptake of LDL and survival, play an important role in atherogenesis (Moore and Tabas, 2011). Human monocytes and macrophages primarily express LPA₁ and LPA₂, and LPA has been shown to mediate the effects of MM-LDL on monocyte activation via LPA₁ (Fueller *et al.*, 2003; D'Aquilio *et al.*, 2007). In agreement with this finding, LPA stimulates the expression of the pro-atherogenic cytokine IL-1 β in a GPCR-dependent manner in a murine macrophage cell line (Chang *et al.*, 2008b). Moreover, LPA increases the uptake of ox-LDL in monocytes/macrophages (Llodra *et al.*, 2004; Chang *et al.*, 2008b). In J774A murine macrophages, which express LPA₂ and LPA₃, LPA-induced lipid accumulation is mediated by the up-regulation of the scavenger receptor A and can be inhibited by the LPA₁/LPA₃ receptor antagonist Ki16425 (Chang *et al.*, 2008a). In addition, LPA binds to the nuclear receptor PPAR γ and induces PPAR γ -dependent CD36 expression in murine macrophages (McIntyre *et al.*, 2003). PPAR γ activation in macrophages has anti-inflammatory effects, and conditional deletion of PPAR γ in macrophages enhances atherosclerosis (Babaev *et al.*, 2005; Bouhrel *et al.*, 2007). Therefore, an atheroprotective role of LPA by activation of PPAR γ could be envisioned, if this mechanism is relevant *in vivo*. Although LPA is known to increase monocyte migration (Zhou *et al.*, 1995; Gustin *et al.*, 2008b), inhibition of reverse transmigration of monocytes by

LPA has been found, which may play a role in the regression of atherosclerosis (Llodra *et al.*, 2004). Moreover, LPA protects macrophages from apoptosis by activating phosphoinositide 3-kinase (PI3K), indicating that LPA impairs the removal of plaque macrophages (Koh *et al.*, 1998).

SMCs accumulate in advanced atherosclerotic plaques, and apoptosis of SMCs is associated with accelerated atherosclerosis. However, exacerbated proliferation of SMCs and the conversion to a pro-inflammatory SMC phenotype may promote the progression of atherosclerosis by enhanced monocyte recruitment (Zeiffer *et al.*, 2004). LPA potently stimulates the proliferation of cultured SMCs and most probably contributes to ox-LDL-induced SMC proliferation (Tokumura *et al.*, 1994; Natarajan *et al.*, 1995; Kim *et al.*, 2006; Damirin *et al.*, 2007; Komachi *et al.*, 2009). LPA mediates the MM-LDL-induced expression of CCL20 in SMCs via LPA receptors, which is secreted from atherosclerotic plaques and elevated in the circulation of patients with hyperlipidaemia (Calvayrac *et al.*, 2011). However, the functional role of LPA-induced CCL20 in atherogenesis needs to be determined. Tissue factor (TF) is a critical determinant of atherosclerotic plaque thrombogenicity. LPA up-regulates TF expression in SMCs via G_i-protein-mediated activation of ERK1/2 (Cui *et al.*, 2003). Furthermore, LPA induces the pro-atherogenic factors CCL2 (MCP-1) and IL-6 in human vascular SMCs *in vitro* (Kaneyuki *et al.*, 2007; Hao *et al.*, 2010).

Taken together, lipoprotein-derived LPA plays a crucial role in atherogenesis by promoting the recruitment of monocytes. Therapeutic targeting of LPA₁ or LPA₃ on endothelial cells may be a promising approach against atherosclerosis.

LPA in vascular remodelling

The arterial vessel wall can adapt to variety of environmental cues, such as changes in pressure, flow, or oxygen supply, by a process called vascular remodelling that mainly includes a response of the SMCs and results in structural alterations of the tunica media (Gibbons and Dzau, 1994). The media may enlarge, for example in arterial or pulmonary hypertension, or diminish, as in the formation of aneurysms. Furthermore, neointimal accumulation of SMCs is characteristic of vascular repair after injury, for example following stent implantation into atherosclerotic arteries (called restenosis). In addition, endothelial cells and inflammatory cells like monocytes/macrophages affect SMC function and the production of extracellular matrix proteins, thereby playing an important role in vascular remodelling (Schober, 2008). The phenotypic switch of SMCs from a contractile, quiescent state towards a synthetic, proliferative state is a common feature during vascular remodelling and may be causally related to neointima formation (Owens *et al.*, 2004).

Both migration and proliferation of medial SMCs following vascular injury have been implicated in neointima formation (Schwartz *et al.*, 1995). *In vitro*, LPA has been shown to induce the proliferation of SMCs via activation of LPA₁ and G_{i/q} proteins and involves PKC, ERK1/2, the PI3K/PKB (Akt) pathway and MAPK cascades (Tokumura *et al.*, 1994; Seewald *et al.*, 1997; 1999; Gennero *et al.*, 1999; Schmitz *et al.*, 2002; Xu *et al.*, 2003; Gouni-Berthold *et al.*, 2004; Baldini *et al.*,

2005; Kim *et al.*, 2006; Komachi *et al.*, 2009). In contrast, combined genetic deletion of LPA₁ and LPA₂ is required to inhibit serum-induced growth of SMCs (Panchatcharam *et al.*, 2008). Furthermore, LPA enhances migratory activity of vascular SMCs by G_{i/q} protein-coupled LPA₁ receptor-mediated activation of the p38MAPK pathway (Kim *et al.*, 2006; Damirin *et al.*, 2007; Komachi *et al.*, 2009; Zhou *et al.*, 2009). However, SMCs from LPA₁^{-/-} mice showed increased migration due to compensatory up-regulation of LPA₃ (Panchatcharam *et al.*, 2008). Whereas the migration of LPA₂-deficient SMCs is not impaired, combined deletion of LPA₁ and LPA₂ inhibits SMC migration (Panchatcharam *et al.*, 2008). These results indicate that LPA₃ also promotes SMC migration in the presence of LPA₂. Neointimal SMCs display a pro-inflammatory phenotype driven by activated NF-κB signalling, which may promote leucocyte recruitment to the injured artery (Zeiffer *et al.*, 2004). LPA increases the expression of IL-6 and CCL2 in vascular SMCs via LPA₁ and PKC-mediated p38 MAPK and NADPH oxidase-dependent generation of reactive oxygen species, respectively (Kaneyuki *et al.*, 2007; Hao *et al.*, 2010). The early growth response gene 1 (Egr1) is crucial for SMC proliferation and neointima formation (Khachigian, 2006). Egr1 is induced by LPA-mediated activation of the transcription factors cAMP response element-binding (CREB) and serum response factor (SRF) in SMCs (Cui *et al.*, 2006). However, the role of LPA-induced Egr1 in vascular remodelling remains to be defined. The reduced expression of contractile proteins is characteristic for synthetic SMCs, and this dedifferentiation process has been implicated in neointima formation (Owens *et al.*, 2004). LPA species with an unsaturated fatty acyl chain selectively down-regulate contractile protein expression in SMCs via sustained activation of ERK1/2 and p38 MAPK (Hayashi *et al.*, 2001). This effect of LPA on SMC dedifferentiation has been attributed to the activation of LPA₃ using LPA receptor-specific antagonists and agonists (Zhou *et al.*, 2010). In agreement with this finding, no differences in LPA-induced SMC dedifferentiation have been observed between wild-type, LPA₁^{-/-}, LPA₂^{-/-}, LPA₁^{-/-}/LPA₂^{-/-} and PPARγ^{-/-} SMCs (Guo *et al.*, 2008; Panchatcharam *et al.*, 2008). However, the PPARγ agonist rosiglitazone reduced the expression of SMC markers, like unsaturated LPA, indicating different pathways for PPARγ- and LPA-induced SMC differentiation (Zhang *et al.*, 2004). Taken together, LPA acts in multiple ways on SMCs *in vitro*, which suggests that LPA promotes vascular remodelling and neointima formation.

To study the role of LPA in vascular remodelling *in vivo*, short-term incubation of the carotid arteries with LPA has been performed. Intriguingly, transient treatment only with unsaturated LPAs, like LPA18:1, LPA20:4 or the LPA analogue 1-AGP18:1 (1-O-octadecenyl glycerophosphate), triggers the formation within weeks of a neointima that primarily consists of SMCs (Yoshida *et al.*, 2003; Zhang *et al.*, 2004; Cheng *et al.*, 2009; Subramanian *et al.*, 2010). Blocking LPA18:1-induced ERK and p38MAPK activation prevented neointima formation (Yoshida *et al.*, 2003). Furthermore, LPA20:4 and 1-AGP18:1-induced neointima formation was reduced by treatment with a PPARγ antagonist and in mice with PPARγ-deficiency in vascular wall cells (Zhang *et al.*, 2004; Cheng *et al.*, 2009). Although LPA20:4-induced neointima formation was partially inhibited by pertussis toxin and the LPA₁/LPA₃




Figure 4

LPA induces neointima formation after vascular injury. Following vascular injury, activated platelets adhere to the denuded surface of the vessel wall and medial SMCs undergo apoptosis. These early events after vascular injury may induce the production of LPA, which increases CXCL12 in the vessel wall through its receptors LPA₁ and LPA₃. CXCL12 is released into the circulation and recruits SPC via its receptor CXCR4 to the injury site. These SPCs differentiate into neointimal SMCs, which form the neointimal lesion.

antagonist diocetyl glycerol pyrophosphate in rats, which do not express LPA₃ in the carotid wall, neointimal growth following 1-AGP18:1 incubation was not impaired in LPA₁^{-/-}, LPA₂^{-/-} and LPA₁^{-/-}/LPA₂^{-/-} mice (Zhang *et al.*, 2004; Cheng *et al.*, 2009). Neointima formation by carotid ligation is increased in LPA₁^{-/-} mice, whereas a combined deficiency of LPA₁ and LPA₂ diminishes neointimal growth (Panchatcharam *et al.*, 2008). The compensatory overexpression of LPA₃ in SMCs from LPA₁^{-/-} mice, which is absent in LPA₁^{-/-}/LPA₂^{-/-} SMCs, is associated with enhanced neointima formation (Panchatcharam *et al.*, 2008). PPAR γ activation in SMCs has been shown to inhibit proliferation and migration and attenuates neointimal hyperplasia after vascular injury (Lim *et al.*, 2006; Lee *et al.*, 2009; Zhang *et al.*, 2011). Accordingly, 1-AGP18:1, but not rosiglitazone, increases injury-induced neointima formation, indicating that LPA-mediated PPAR γ activation does not play a role in vascular remodelling following vascular injury (Cheng *et al.*, 2009). Taken together, these results show that molecular mechanisms of LPA-dependent neointima formation may differ between local treatment with LPA and in established animal models of vascular remodelling.

In atherosclerosis-prone Apoe^{-/-} mice, treatment with the LPA₁/LPA₃ antagonist Ki16425 diminished neointimal hyperplasia after carotid wire injury by reducing the SMC and macrophage content in the lesions, indicating that LPA promotes injury-induced neointima formation via LPA receptors (Subramanian *et al.*, 2010). Interestingly, unsaturated but not saturated LPA induces sustained up-regulation of the chemokine CXCL12 (stromal cell-derived factor 1) in the vessel wall, which mobilizes and recruits smooth muscle progenitor cells (SPCs) from the bone marrow into the injured artery (Schober, 2008; Subramanian *et al.*, 2010) (Figure 4). Silencing of either LPA₁ or LPA₃ in the vessel wall impaired the LPA20:4-induced neointima formation and SPC mobilization, suggesting that, similar to atherogenic monocyte recruitment, these two LPA receptors independently trigger the vascular response (Subramanian *et al.*, 2010). It remains to be determined whether this functional independency might be due to the formation of heterodimers between LPA₁

and LPA₃, and whether LPA₁/LPA₃ heterodimers have an increased binding affinity to unsaturated LPAs (Zaslavsky *et al.*, 2006). The different mechanisms that were observed in LPA-induced neointimal hyperplasia might be dose-dependent. Whereas 40 μ M of LPA20:4 has been found to trigger CXCL12 expression, unsaturated LPAs in a dosage range between 1 and 10 μ M induce ERK1/2/p38MAPK- or PPAR γ -dependent neointima formation (Yoshida *et al.*, 2003; Zhang *et al.*, 2004; Cheng *et al.*, 2009; Subramanian *et al.*, 2010). However, the fact that pharmacological inhibition of LPA₁ and LPA₃ after carotid injury inhibits CXCL12 expression and impairs CXCL12-dependent SPC mobilization indicates that the 40 μ M LPA20:4 dose used for local carotid treatment closely mirrors the role of LPA after vascular injury (Subramanian *et al.*, 2010). Microvesicles released from apoptotic SMCs following injury have been found to up-regulate CXCL12 expression in non-injured SMCs, and blocking the apoptosis of SMCs following vascular injury greatly inhibits CXCL12 expression (Zernecke *et al.*, 2005). Taking into account the role of microvesicles in LPA generation (Fourcade *et al.*, 1995), increased LPA production due to the release of microvesicles from apoptotic SMCs may contribute to the up-regulation of CXCL12 in the injured arteries. In addition, activated platelets adhering to the denuded vascular surface may not only present CXCL12 during the recruitment of SPCs, but also increase CXCL12 expression by enhancing LPA formation (Zernecke *et al.*, 2005).

Remodelling of the pulmonary vasculature occurs during the hypoxia-driven development of pulmonary hypertension and is characterized by the thickening of the media and adventitia in pulmonary arteries and muscularization of non-muscular alveolar wall vessels (Stenmark *et al.*, 2006). Interestingly, hypoxia-induced pulmonary hypertension and remodelling is accelerated in both ATX^{+/-} mice and LPA₁^{-/-}/LPA₂^{-/-} mice (Cheng *et al.*, 2012). In LPA₁^{-/-}/LPA₂^{-/-} mice, thickening of the arteriolar vessel wall is already evident during normoxia, and aging results in severe pulmonary hypertension (Cheng *et al.*, 2012). This increased remodelling of pulmonary vessels was associated with increased expression of genes involved in endothelin-1 signalling (Cheng

et al., 2012). In contrast, the deficiency of either LPA₁ or LPA₂ is not associated with pulmonary hypertension. This indicates that LPA signalling via LPA₁ and LPA₂ in the pulmonary vasculature has an important homeostatic role and appears to have protective effects in hypoxia-induced pulmonary hypertension.

Conclusions and future directions

LPA has atherogenic actions, and specific LPA receptors have been identified that may mediate some of these harmful LPA activities. The experimental evidence is based not only on numerous *in vitro* studies, but also on several *in vivo* studies. LPA does not only enhance atherosclerosis, but also *vice versa*, cardiovascular risk factors affect the ATX/LPA axis thereby further aggravating the progression of atherosclerotic diseases. Obesity and hyperlipidaemia enhance ATX expression and activity, and the generation of LPA in the circulation. Increased ATX-dependent LPA formation from LDL-derived LPC promotes endothelial chemokine (CXCL1) secretion leading to the recruitment of monocytes to the atherosclerotic lesion, and thereby enhancing atherosclerosis (Figure 3). Endothelial LPA_{1/3} receptors and Rho-kinase are critical here. LPA also increases the uptake of ox-LDL in monocytes/macrophages by up-regulation of the scavenger receptor A, and inhibits monocyte egression from plaques. LPA may also induce these effects by binding to LPA_{1/3} and to the nuclear receptor PPAR γ . LPA also stimulates VSMC dedifferentiation, proliferation and neointima formation. Also here, LPA₁ and LPA₃ receptors as well as PPAR γ seem to be involved; the situation is, however, complex, as the mechanism of LPA-dependent neointima formation seems to differ between local treatment with LPA and established animal models of vascular remodelling. Lastly but not least, LPA accumulating in the lipid-rich core may contribute to acute thrombus formation after plaque rupture by stimulation of the platelet LPA₃ receptor and Rho-kinase (Figures 1, 2). Future studies might be directed to explore the role of the non-Edg receptor family of LPA receptors in various animal models of atherosclerosis, to determine the expression of individual LPA receptors in human atherosclerotic lesions *in situ* and to evaluate the effects of novel LPA receptor antagonists on the development, progression and regression of atherosclerosis in animal models, as well as on thrombosis induced by atherosclerotic plaques in human models. Clinical studies could determine whether plasma levels of defined LPA molecular species could be a novel biomarker of cardiovascular risk. In such studies, the addition of pharmacological ATX inhibitors to the blood collection tube might be important to minimize artificial *in vitro* LPA formation during blood handling.

Acknowledgements

This publication was made possible by grants to AS and WS from the Deutsche Forschungsgemeinschaft (Scho 1056/2, Scho 1056/3, Si 274/9; Si 274/11), the August-Lenz-Stiftung and the Bayern University.

Statement of conflict of interest

The authors have no conflict of interest.

References

- Adinolfi LE, Restivo L, Zampino R, Guerrero B, Lonardo A, Ruggiero L *et al.* (2012). Chronic HCV infection is a risk of atherosclerosis. Role of HCV and HCV-related steatosis. *Atherosclerosis* 221: 496–502.
- Aird WC (2005). Spatial and temporal dynamics of the endothelium. *J Thromb Haemost* 3: 1392–1406.
- Albers HM, Dong A, van Meeteren LA, Egan DA, Sunkara M, van Tilburg EW *et al.* (2010). Boronic acid-based inhibitor of autotaxin reveals rapid turnover of LPA in the circulation. *Proc Natl Acad Sci U S A* 107: 7257–7262.
- Albert MA, Glynn RJ, Wolfert RL, Ridker PM (2005). The effect of statin therapy on lipoprotein associated phospholipase A2 levels. *Atherosclerosis* 182: 193–198.
- Alexander JS, Patton WF, Christman BW, Cuiper LL, Haselton FR (1998). Platelet-derived lysophosphatidic acid decreases endothelial permeability *in vitro*. *Am J Physiol* 274(1 Pt 2): H115–H122.
- Amisten S, Braun OO, Bengtsson A, Erlinge D (2008). Gene expression profiling for the identification of G-protein coupled receptors in human platelets. *Thromb Res* 122: 47–57.
- Aoki J, Taira A, Takanezawa Y, Kishi Y, Hama K, Kishimoto T *et al.* (2002). Serum lysophosphatidic acid is produced through diverse phospholipase pathways. *J Biol Chem* 277: 48737–48744.
- Babaev VR, Yancey PG, Ryzhov SV, Kon V, Breyer MD, Magnuson MA *et al.* (2005). Conditional knockout of macrophage PPAR γ increases atherosclerosis in C57BL/6 and low-density lipoprotein receptor-deficient mice. *Arterioscler Thromb Vasc Biol* 25: 1647–1653.
- Baker DL, Desiderio DM, Miller DD, Tolley B, Tigyi GJ (2001). Direct quantitative analysis of lysophosphatidic acid molecular species by stable isotope dilution electrospray ionization liquid chromatography-mass spectrometry. *Anal Biochem* 292: 287–295.
- Baldini PM, De Vito P, D'Aquilio F, Vismara D, Zalfa F, Bagni C *et al.* (2005). Role of atrial natriuretic peptide in the suppression of lysophosphatidic acid-induced rat aortic smooth muscle (RASM) cell growth. *Mol Cell Biochem* 272: 19–28.
- le Balle F, Simon MF, Meijer S, Fourcade O, Chap H (1999). Membrane sidedness of biosynthetic pathways involved in the production of lysophosphatidic acid. *Adv Enzyme Regul* 39: 275–284.
- Bauer M, Retzer M, Wilde JI, Maschberger P, Essler M, Aepfelbacher M *et al.* (1999). Dichotomous regulation of myosin phosphorylation and shape change by Rho-kinase and calcium in intact human platelets. *Blood* 94: 1665–1672.
- Benitez S, Sanchez-Quesada JL, Ribas V, Jorba O, Blanco-Vaca F, Gonzalez-Sastre F *et al.* (2003). Platelet-activating factor acetylhydrolase is mainly associated with electronegative low-density lipoprotein subfraction. *Circulation* 108: 92–96.
- Berliner JA, Territo MC, Sevanian A, Ramin S, Kim JA, Bamshad B *et al.* (1990). Minimally modified low density lipoprotein stimulates monocyte endothelial interactions. *J Clin Invest* 85: 1260–1266.

- Billah MM, Lapetina EG, Cuatrecasas P (1981). Phospholipase A2 activity specific for phosphatidic acid. A possible mechanism for the production of arachidonic acid in platelets. *J Biol Chem* 256: 5399–5403.
- Bolen AL, Naren AP, Yarlagadda S, Beranova-Giorgianni S, Chen L, Norman D *et al.* (2011). The phospholipase A1 activity of lysophospholipase A-I links platelet activation to LPA production during blood coagulation. *J Lipid Res* 52: 958–970.
- Born GV, Wehmeier A (1979). Inhibition of platelet thrombus formation by chlorpromazine acting to diminish haemolysis. *Nature* 282: 212–213.
- Bot M, Bot I, Lopez-Vales R, van de Lest CHA, Saulnier-Blache JS, Helms JB *et al.* (2010). Atherosclerotic lesion progression changes lysophosphatidic acid homeostasis to favor its accumulation. *Am J Pathol* 176: 3073–3084.
- Boucher J, Quilliot D, Praderes JP, Simon MF, Gres S, Guigne C *et al.* (2005). Potential involvement of adipocyte insulin resistance in obesity-associated up-regulation of adipocyte lysophospholipase D/autotaxin expression. *Diabetologia* 48: 569–577.
- Bouhrel MA, Derudas B, Rigamonti E, Dièvert R, Brozek J, Haulon S *et al.* (2007). PPAR γ activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metab* 6: 137–143.
- Calvayrac O, Rodriguez-Calvo R, Alonso J, Orbe J, Martin-Ventura JL, Guadall A *et al.* (2011). CCL20 is increased in hypercholesterolemic subjects and is upregulated by LDL in vascular smooth muscle cells: role of NF- κ B. *Arterioscler Thromb Vasc Biol* 31: 2733–2741.
- Chang CL, Hsu HY, Lin HY, Chiang W, Lee H (2008a). Lysophosphatidic acid-induced oxidized low-density lipoprotein uptake is class A scavenger receptor-dependent in macrophages. *Prostaglandins Other Lipid Mediat* 87: 20–25.
- Chang CL, Lin ME, Hsu HY, Yao CL, Hwang SM, Pan CY *et al.* (2008b). Lysophosphatidic acid-induced interleukin-1 β expression is mediated through Gi/Rho and the generation of reactive oxygen species in macrophages. *J Biomed Sci* 15: 357–363.
- Chappell D, Jacob M, Paul O, Rehm M, Welsch U, Stoeckelhuber M *et al.* (2009). The glycocalyx of the human umbilical vein endothelial cell: an impressive structure ex vivo but not in culture. *Circ Res* 104: 1313–1317.
- Charo IF, Taub R (2011). Anti-inflammatory therapeutics for the treatment of atherosclerosis. *Nat Rev Drug Discov* 10: 365–376.
- Cheng HY, Dong A, Panchatcharam M, Mueller P, Yang F, Li Z *et al.* (2012). Lysophosphatidic Acid signaling protects pulmonary vasculature from hypoxia-induced remodeling. *Arterioscler Thromb Vasc Biol* 32: 24–32.
- Cheng Y, Makarova N, Tsukahara R, Guo H, Shuyu E, Farrar P *et al.* (2009). Lysophosphatidic acid-induced arterial wall remodeling: requirement of PPAR γ but not LPA1 or LPA2 GPCR. *Cell Signal* 21: 1874–1884.
- Choi JW, Herr DR, Noguchi K, Yung YC, Lee CW, Mutoh T *et al.* (2010). LPA receptors: subtypes and biological actions. *Annu Rev Pharmacol Toxicol* 50: 157–186.
- Chun J, Hla T, Lynch KR, Spiegel S, Moolenaar WH (2010). International Union of Basic and Clinical Pharmacology. LXXVIII. Lysophospholipid receptor nomenclature. *Pharmacol Rev* 62: 579–587.
- Corti R, Badimon JJ (2002). Biologic aspects of vulnerable plaque. *Curr Opin Cardiol* 17: 616–625.
- Cui MZ, Zhao G, Winokur AL, Laag E, Bydash JR, Penn MS *et al.* (2003). Lysophosphatidic acid induction of tissue factor expression in aortic smooth muscle cells. *Arterioscler Thromb Vasc Biol* 23: 224–230.
- Cui MZ, Laag E, Sun L, Tan M, Zhao G, Xu X (2006). Lysophosphatidic acid induces early growth response gene 1 expression in vascular smooth muscle cells: CRE and SRE mediate the transcription. *Arterioscler Thromb Vasc Biol* 26: 1029–1035.
- D'Aquilio F, Procaccini M, Izzi V, Chiurchiu V, Giambra V, Carotenuto F *et al.* (2007). Activatory properties of lysophosphatidic acid on human THP-1 cells. *Inflammation* 30: 167–177.
- Damirin A, Tomura H, Komachi M, Liu JP, Mogi C, Tobo M *et al.* (2007). Role of lipoprotein-associated lysophospholipids in migratory activity of coronary artery smooth muscle cells. *Am J Physiol Heart Circ Physiol* 292: H2513–H2522.
- Dohi T, Miyauchi K, Ohkawa R, Nakamura K, Kishimoto T, Miyazaki T *et al.* (2012). Increased circulating plasma lysophosphatidic acid in patients with acute coronary syndrome. *Clin Chim Acta* 413: 207–212.
- Dusaucy R, Rancoule C, Gres S, Wanecq E, Colom A, Guigne C *et al.* (2011). Adipose-specific disruption of autotaxin enhances nutritional fattening and reduces plasma lysophosphatidic acid. *J Lipid Res* 52: 1247–1255.
- Eichholtz T, Jalink K, Fahrenfort I, Moolenaar WH (1993). The bioactive phospholipid lysophosphatidic acid is released from activated platelets. *Biochem J* 291 (Pt 3): 677–680.
- Fernandez-Ortiz A, Badimon JJ, Falk E, Fuster V, Meyer B, Mailhac A *et al.* (1994). Characterization of the relative thrombogenicity of atherosclerotic plaque components: implications for consequences of plaque rupture. *J Am Coll Cardiol* 23: 1562–1569.
- FitzGerald GA, Healy C, Daugherty J (1987). Thromboxane A2 biosynthesis in human disease. *Fed Proc* 46: 154–158.
- Fourcade O, Simon MF, Viode C, Rugani N, Leballe F, Ragab A *et al.* (1995). Secretory phospholipase A2 generates the novel lipid mediator lysophosphatidic acid in membrane microvesicles shed from activated cells. *Cell* 80: 919–927.
- Fueller M, Wang DA, Tigyi G, Siess W (2003). Activation of human monocytic cells by lysophosphatidic acid and sphingosine-1-phosphate. *Cell Signal* 15: 367–375.
- Fujiwara Y (2008). Cyclic phosphatidic acid – a unique bioactive phospholipid. *Biochim Biophys Acta* 1781: 519–524.
- Fulkerson Z, Wu T, Sunkara M, Kooi CV, Morris AJ, Smyth SS (2011). Binding of autotaxin to integrins localizes lysophosphatidic acid production to platelets and mammalian cells. *J Biol Chem* 286: 34654–34663.
- Fuster V, Moreno PR, Fayad ZA, Corti R, Badimon JJ (2005). Atherothrombosis and high-risk plaque: part I: evolving concepts. *J Am Coll Cardiol* 46: 937–954.
- Gaits F, Fourcade O, Le Balle F, Gueguen G, Gaige B, Gassama-Diagne A *et al.* (1997). Lysophosphatidic acid as a phospholipid mediator: pathways of synthesis. *FEBS Lett* 410: 54–58.
- Gennero I, Xuereb JM, Simon MF, Girolami JP, Bascands JL, Chap H *et al.* (1999). Effects of lysophosphatidic acid on proliferation and cytosolic Ca⁺⁺ of human adult vascular smooth muscle cells in culture. *Thromb Res* 94: 317–326.

- Gibbons GH, Dzau VJ (1994). The emerging concept of vascular remodeling. *N Engl J Med* 330: 1431–1438.
- Gierse J, Thorarensen A, Beltey K, Bradshaw-Pierce E, Cortes-Burgos L, Hall T *et al.* (2010). A novel autotaxin inhibitor reduces lysophosphatidic acid levels in plasma and the site of inflammation. *J Pharmacol Exp Ther* 334: 310–317.
- Goncalves I, Edsfeldt A, Ko NY, Grufman H, Berg K, Bjorkbacka H *et al.* (2012). Evidence supporting a key role of Lp-PLA2-generated lysophosphatidylcholine in human atherosclerotic plaque inflammation. *Arterioscler Thromb Vasc Biol* 32: 1505–1512.
- Gouni-Berthold I, Seewald S, Hescheler J, Sachinidis A (2004). Regulation of mitogen-activated protein kinase cascades by low density lipoprotein and lysophosphatidic acid. *Cell Physiol Biochem* 14: 167–176.
- Gratacap MP, Payraastre B, Nieswandt B, Offermanns S (2001). Differential regulation of Rho and Rac through heterotrimeric G-proteins and cyclic nucleotides. *J Biol Chem* 276: 47906–47913.
- Guo H, Makarova N, Cheng Y, E S, Ji RR, Zhang C *et al.* (2008). The early- and late stages in phenotypic modulation of vascular smooth muscle cells: differential roles for lysophosphatidic acid. *Biochim Biophys Acta* 1781: 571–581.
- Gupte R, Patil R, Liu J, Wang Y, Lee SC, Fujiwara Y *et al.* (2011). Benzyl and naphthalene methylphosphonic acid inhibitors of autotaxin with anti-invasive and anti-metastatic activity. *ChemMedChem* 6: 922–935.
- Gustin C, Delaive E, Dieu M, Calay D, Raes M (2008a). Upregulation of pentraxin-3 in human endothelial cells after lysophosphatidic acid exposure. *Arterioscler Thromb Vasc Biol* 28: 491–497.
- Gustin C, Van Steenbrugge M, Raes M (2008b). LPA modulates monocyte migration directly and via LPA-stimulated endothelial cells. *Am J Physiol Cell Physiol* 295: C905–C914.
- Hansson GK, Hermansson A (2011). The immune system in atherosclerosis. *Nat Immunol* 12: 204–212.
- Hao F, Tan M, Wu DD, Xu X, Cui MZ (2010). LPA induces IL-6 secretion from aortic smooth muscle cells via an LPA1-regulated, PKC-dependent, and p38[alpha]-mediated pathway. *Am J Physiol Heart Circ Physiol* 298: H974–H983.
- Haseruck N, Erl W, Pandey D, Tigyi G, Ohlmann P, Ravanat C *et al.* (2004). The plaque lipid lysophosphatidic acid stimulates platelet activation and platelet-monocyte aggregate formation in whole blood: involvement of P2Y1 and P2Y12 receptors. *Blood* 103: 2585–2592.
- Hausmann J, Kamtekar S, Christodoulou E, Day JE, Wu T, Fulkerson Z *et al.* (2011). Structural basis of substrate discrimination and integrin binding by autotaxin. *Nat Struct Mol Biol* 18: 198–204.
- Hayashi K, Takahashi M, Nishida W, Yoshida K, Ohkawa Y, Kitabatake A *et al.* (2001). Phenotypic modulation of vascular smooth muscle cells induced by unsaturated lysophosphatidic acids. *Circ Res* 89: 251–258.
- Hirakawa M, Oike M, Karashima Y, Ito Y (2004). Sequential activation of RhoA and FAK/paxillin leads to ATP release and actin reorganization in human endothelium. *J Physiol* 558 (Pt 2): 479–488.
- Hirase T, Kawashima S, Wong EY, Ueyama T, Rikitake Y, Tsukita S *et al.* (2001). Regulation of tight junction permeability and occludin phosphorylation by RhoA-p160ROCK-dependent and -independent mechanisms. *J Biol Chem* 276: 10423–10431.
- Holvoet P, Vanhaecke J, Janssens S, Van de Werf F, Collen D (1998). Oxidized LDL and malondialdehyde-modified LDL in patients with acute coronary syndromes and stable coronary artery disease. *Circulation* 98: 1487–1494.
- Hoofnagle AN, Heinecke JW (2009). Lipoproteomics: using mass spectrometry-based proteomics to explore the assembly, structure, and function of lipoproteins. *J Lipid Res* 50: 1967–1975.
- Hoshiga M, Alpers CE, Smith LL, Giachelli CM, Schwartz SM (1995). Alpha-v beta-3 integrin expression in normal and atherosclerotic artery. *Circ Res* 77: 1129–1135.
- Hosogaya S, Yatomi Y, Nakamura K, Ohkawa R, Okubo S, Yokota H *et al.* (2008). Measurement of plasma lysophosphatidic acid concentration in healthy subjects: strong correlation with lysophospholipase D activity. *Ann Clin Biochem* 45: 364–368.
- Huo Y, Weber C, Forlow SB, Sperandio M, Thatté J, Mack M *et al.* (2001). The chemokine KC, but not monocyte chemoattractant protein-1, triggers monocyte arrest on early atherosclerotic endothelium. *J Clin Invest* 108: 1307–1314.
- Kaneyuki U, Ueda S, Yamagishi S, Kato S, Fujimura T, Shibata R *et al.* (2007). Pitavastatin inhibits lysophosphatidic acid-induced proliferation and monocyte chemoattractant protein-1 expression in aortic smooth muscle cells by suppressing Rac-1-mediated reactive oxygen species generation. *Vascul Pharmacol* 46: 286–292.
- Khachigian LM (2006). Early growth response-1 in cardiovascular pathobiology. *Circ Res* 98: 186–191.
- Khandoga AL, Fujiwara Y, Goyal P, Pandey D, Tsukahara R, Bolen A *et al.* (2008). Lysophosphatidic acid-induced platelet shape change revealed through LPA(1-5) receptor-selective probes and albumin. *Platelets* 19: 415–427.
- Khandoga AL, Pandey D, Welsch U, Brandl R, Siess W (2011). GPR92/LPA lysophosphatidate receptor mediates megakaryocytic cell shape change induced by human atherosclerotic plaques. *Cardiovasc Res* 90: 157–164.
- Kim J, Keys JR, Eckhart AD (2006). Vascular smooth muscle migration and proliferation in response to lysophosphatidic acid (LPA) is mediated by LPA receptors coupling to Gq. *Cell Signal* 18: 1695–1701.
- Kimura K, Ito M, Amano M, Chihara K, Fukata Y, Nakafuku M *et al.* (1996). Regulation of myosin phosphatase by Rho and Rho-associated kinase (Rho-kinase). *Science* 273: 245–248.
- Kishimoto T, Matsuoka T, Imamura S, Mizuno K (2003). A novel colorimetric assay for the determination of lysophosphatidic acid in plasma using an enzymatic cycling method. *Clin Chim Acta* 333: 59–67.
- Klages B, Brandt U, Simon MI, Schultz G, Offermanns S (1999). Activation of G12/G13 results in shape change and Rho/Rho-kinase-mediated myosin light chain phosphorylation in mouse platelets. *J Cell Biol* 144: 745–754.
- Kobayashi T, Tanaka-Ishii R, Taguchi R, Ikezawa H, Murakami-Murofushi K (1999). Existence of a bioactive lipid, cyclic phosphatidic acid, bound to human serum albumin. *Life Sci* 65: 2185–2191.
- Koenig W, Khuseyinova N (2009). Lipoprotein-associated and secretory phospholipase A2 in cardiovascular disease: the epidemiological evidence. *Cardiovasc Drugs Ther* 23: 85–92.
- Koh JS, Lieberthal W, Heydrick S, Levine JS (1998). Lysophosphatidic acid is a major serum noncytokine survival factor for murine macrophages which acts via the phosphatidylinositol 3-kinase signaling pathway. *J Clin Invest* 102: 716–727.

- Kolodgie FD, Burke AP, Wight TN, Virmani R (2004). The accumulation of specific types of proteoglycans in eroded plaques: a role in coronary thrombosis in the absence of rupture. *Curr Opin Lipidol* 15: 575–582.
- Komachi M, Damirin A, Malchinkhuu E, Mogi C, Tobo M, Ohta H *et al.* (2009). Signaling pathways involved in DNA synthesis and migration in response to lysophosphatidic acid and low-density lipoprotein in coronary artery smooth muscle cells. *Vascul Pharmacol* 50: 178–184.
- Lee CS, Kwon YW, Yang HM, Kim SH, Kim TY, Hur J *et al.* (2009). New mechanism of rosiglitazone to reduce neointimal hyperplasia: activation of glycogen synthase kinase-3 β followed by inhibition of MMP-9. *Arterioscler Thromb Vasc Biol* 29: 472–479.
- Lee H, Lin CI, Liao J-J LY-W, Yang HY, Lee C-Y *et al.* (2004). Lysophospholipids increase ICAM-1 expression in HUVEC through a Gi- and NF- κ B-dependent mechanism. *Am J Physiol Cell Physiol* 287: C1657–C1666.
- Levitani I, Volkov S, Subbiah PV (2010). Oxidized LDL: diversity, patterns of recognition, and pathophysiology. *Antioxid Redox Signal* 13: 39–75.
- Li ZG, Yu ZC, Wang DZ, Ju WP, Zhan X, Wu QZ *et al.* (2008). Influence of acetylsalicylate on plasma lysophosphatidic acid level in patients with ischemic cerebral vascular diseases. *Neurol Res* 30: 366–369.
- Li ZG, Yu ZC, Yu YP, Ju WP, Wang DZ, Zhan X *et al.* (2010). Lysophosphatidic acid level and the incidence of silent brain infarction in patients with nonvalvular atrial fibrillation. *Int J Mol Sci* 11: 3988–3998.
- Libby P, Ridker PM, Hansson GK (2011). Progress and challenges in translating the biology of atherosclerosis. *Nature* 473: 317–325.
- Liebisch G, Scherer M (2012). Quantification of bioactive sphingo- and glycerophospholipid species by electrospray ionization tandem mass spectrometry in blood. *J Chromatogr B Analyt Technol Biomed Life Sci* 883–884: 141–146.
- Lim S, Jin CJ, Kim M, Chung SS, Park HS, Lee IK *et al.* (2006). PPAR γ gene transfer sustains apoptosis, inhibits vascular smooth muscle cell proliferation, and reduces neointima formation after balloon injury in rats. *Arterioscler Thromb Vasc Biol* 26: 808–813.
- Lin CI, Chen CN, Chen JH, Lee H (2006). Lysophospholipids increase IL-8 and MCP-1 expressions in human umbilical cord vein endothelial cells through an IL-1-dependent mechanism. *J Cell Biochem* 99: 1216–1232.
- Lin CI, Chen CN, Lin PW, Chang KJ, Hsieh FJ, Lee H (2007). Lysophosphatidic acid regulates inflammation-related genes in human endothelial cells through LPA1 and LPA3. *Biochem Biophys Res Commun* 363: 1001–1008.
- Llodra J, Angeli V, Liu J, Trogan E, Fisher EA, Randolph GJ (2004). Emigration of monocyte-derived cells from atherosclerotic lesions characterizes regressive, but not progressive, plaques. *Proc Natl Acad Sci U S A* 101: 11779–11784.
- McConnell JP, Hoefner DM (2006). Lipoprotein-associated phospholipase A2. *Clin Lab Med* 26: 679–697, vii.
- McIntyre TM, Pontsler AV, Silva AR, St Hilaire A, Xu Y, Hinshaw JC *et al.* (2003). Identification of an intracellular receptor for lysophosphatidic acid (LPA): LPA is a transcellular PPAR γ agonist. *Proc Natl Acad Sci U S A* 100: 131–136.
- Maschberger P, Bauer M, Baumann-Siemons J, Zangl KJ, Negrescu EV, Reininger AJ *et al.* (2000). Mildly oxidized low density lipoprotein rapidly stimulates via activation of the lysophosphatidic acid receptor Src family and Syk tyrosine kinases and Ca²⁺ influx in human platelets. *J Biol Chem* 275: 19159–19166.
- Matsumoto T, Kobayashi T, Kamata K (2007). Role of lysophosphatidylcholine (LPC) in atherosclerosis. *Curr Med Chem* 14: 3209–3220.
- van Meeteren LA, Moolenaar WH (2007). Regulation and biological activities of the autotaxin-LPA axis. *Prog Lipid Res* 46: 145–160.
- van Meeteren LA, Ruurs P, Stortelers C, Bouwman P, van Rooijen MA, Pradere JP *et al.* (2006). Autotaxin, a secreted lysophospholipase D, is essential for blood vessel formation during development. *Mol Cell Biol* 26: 5015–5022.
- Minnear FL, Patil S, Bell D, Gainor JP, Morton CA (2001). Platelet lipid(s) bound to albumin increases endothelial electrical resistance: mimicked by LPA. *Am J Physiol Lung Cell Mol Physiol* 281: L1337–L1344.
- Moers A, Nieswandt B, Massberg S, Wettschureck N, Gruner S, Konrad I *et al.* (2003). G13 is an essential mediator of platelet activation in hemostasis and thrombosis. *Nat Med* 9: 1418–1422.
- Moore KJ, Tabas I (2011). Macrophages in the pathogenesis of atherosclerosis. *Cell* 145: 341–355.
- Morris AJ, Panchatcharam M, Cheng HY, Federico L, Fulkerson Z, Selim S *et al.* (2009). Regulation of blood and vascular cell function by bioactive lysophospholipids. *J Thromb Haemost* 7 (Suppl 1): 38–43.
- Nakamura K, Kishimoto T, Ohkawa R, Okubo S, Tozuka M, Yokota H *et al.* (2007). Suppression of lysophosphatidic acid and lysophosphatidylcholine formation in the plasma in vitro: proposal of a plasma sample preparation method for laboratory testing of these lipids. *Anal Biochem* 367: 20–27.
- Nakamura K, Igarashi K, Ide K, Ohkawa R, Okubo S, Yokota H *et al.* (2008). Validation of an autotaxin enzyme immunoassay in human serum samples and its application to hypoalbuminemia differentiation. *Clin Chim Acta* 388: 51–58.
- Nakanaga K, Hama K, Aoki J (2010). Autotaxin – an LPA producing enzyme with diverse functions. *J Biochem* 148: 13–24.
- Natarajan V, Scribner WM, Hart CM, Parthasarathy S (1995). Oxidized low density lipoprotein-mediated activation of phospholipase D in smooth muscle cells: a possible role in cell proliferation and atherogenesis. *J Lipid Res* 36: 2005–2016.
- van Nieuw Amerongen GP, Vermeer MA, van Hinsbergh VW (2000). Role of RhoA and Rho kinase in lysophosphatidic acid-induced endothelial barrier dysfunction. *Arterioscler Thromb Vasc Biol* 20: E127–E133.
- Nishimasu H, Okudaira S, Hama K, Mihara E, Dohmae N, Inoue A *et al.* (2011). Crystal structure of autotaxin and insight into GPCR activation by lipid mediators. *Nat Struct Mol Biol* 18: 205–212.
- Owens GK, Kumar MS, Wamhoff BR (2004). Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol Rev* 84: 767–801.
- Oyenebraten I (2004). Rapid chemokine secretion from endothelial cells originates from 2 distinct compartments. *Blood* 104: 314–320.
- Packard CJ, O'Reilly DS, Caslake MJ, McMahon AD, Ford I, Cooney J *et al.* (2000). Lipoprotein-associated phospholipase A2 as an independent predictor of coronary heart disease. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 343: 1148–1155.
- Palmethofer A, Robson SC, Nehls V (1999). Lysophosphatidic acid activates nuclear factor kappa B and induces proinflammatory gene expression in endothelial cells. *Thromb Haemost* 82: 1532–1537.

- Pamuklar Z, Lee JS, Cheng HY, Panchatcharam M, Steinhubl S, Morris AJ *et al.* (2008). Individual heterogeneity in platelet response to lysophosphatidic acid: evidence for a novel inhibitory pathway. *Arterioscler Thromb Vasc Biol* 28: 555–561.
- Pamuklar Z, Federico L, Liu S, Umezu-Goto M, Dong A, Panchatcharam M *et al.* (2009). Autotaxin/lysopholipase D and lysophosphatidic acid regulate murine hemostasis and thrombosis. *J Biol Chem* 284: 7385–7394.
- Panchatcharam M, Miriyala S, Yang F, Rojas M, End C, Vallant C *et al.* (2008). Lysophosphatidic acid receptors 1 and 2 play roles in regulation of vascular injury responses but not blood pressure. *Circ Res* 103: 662–670.
- Pandey D, Goyal P, Bamburg JR, Siess W (2006). Regulation of LIM-kinase 1 and cofilin in thrombin-stimulated platelets. *Blood* 107: 575–583.
- Pandey D, Goyal P, Siess W (2007). Lysophosphatidic acid stimulation of platelets rapidly induces Ca²⁺-dependent dephosphorylation of cofilin that is independent of dense granule secretion and aggregation. *Blood Cells Mol Dis* 38: 269–279.
- Panetti TS, Nowlen J, Mosher DF (2000). Sphingosine-1-phosphate and lysophosphatidic acid stimulate endothelial cell migration. *Arterioscler Thromb Vasc Biol* 20: 1013–1019.
- Panetti TS, Hannah DF, Avraamides C, Gaughan JP, Marcinkiewicz C, Huttenlocher A *et al.* (2004). Extracellular matrix molecules regulate endothelial cell migration stimulated by lysophosphatidic acid. *J Thromb Haemost* 2: 1645–1656.
- Penz S, Reininger AJ, Brandl R, Goyal P, Rabie T, Bernlochner I *et al.* (2005). Human atheromatous plaques stimulate thrombus formation by activating platelet glycoprotein VI. *FASEB J* 19: 898–909.
- Penz SM, Reininger AJ, Toth O, Deckmyn H, Brandl R, Siess W (2007). Glycoprotein Ib/alpha inhibition and ADP receptor antagonists, but not aspirin, reduce platelet thrombus formation in flowing blood exposed to atherosclerotic plaques. *Thromb Haemost* 97: 435–443.
- Petta S, Torres D, Fazio G, Camma C, Cabibi D, Di Marco V *et al.* (2011). Carotid atherosclerosis and chronic hepatitis C: a prospective study of risk associations. *Hepatology* 55: 1317–1323.
- Portman OW, Soltys P, Alexander M, Osuga T (1970). Metabolism of lysolecithin in vivo: effects of hyperlipemia and atherosclerosis in squirrel monkeys. *J Lipid Res* 11: 596–604.
- Ptaszynska MM, Pendrak ML, Stracke ML, Roberts DD (2010). Autotaxin signaling via lysophosphatidic acid receptors contributes to vascular endothelial growth factor-induced endothelial cell migration. *Mol Cancer Res* 8: 309–321.
- Purnak T, Efe C, Beyazit Y, Ozaslan E, Astan R, Milanoglu A *et al.* (2011). Recent insights into the relationship between inflammatory liver diseases and atherosclerosis. *J Invest Med* 59: 904–911.
- Reininger AJ, Bernlochner I, Penz SM, Ravanat C, Smethurst P, Farndale RW *et al.* (2010). A 2-step mechanism of arterial thrombus formation induced by human atherosclerotic plaques. *J Am Coll Cardiol* 55: 1147–1158.
- Retzer M, Essler M (2000). Lysophosphatidic acid-induced platelet shape change proceeds via Rho/Rho kinase-mediated myosin light-chain and moesin phosphorylation. *Cell Signal* 12: 645–648.
- Rizza C, Leitinger N, Yue J, Fischer DJ, Wang DA, Shih PT *et al.* (1999). Lysophosphatidic acid as a regulator of endothelial/leukocyte interaction. *Lab Invest* 79: 1227–1235.
- Rosenson RS (2010). Phospholipase A2 inhibition and atherosclerotic vascular disease: prospects for targeting secretory and lipoprotein-associated phospholipase A2 enzymes. *Curr Opin Lipidol* 21: 473–480.
- Rother E, Brandl R, Baker DL, Goyal P, Gebhard H, Tigyi G *et al.* (2003). Subtype-selective antagonists of lysophosphatidic acid receptors inhibit platelet activation triggered by the lipid core of atherosclerotic plaques. *Circulation* 108: 741–747.
- Samadi N, Bekele R, Capatos D, Venkatraman G, Sariahmetoglu M, Brindley DN (2011). Regulation of lysophosphatidate signaling by autotaxin and lipid phosphate phosphatases with respect to tumor progression, angiogenesis, metastasis and chemo-resistance. *Biochimie* 93: 61–70.
- Sanchez-Quesada JL, Benitez S, Otal C, Franco M, Blanco-Vaca F, Ordonez-Llanos J (2002). Density distribution of electronegative LDL in normolipemic and hyperlipemic subjects. *J Lipid Res* 43: 699–705.
- Sanchez-Quesada JL, Camacho M, Anton R, Benitez S, Vila L, Ordonez-Llanos J (2003). Electronegative LDL of FH subjects: chemical characterization and induction of chemokine release from human endothelial cells. *Atherosclerosis* 166: 261–270.
- Sano T, Baker D, Virag T, Wada A, Yatomi Y, Kobayashi T *et al.* (2002). Multiple mechanisms linked to platelet activation result in lysophosphatidic acid and sphingosine 1-phosphate generation in blood. *J Biol Chem* 277: 21197–21206.
- Sarker MH, Hu DE, Fraser PA (2010). Regulation of cerebrovascular permeability by lysophosphatidic acid. *Microcirculation* 17: 39–46.
- Saulnier-Blache JS, Girard A, Simon MF, Lafontan M, Valet P (2000). A simple and highly sensitive radioenzymatic assay for lysophosphatidic acid quantification. *J Lipid Res* 41: 1947–1951.
- Schmitz G, Ruebsaamen K (2010). Metabolism and atherogenic disease association of lysophosphatidylcholine. *Atherosclerosis* 208: 10–18.
- Schmitz U, Thommes K, Beier I, Vetter H (2002). Lysophosphatidic acid stimulates p21-activated kinase in vascular smooth muscle cells. *Biochem Biophys Res Commun* 291: 687–691.
- Schober A (2008). Chemokines in vascular dysfunction and remodeling. *Arterioscler Thromb Vasc Biol* 28: 1950–1959.
- Schulz C, Penz S, Hoffmann C, Langer H, Gillitzer A, Schneider S *et al.* (2008). Platelet GPVI binds to collagenous structures in the core region of human atheromatous plaque and is critical for atheroprotection in vivo. *Basic Res Cardiol* 103: 356–367.
- Schumacher KA, Classen HG, Spath M (1979). Platelet aggregation evoked in vitro and in vivo by phosphatidic acids and lysoderivatives: identity with substances in aged serum (DAS). *Thromb Haemost* 42: 631–640.
- Schwartz D, Andalibi A, Chaverri-Almada L, Berliner JA, Kirchgessner T, Fang ZT *et al.* (1994). Role of the GRO family of chemokines in monocyte adhesion to MM-LDL-stimulated endothelium. *J Clin Invest* 94: 1968–1973.
- Schwartz SM, deBlois D, O'Brien ER (1995). The intima. Soil for atherosclerosis and restenosis. *Circ Res* 77: 445–465.
- Seewald S, Sachinidis A, Dusing R, Ko Y, Seul C, Epping P *et al.* (1997). Lysophosphatidic acid and intracellular signalling in vascular smooth muscle cells. *Atherosclerosis* 130: 121–131.
- Seewald S, Schmitz U, Seul C, Ko Y, Sachinidis A, Vetter H (1999). Lysophosphatidic acid stimulates protein kinase C isoforms alpha, beta, epsilon, and zeta in a pertussis toxin sensitive pathway in vascular smooth muscle cells. *Am J Hypertens* 12: 532–537.

- Shan L, Jaffe K, Li S, Davis L (2008a). Quantitative determination of lysophosphatidic acid by LC/ESI/MS/MS employing a reversed phase HPLC column. *J Chromatogr B Analyt Technol Biomed Life Sci* 864: 22–28.
- Shan L, Li S, Jaffe K, Davis L (2008b). Quantitative determination of cyclic phosphatidic acid in human serum by LC/ESI/MS/MS. *J Chromatogr B Analyt Technol Biomed Life Sci* 862: 161–167.
- Siess W (2002). Athero- and thrombogenic actions of lysophosphatidic acid and sphingosine-1-phosphate. *Biochim Biophys Acta* 1582: 204–215.
- Siess W (2006). Platelet interaction with bioactive lipids formed by mild oxidation of low-density lipoprotein. *Pathophysiol Haemost Thromb* 35: 292–304.
- Siess W, Tigyi G (2004). Thrombogenic and atherogenic activities of lysophosphatidic acid. *J Cell Biochem* 92: 1086–1094.
- Siess W, Zangl KJ, Essler M, Bauer M, Brandl R, Corrinth C *et al.* (1999). Lysophosphatidic acid mediates the rapid activation of platelets and endothelial cells by mildly oxidized low density lipoprotein and accumulates in human atherosclerotic lesions. *Proc Natl Acad Sci U S A* 96: 6931–6936.
- Simon MF, Chap H, Douste-Blazy L (1982). Human platelet aggregation induced by 1-alkyl-lysophosphatidic acid and its analogs: a new group of phospholipid mediators? *Biochem Biophys Res Commun* 108: 1743–1750.
- Smyth SS, Cheng HY, Miriyala S, Panchatcharam M, Morris AJ (2008). Roles of lysophosphatidic acid in cardiovascular physiology and disease. *Biochim Biophys Acta* 1781: 563–570.
- Stafforini DM (2009). Biology of platelet-activating factor acetylhydrolase (PAF-AH, lipoprotein associated phospholipase A2). *Cardiovasc Drugs Ther* 23: 73–83.
- Steinbrecher UP, Parthasarathy S, Leake DS, Witztum JL, Steinberg D (1984). Modification of low density lipoprotein by endothelial cells involves lipid peroxidation and degradation of low density lipoprotein phospholipids. *Proc Natl Acad Sci U S A* 81: 3883–3887.
- Stenmark KR, Fagan KA, Frid MG (2006). Hypoxia-induced pulmonary vascular remodeling: cellular and molecular mechanisms. *Circ Res* 99: 675–691.
- Stocker R, Keaney JF, Jr (2004). Role of oxidative modifications in atherosclerosis. *Physiol Rev* 84: 1381–1478.
- Subramanian P, Karshovska E, Reinhard P, Megens RTA, Zhou Z, Akhtar S *et al.* (2010). Lysophosphatidic acid receptors LPA1 and LPA3 promote CXCL12-mediated smooth muscle progenitor cell recruitment in neointima formation. *Circ Res* 107: 96–105.
- Swirski FK (2006). Monocyte accumulation in mouse atherogenesis is progressive and proportional to extent of disease. *Proc Natl Acad Sci U S A* 103: 10340–10345.
- Tanaka M, Okudaira S, Kishi Y, Ohkawa R, Iseki S, Ota M *et al.* (2006). Autotaxin stabilizes blood vessels and is required for embryonic vasculature by producing lysophosphatidic acid. *J Biol Chem* 281: 25822–25830.
- Thim T, Hagensen MK, Bentzon JF, Falk E (2008). From vulnerable plaque to atherothrombosis. *J Intern Med* 263: 506–516.
- Thompson A, Gao P, Orfei L, Watson S, Di Angelantonio E, Kaptoge S *et al.* (2010). Lipoprotein-associated phospholipase A(2) and risk of coronary disease, stroke, and mortality: collaborative analysis of 32 prospective studies. *Lancet* 375: 1536–1544.
- Thumser AE, Voysey JE, Wilton DC (1994). The binding of lysophospholipids to rat liver fatty acid-binding protein and albumin. *Biochem J* 301: (Pt 3): 801–806.
- Tigyi G (2001). Physiological responses to lysophosphatidic acid and related glycerophospholipids. *Prostaglandins Other Lipid Mediat* 64: 47–62.
- Tigyi G (2010). Aiming drug discovery at lysophosphatidic acid targets. *Br J Pharmacol* 161: 241–270.
- Tigyi G, Miledi R (1992). Lysophosphatidates bound to serum albumin activate membrane currents in *Xenopus* oocytes and neurite retraction in PC12 pheochromocytoma cells. *J Biol Chem* 267: 21360–21367.
- Tigyi G, Henschen A, Miledi R (1991). A factor that activates oscillatory chloride currents in *Xenopus* oocytes copurifies with a subfraction of serum albumin. *J Biol Chem* 266: 20602–20609.
- Tokumura A (1995). A family of phospholipid autacoids: occurrence, metabolism and bioactions. *Prog Lipid Res* 34: 151–184.
- Tokumura A, Fukuzawa K, Isobe J, Tsukatani H (1981). Lysophosphatidic acid-induced aggregation of human and feline platelets: structure-activity relationship. *Biochem Biophys Res Commun* 99: 391–398.
- Tokumura A, Iimori M, Nishioka Y, Kitahara M, Sakashita M, Tanaka S (1994). Lysophosphatidic acids induce proliferation of cultured vascular smooth muscle cells from rat aorta. *Am J Physiol* 267 (1 Pt 1): C204–C210.
- Tokumura A, Kanaya Y, Kitahara M, Miyake M, Yoshioka Y, Fukuzawa K (2002a). Increased formation of lysophosphatidic acids by lysophospholipase D in serum of hypercholesterolemic rabbits. *J Lipid Res* 43: 307–315.
- Tokumura A, Kanaya Y, Miyake M, Yamano S, Irahara M, Fukuzawa K (2002b). Increased production of bioactive lysophosphatidic acid by serum lysophospholipase D in human pregnancy. *Biol Reprod* 67: 1386–1392.
- Tokumura A, Majima E, Kariya Y, Tominaga K, Kogure K, Yasuda K *et al.* (2002c). Identification of human plasma lysophospholipase D, a lysophosphatidic acid-producing enzyme, as autotaxin, a multifunctional phosphodiesterase. *J Biol Chem* 277: 39436–39442.
- Tokumura A, Sinomiya J, Kishimoto S, Tanaka T, Kogure K, Sugiura T *et al.* (2002d). Human platelets respond differentially to lysophosphatidic acids having a highly unsaturated fatty acyl group and alkyl ether-linked lysophosphatidic acids. *Biochem J* 365 (Pt 3): 617–628.
- Tomsig JL, Snyder AH, Berdyshev EV, Skobeleva A, Mataya C, Natarajan V *et al.* (2009). Lipid phosphate phosphohydrolase type 1 (LPP1) degrades extracellular lysophosphatidic acid in vivo. *Biochem J* 419: 611–618.
- Toschi V, Gallo R, Lettino M, Fallon JT, Gertz SD, Fernandez-Ortiz A *et al.* (1997). Tissue factor modulates the thrombogenicity of human atherosclerotic plaques. *Circulation* 95: 594–599.
- Tsuda S, Okudaira S, Moriya-Ito K, Shimamoto C, Tanaka M, Aoki J *et al.* (2006). Cyclic phosphatidic acid is produced by autotaxin in blood. *J Biol Chem* 281: 26081–26088.
- Tsukahara T, Tsukahara R, Fujiwara Y, Yue J, Cheng Y, Guo H *et al.* (2010). Phospholipase D2-dependent inhibition of the nuclear hormone receptor PPARgamma by cyclic phosphatidic acid. *Mol Cell* 39: 421–432.

- Umez-Goto M, Kishi Y, Taira A, Hama K, Dohmae N, Takio K *et al.* (2002). Autotaxin has lysophospholipase D activity leading to tumor cell growth and motility by lysophosphatidic acid production. *J Cell Biol* 158: 227–233.
- Wang WY, Zhang J, Wu WY, Li J, Ma YL, Chen WH *et al.* (2011). Inhibition of lipoprotein-associated phospholipase A2 ameliorates inflammation and decreases atherosclerotic plaque formation in ApoE-deficient mice. *PLoS ONE* 6: e23425.
- Watanabe N, Ikeda H, Nakamura K, Ohkawa R, Kume Y, Aoki J *et al.* (2007a). Both plasma lysophosphatidic acid and serum autotaxin levels are increased in chronic hepatitis C. *J Clin Gastroenterol* 41: 616–623.
- Watanabe N, Ikeda H, Nakamura K, Ohkawa R, Kume Y, Tomiya T *et al.* (2007b). Plasma lysophosphatidic acid level and serum autotaxin activity are increased in liver injury in rats in relation to its severity. *Life Sci* 81: 1009–1015.
- Watson SP, McConnell RT, Lapetina EG (1985). Decanoyl lysophosphatidic acid induces platelet aggregation through an extracellular action. Evidence against a second messenger role for lysophosphatidic acid. *Biochem J* 232: 61–66.
- Weber C, Noels H (2011). Atherosclerosis: current pathogenesis and therapeutic options. *Nat Med* 17: 1410–1422.
- Weber KS, von Hundelshausen P, Clark-Lewis I, Weber PC, Weber C (1999). Differential immobilization and hierarchical involvement of chemokines in monocyte arrest and transmigration on inflamed endothelium in shear flow. *Eur J Immunol* 29: 700–712.
- Wiesner P, Leidl K, Boettcher A, Schmitz G, Liebisch G (2009). Lipid profiling of FPLC-separated lipoprotein fractions by electrospray ionization tandem mass spectrometry. *J Lipid Res* 50: 574–585.
- Williams JR, Khandoga AL, Goyal P, Fells JL, Perygin DH, Siess W *et al.* (2009). Unique ligand selectivity of the GPR92/LPA5 lysophosphatidate receptor indicates role in human platelet activation. *J Biol Chem* 284: 17304–17319.
- Xu YJ, Rath SS, Chapman DC, Arneja AS, Dhalla NS (2003). Mechanisms of lysophosphatidic acid-induced DNA synthesis in vascular smooth muscle cells. *J Cardiovasc Pharmacol* 41: 381–387.
- Yanagida K, Ishii S (2011). Non-Edg family LPA receptors: the cutting edge of LPA research. *J Biochem* 150: 223–232.
- Yoshida K, Nishida W, Hayashi K, Ohkawa Y, Ogawa A, Aoki J *et al.* (2003). Vascular remodeling induced by naturally occurring unsaturated lysophosphatidic acid in vivo. *Circulation* 108: 1746–1752.
- van Zanten GH, de Graaf S, Slootweg PJ, Heijnen HF, Connolly TM, de Groot PG *et al.* (1994). Increased platelet deposition on atherosclerotic coronary arteries. *J Clin Invest* 93: 615–632.
- Zaslavsky A, Singh LS, Tan H, Ding H, Liang Z, Xu Y (2006). Homo- and hetero-dimerization of LPA/S1P receptors, OGR1 and GPR4. *Biochim Biophys Acta* 1761: 1200–1212.
- Zeiffer U, Schober A, Lietz M, Liehn EA, Erl W, Emans N *et al.* (2004). Neointimal smooth muscle cells display a proinflammatory phenotype resulting in increased leukocyte recruitment mediated by P-selectin and chemokines. *Circ Res* 94: 776–784.
- Zernecke A, Schober A, Bot I, von Hundelshausen P, Liehn EA, Mopps B *et al.* (2005). SDF-1 α /CXCR4 axis is instrumental in neointimal hyperplasia and recruitment of smooth muscle progenitor cells. *Circ Res* 96: 784–791.
- Zhang C, Baker DL, Yasuda S, Makarova N, Balazs L, Johnson LR *et al.* (2004). Lysophosphatidic acid induces neointima formation through PPAR γ activation. *J Exp Med* 199: 763–774.
- Zhang LL, Gao CY, Fang CQ, Wang YJ, Gao D, Yao GE *et al.* (2011). PPAR γ attenuates intimal hyperplasia by inhibiting TLR4-mediated inflammation in vascular smooth muscle cells. *Cardiovasc Res* 92: 484–493.
- Zhao Z, Xu Y (2009). Measurement of endogenous lysophosphatidic acid by ESI-MS/MS in plasma samples requires pre-separation of lysophosphatidylcholine. *J Chromatogr B Analyt Technol Biomed Life Sci* 877: 3739–3742.
- Zhou D, Luini W, Bernasconi S, Diomedea L, Salmons M, Mantovani A *et al.* (1995). Phosphatidic acid and lysophosphatidic acid induce haptotactic migration of human monocytes. *J Biol Chem* 270: 25549–25556.
- Zhou Z, Niu J, Zhang Z (2010). The role of lysophosphatidic acid receptors in phenotypic modulation of vascular smooth muscle cells. *Mol Biol Rep* 37: 2675–2686.
- Zhou Z, Subramanian P, Sevilimis G, Globke B, Soehnlein O, Karshovska E *et al.* (2011). Lipoprotein-derived lysophosphatidic acid promotes atherosclerosis by releasing CXCL1 from the endothelium. *Cell Metab* 13: 592–600.
- Zhou ZB, Niu JP, Zhang ZJ (2009). Receptor-mediated vascular smooth muscle migration induced by LPA involves p38 mitogen-activated protein kinase pathway activation. *Int J Mol Sci* 10: 3194–3208.