

REVIEW

Roles of proteolysis in
regulation of GPCR function

GS Cottrell

Reading School of Pharmacy, University of Reading, Reading, UK

Correspondence

Graeme S Cottrell, Reading
School of Pharmacy, University
of Reading, Whiteknights,
Reading RG6 6UB, UK. E-mail:
g.s.cottrell@reading.ac.uk

Keywords

peptidase; neuropeptide; GPCR;
trafficking; signalling; proteolysis;
endosome; cell-surface; ubiquitin;
 β -arrestins

Received

11 July 2012

Revised

3 September 2012

Accepted

24 September 2012

The enzymatic activity of peptidases must be tightly regulated to prevent uncontrolled hydrolysis of peptide bonds, which could have devastating effects on biological systems. Peptidases are often generated as inactive propeptidases, secreted with endogenous inhibitors, or they are compartmentalized. Propeptidases become active after proteolytic removal of N-terminal activation peptides by other peptidases. Some peptidases only become active towards substrates only at certain pHs, thus confining activity to specific compartments or conditions. This review discusses the different roles proteolysis plays in regulating GPCRs. At the cell-surface, certain GPCRs are regulated by the hydrolytic inactivation of bioactive peptides by membrane-anchored peptidases, which prevent signalling. Conversely, cell-surface peptidases can also generate bioactive peptides, which directly activate GPCRs. Alternatively, cell-surface peptidases activated by GPCRs, can generate bioactive peptides to cause transactivation of receptor tyrosine kinases, thereby promoting signalling. Certain peptidases can signal directly to cells, by cleaving GPCR to initiate intracellular signalling cascades. Intracellular peptidases also regulate GPCRs; lysosomal peptidases destroy GPCRs in lysosomes to permanently terminate signalling and mediate down-regulation; endosomal peptidases cleave internalized peptide agonists to regulate GPCR recycling, resensitization and signalling; and soluble intracellular peptidases also participate in GPCR function by regulating the ubiquitination state of GPCRs, thereby altering GPCR signalling and fate. Although the use of peptidase inhibitors has already brought success in the treatment of diseases such as hypertension, the discovery of new regulatory mechanisms involving proteolysis that control GPCRs may provide additional targets to modulate dysregulated GPCR signalling in disease.

Abbreviations

ADAM, a disintegrin and metallopeptidase; CGRP, calcitonin gene-related peptide; CLR, calcitonin receptor-like receptor; CRF₁ receptor, corticotropin-releasing factor 1 receptor; CXCR, C-X-C chemokine receptor; ECE-1, endothelin-converting enzyme-1; EGF, epidermal growth factor; HB-EGF, membrane-anchored heparin-binding EGF-like growth factor; NEP, neprilysin; NK receptor, neurokinin receptor; NTS₁ receptor, neurotensin 1 receptor; PAR, proteinase-activated receptor; PP2A, protein phosphatase 2A; RAMP, receptor activity-modifying protein; sst receptor, somatostatin receptor; SP, substance P; USP, ubiquitin-specific peptidase; Unc1, urocortin 1

Nomenclature

For receptors, the British Journal of Pharmacology's *Guide to Receptors and Channels* was used (Alexander *et al.*, 2011). For peptidases, the MEROPS database was used (Rawlings *et al.*, 2012).

The traditional notion of proteolysis

Proteolysis is the hydrolytic breakdown of a peptide bond, C(O)-NH found between amino acids in peptides, poly-

peptides and protein structures. Peptide bonds can spontaneously break in the presence of water, but do so, at an extremely slow rate. Therefore, in biological systems, enzymes are required to facilitate the breakage of these bonds. These enzymes or more specifically, peptidases (also known as proteases or proteinases) are themselves made of amino acids and are currently classified into six groups according to the critical residue in their catalytic site: (i) serine peptidases; (ii) cysteine peptidases; (iii) aspartic peptidases; (iv) threonine peptidases; (v) metallopeptidases; and (vi) glutamic peptidases (Rawlings *et al.*, 2012). Traditionally, peptidases are mainly thought of as enzymes of digestion,

breaking down food in the stomach and in the intestine. However, it is now clear that peptidases can contribute to the regulation of cell function by controlling levels of bioactive peptides and by cleaving cell-surface receptors and ion channels to regulate signalling pathways.

Cell-surface peptidases regulate the availability of GPCR ligands

The role of cell-surface peptides in the regulation of GPCR signalling is well documented. These peptidases regulate levels of circulating bioactive peptides, which function to initiate GPCR-mediated signalling. ACE compound peptidase (EC 3.4.15.1) is a zinc-dependent metallopeptidase that converts the inactive angiotensin I to angiotensin II by releasing the C-terminal dipeptide, His⁹-Leu¹⁰ (Figure 1A). Angiotensin II is a vasoconstrictor and exerts its effect through two types of angiotensin II receptors, AT₁ and AT₂. The AT₁ receptor mediates most of the physiological and pathophysiological actions of angiotensin II and is the predominant receptor subtype expressed in the cardiovascular




Figure 1

Cell-surface peptidases regulate GPCR-mediated signalling. (A) ACE compound peptidase cleaves angiotensin I to generate angiotensin II to promote activation of angiotensin II receptors. Conversely, NEP hydrolyses SP to prevent activation of SP of neurokinin 1 receptors. (B) Activation of certain GPCRs promotes the ADAMs that generate EGF-like ligands (e.g. heparin-binding EGF-like factor, amphiregulin). In turn, these ligands transactivate ErbB receptors to activate intracellular signalling pathways.

system. Interaction of the AT₁ receptor with angiotensin II activates G_{q/11}, G_i, G₁₂ and G₁₃ proteins, leading to the mobilization of intracellular calcium, generation of reactive oxygen species and activation of numerous PKs and mitogenic signalling pathways [reviewed in Mogi *et al.* (2009)]. AT₂ receptors couple to G proteins to activate PLC, promoting the mobilization of intracellular calcium and activation of PKC [reviewed in Porrello *et al.* (2009)]. Angiotensin II causes a plethora of effects, including tissue remodelling, leukocyte infiltration, inflammation, atherosclerosis, endothelial dysfunction, myocardial infarction, stroke, and heart and renal failure [reviewed in Cheng *et al.* (2005)]. Similarly, endothelin-converting enzyme-1 (ECE-1, EC 3.4.24.71) is responsible for the production of the vasoconstrictor endothelin-1 from big endothelin. Endothelin-1 can activate both endothelin A (ET_A) and ET_B receptors to elicit a broad range of signalling responses [reviewed in Khimji and Rockey (2010)]. Activation of ET_A receptors by endothelin-1 promotes mobilization of intracellular calcium, indicating that ET_A receptors are coupled to G_{q/11} proteins. ET_A receptors primarily mediate the vasoactive and proliferative effects of endothelin-1. ET_B receptors have been proposed to act as endothelin-1 scavengers and thus, reduce circulating endothelin-1 levels. Thus, proteolysis acts to promote activation of GPCRs. In contrast, proteolysis can also act to prevent activation of GPCRs. For example, neprilysin (NEP, neutral endopeptidase 24.11, EC 3.4.24.11) cleaves and inactivates both bradykinin (Gly⁴↓Phe⁵ and Pro⁷↓Phe⁸) and substance P (SP, Gln⁶↓Phe↓Phe↓Gly↓Leu¹⁰) (Matsas *et al.*, 1984), thus preventing activation of their respective GPCRs, the bradykinin 2 receptor (B₂ receptor) and the neurokinin 1 receptor (NK₁ receptor) (Figure 1A). Peptidases can therefore play a major role in the production of vasoactive peptides and, therefore, regulate vascular functions and dysregulation can lead to vascular diseases. A great deal of time and effort has been spent on the development of peptidase inhibitors for the treatment of hypertension. It is over 30 years since the first ACE inhibitor, captopril, was designed (Ondetti *et al.*, 1977). Now, however, other ACE inhibitors with improved pharmacokinetics and pharmacodynamics have since been developed and include enalaprilat (MK-421) (Gross *et al.*, 1981) and imidaprilat (Ikeo *et al.*, 1992). ACE inhibitors are effective treatments for hypertension and congestive heart failure (CONSENSUS, 1987; SOLVD, 1991) and are also beneficial for patients with atherosclerosis (Yusuf *et al.*, 2000). With the success of ACE inhibitors for the treatment of hypertension, it was thought that the development of compounds able to inhibit multiple peptidases, thereby potentiating the effects of dilator peptides such as bradykinin, while reducing the availability of constrictors such as angiotensin II, may offer an even better way of treating diseases such as hypertension. Indeed, dual or triple vasoactive peptidase inhibitors have been developed. These compounds inhibit the proteolytic activities of ACE, ECE-1 and NEP in various combinations. The first described dual inhibitors were alatriopril and glycoprilat. Both compounds inhibit the activities of both ACE and NEP (Gros *et al.*, 1991). Alatriopril was shown to be more effective than captopril alone in reducing cardiac hypertrophy in rats with myocardial infarction (Bralet *et al.*, 1994). Later, omapatrilat (BMS-186716) was developed and was shown to

Table 1

Future and current uses for peptidase inhibitors in the treatment of disease

Peptidase	Drug perspectives	Targeted GPCRs	References/reviews
Cell-surface			
ACE, ECE-1, NEP ^a	Hypertension	AT ₁ , AT ₂ , ET _A , ET _B , B ₂	Dive <i>et al.</i> (2009)
NEP, aminopeptidase N	Pain	Opioid receptors	Thanawala <i>et al.</i> (2008)
ADAMs	Cancer progression	GPCRs causing ErbB receptor transactivation	Paolillo and Schinelli (2008)
Extracellular			
Thrombin, Coagulation Factor Xa	Anticoagulants, thrombosis	PAR ₁ , PAR ₃ , PAR ₄	Showkathali and Natarajan (2012)
Trypsin, Trypsinase	Inflammatory bowel diseases	PAR ₂	Rothmeier and Ruf (2012; Yoshida and Yoshikawa (2008)
Lysosomal			
Cathepsins B, C, D, E, G and L, Carboxypeptidase A	No known functional consequences of preventing GPCR degradation	— ^b	N/A
Endosomal			
ECE-1	Modulation of GPCR trafficking and signalling	NK ₁ , CLR•RAMP1, sst ₂ , NT ₁ , CRF ₁	Murphy <i>et al.</i> (2009; 2011)
USPs	Modulation of GPCR trafficking and signalling	GPCR function affected by de-ubiquitination of β-arrestins	Murphy <i>et al.</i> (2009; Shenoy <i>et al.</i> (2009)

^aDual and triple ACE, ECE-1 and NEP inhibitors are in development.

^bToo numerous to list in table. N/A, not applicable.

be more effective in reducing blood pressure in humans than either placebo or ACE inhibitors (Neal *et al.*, 2002; Regamey *et al.*, 2002). However, omapatrilat failed in phase III clinical trials and was discontinued due to an increased incidence of angioedema as an unwanted side effect. CGS 35601, a triple vasopeptidase inhibitor, prevents the activities of ACE, NEP and ECE-1 (Trapani *et al.*, 2004) and significantly reduced both systolic and diastolic blood pressure in a number of preclinical rodent models of hypertension (Daull *et al.*, 2005; 2006a). Further, a preclinical safety profile assessment of CGS 35601 showed it to have no effect on either hepatic or liver toxicities (Daull *et al.*, 2006b). Although these triple inhibitors may represent the future of peptidase inhibitors for the treatment of disease (Table 1), no clinical trials using triple peptidase inhibitors have yet been conducted. So, although these dual and triple peptidase inhibitor compounds are promising in humans and animal models of hypertension, none have yet been approved for the treatment of human disease. Thus, ACE inhibitors remain the compounds of choice for the treatment of hypertension, often in combination with angiotensin or β-adrenoceptor antagonists or diuretics. NEP also plays a major role in the catabolism of endogenous opioid peptides such as the enkephalins and dynorphins. Thus, together with aminopeptidase N (EC 3.4.11.2), NEP represents a major target for the development of drugs for the treatment of acute and chronic pain [reviewed in Thanawala *et al.* (2008)].

GPCRs promote proteolysis to transactivate epidermal growth factor receptors

Certain GPCRs can activate intracellular signalling pathways via transactivation of cell-surface receptor tyrosine kinases such as PDGF (Linseman *et al.*, 1995), insulin-like growth factor-1 receptor (Rao *et al.*, 1995), ErbB receptor (Daub *et al.*, 1996) and Trk neurotrophin receptors (Lee and Chao, 2001). Interestingly, only transactivation of ErbB receptors requires the activity of peptidases (Figure 1B). The GPCR agonists, endothelin-1, lysophosphatic acid and thrombin, were shown to induce phosphorylation of the ErbB receptor in Rat-1 cells (Daub *et al.*, 1996). Specific inhibition of ErbB receptors using the selective tyrphostin AG1478 or expression of a dominant-negative ErbB receptor mutant that is unable to signal, suppressed the GPCR-induced activation of ERK1/2 (Daub *et al.*, 1996). Later studies revealed that transactivation of ErbB receptors, involves the processing of membrane-anchored heparin-binding EGF-like growth factor by a metalloproteinase activity that is rapidly induced following GPCR activation. Inhibition of the peptidase activity using the metalloproteinase inhibitor batimastat prevented ErbB receptor transactivation and downstream signals (Prenzel *et al.*, 1999). Additional work, indicated that this GPCR-mediated transactivation not only lead to the initiation of signalling cascades, but also ErbB receptor dimerization, tyrosine autophosphor-

ylation and internalization (Maudsley *et al.*, 2000). The identity of the metallopeptidase responsible for the processing of the membrane-anchored ligand remained elusive until a study conducted in cardiomyocytes showed that the metallopeptidase inhibitor, KB-R7785 and a dominant-negative of a disintegrin and metallopeptidase (ADAM) 12 peptidase attenuated GPCR-induced signalling (Asakura *et al.*, 2002). The same study also showed that KB-R7785 inhibited the shedding of heparin-binding EGF-like growth factor and attenuated thoracic aortic constriction-induced thickening of the heart muscle in intact mice. A subsequent study suggested that ADAM10 peptidase (EC 3.4.24.81) is also implicated in bombesin-mediated ErbB receptor transactivation (Yan *et al.*, 2002). As more than one peptidase is involved in ErbB receptor transactivation, it came as no surprise that additional ligands were also involved. In squamous cell carcinoma cells, the GPCR agonists, lysophosphatic acid and carbachol, which activate lysophospholipid and muscarinic receptors, respectively, specifically activate metallopeptidase-dependent release of amphiregulin by another ADAM peptidase, ADAM17 peptidase (TNF- α -converting enzyme, TACE; EC 3.4.24.86) to regulate proliferation and motility (Gschwind *et al.*, 2003). However, in TccSup cancer cells, lysophosphatic acid-induced transactivation is mediated by ADAM15 peptidase and promotes cell survival (Schafer *et al.*, 2004).

Following on from the fact that GPCRs can promote the activity of many different metallopeptidases to cause ErbB receptor transactivation, evidence exists for the involvement of numerous GPCR-induced pathways in the generation of EGF-like ligands [reviewed in Ohtsu *et al.* (2006)]. The β_2 -adrenoceptor agonist, isoprenaline causes ErbB receptor transactivation via a mechanism that requires G $\beta\gamma$ subunits and c-Src activity (Pierce *et al.*, 2001). Similarly, glucagon-like peptide-1-induced transactivation also requires c-Src activity (Buteau *et al.*, 2003). The involvement of G proteins has also been observed in other studies. Lysophosphatic acid-mediated transactivation by ADAM17 was partially blocked by pertussis toxin in squamous cell carcinoma-9 cells (Gschwind *et al.*, 2003) and in MDA-MB-231 cells, lysophosphatic acid- and sphingosine-1 phosphate-dependent ErbB receptor transactivation involving ADAM15, was also prevented by pertussis toxin (Hart *et al.*, 2005). A requirement for the activity of phospholipase C has also been observed for angiotensin II-induced, ADAM17-dependent ErbB receptor transactivation by the AT $_1$ receptor (Mifune *et al.*, 2005). The elevation of intracellular calcium levels by bradykinin (Zwick *et al.*, 1999) and angiotensin II (Eguchi *et al.*, 2003) has also been implicated in the transactivation of ErbB receptors. It is assumed that many of these GPCR-induced pathways lead to the phosphorylation of the metallopeptidases, enhancing the proteolytic activity of the peptidase. Although it is well established that numerous PKs can phosphorylate metallopeptidases to promote proteolytic activity [reviewed in Huovila *et al.* (2005)], none of the present studies have presented evidence of a direct GPCR-mediated effect on metallopeptidases.

The generation of reactive oxygen species by GPCRs has also been proposed to play an important role in promoting metallopeptidase activity and includes ErbB receptor transactivation mediated by 5-HT $_{2B}$ receptors and α_{1D} -adrenoceptors (Pietri *et al.*, 2005), AT $_1$ receptor (Eguchi and Inagami, 2000)

and purine P2Y receptors (Myers *et al.*, 2009). The reactive oxygen species are thought to induce oxidation of a cysteine residue that lies within an inhibitory motif, thereby activating the peptidase (Zhang *et al.*, 2001). A more recent development in elucidating the mechanism by which metallopeptidases become active following GPCR activation to promote ErbB receptor activation is the involvement of integrins (Gooz *et al.*, 2012). 5-HT induces kidney mesangial cell proliferation through ADAM17 activation and ErbB receptor transactivation. In unstimulated cells, ADAM17 binds to $\alpha_5\beta_1$ integrin, an interaction that prevents the proteolytic activity of ADAM17. However, following application of 5-HT, this interaction is disrupted, presumably promoting peptidase activity (Gooz *et al.*, 2012).

A further consequence of the processing of EGF-like ligands from cell-surface precursors, is the generation of C-terminal fragments. It is now established that these C-terminal fragments also have a biological role [reviewed in Tanida *et al.* (2010)]. IL-8 induces cell proliferation and migration of the colon cancer cell lines, HT-29 and Caco2 by an ADAM-dependent intranuclear translocation pathway of HB-EGF-C-terminal fragment (Itoh *et al.*, 2005). Thus, understanding how GPCRs activate cell-surface peptidases to generate the multiple ligands responsible for receptor transactivation is essential to develop new pharmacological interventions. The importance of metallopeptidases in generating EGF-like ligands that regulate cell proliferation and differentiation, has lead these peptidases to be considered as useful molecular targets in the treatment of cancer. Specific peptidase inhibitors may also be used to modulate GPCR-induced signalling and this may be an effective treatment for GPCR-driven diseases.

Peptidases can act as positive or negative regulators of GPCR signalling

It was long recognized that peptidases such as thrombin (EC 3.4.21.5) and trypsin could signal directly to cells (Burger, 1970; Sefton and Rubin, 1970; Chen and Buchanan, 1975; Carney and Cunningham, 1977, 1978). However, the mechanism through which these peptidases could activate intracellular signalling cascades remained unknown until the cloning of the thrombin receptor, now called proteinase-activated receptor 1 (PAR $_1$) (Vu *et al.*, 1991). PAR $_1$ is a GPCR and thrombin activates PAR $_1$ by direct cleavage of the receptor in a two-step process. Firstly, thrombin binds to PAR $_1$ either side of the proteolytic cleavage site. One of these sites (D 51 KYEPF 56) is similar to that of hirudin, an anticoagulant found in the saliva of leech (Vu *et al.*, 1991). This hirudin-like binding domain increases the affinity of thrombin for PAR $_1$. Following binding, thrombin cleaves between Arg 41 and Ser 42 to expose the new N-terminus starting with S 42 FLLRN 47 (Vu *et al.*, 1991). This tethered ligand domain then interacts with residues on the second extracellular loop of the receptor and presumably induces a conformational change, which activates the receptor. PAR $_1$ activation by thrombin is the most potent known trigger for platelet aggregation (Sambrano *et al.*, 2001) and as such, thrombin signalling and PAR $_1$ are key targets for the prevention of thrombosis.




Figure 2

Peptidases act as biased agonists at GPCRs. (A) Trypsin cleaves PAR₂ to create a new N-terminus that activates PAR₂ mobilizing intracellular calcium and promoting phosphorylation (p) of ERK1/2. (B) Cathepsin G cleaves PAR₂, but does not elicit any known signaling and prevents activation by other peptidases such as trypsin. (C) Other peptidases such as elastase-2, cleave PAR₂ at a site distinct from trypsin. The action of this peptidase does not mobilize intracellular calcium, but does activate ERK1/2.

PAR₂ is the second member of this receptor family and is activated by trypsin (Figure 2A) (Nystedt *et al.*, 1994; 1995a,b; Bohm *et al.*, 1996). Cleavage of PAR₂ occurs between Arg³⁶ and Ser³⁷ to reveal the tethered ligand and new amino termi-

nus of S³⁷LIGKV⁴². The use of PAR₂-specific agonistic and antagonistic peptides and studies in PAR₂-deficient mice, have helped to identify critical roles for PAR₂ in inflammation, development, angiogenesis and immune responses [reviewed in Rothmeier and Ruf (2012)]. Subsequently, two other members of this subfamily of GPCRs have been cloned, PAR₃ (Ishihara *et al.*, 1997; Scase *et al.*, 1997) and PAR₄ (Kahn *et al.*, 1998; Xu *et al.*, 1998), both of which are activated by thrombin. PAR₄ can also be activated by trypsin (Xu *et al.*, 1998). Due to their sensitivity to thrombin, both PAR₃ and PAR₄ are important in platelet function (Ishihara *et al.*, 1998; Kahn *et al.*, 1998).

PARs are promiscuous receptors and may be activated by multiple peptidases. For example, PAR₁ can also be activated by coagulation factor Xa (EC 3.4.21.6) (Camerer *et al.*, 2000), activated protein C (EC 3.4.21.69) (Riewald *et al.*, 2002), matrix metalloproteinase-1 (EC 3.4.24.7) (Boire *et al.*, 2005) and plasmin (EC 3.4.21.7) (Mannaioni *et al.*, 2008). Similarly, PAR₂ is also cleaved by multiple peptidases, including mast cell tryptases (Molino *et al.*, 1997), coagulation factor VIIa (EC 3.4.21.21) in complex with tissue factor and coagulation factor Xa (Camerer *et al.*, 2000), matriptase-1 (membrane-type serine peptidase 1) (Takeuchi *et al.*, 2000) and trypsin IV (Cottrell *et al.*, 2004). PAR₄ is cleaved and activated by trypsin (Xu *et al.*, 1998), cathepsin G (Sambrano *et al.*, 2000), trypsin IV (Cottrell *et al.*, 2004) and plasmin (Quinton *et al.*, 2004). Peptidases that cleave PARs to disarm the receptor and prevent subsequent activation by removing the tethered ligand have also been identified. For example, cathepsin G (EC 3.4.21.20) (Dulon *et al.*, 2003), pseudolysin (EC 3.4.24.26) (Dulon *et al.*, 2005) and myeloblastin (neutrophil leukocyte proteinase 3, EC 3.4.21.76) (Ramachandran *et al.*, 2011) cleave PAR₂ downstream of the tryptic cleavage site inactivating the receptor (Figure 2B). Cathepsin G and elastase-2 (neutrophil elastase, EC 3.4.21.37) abolish signalling by thrombin in PAR₃-transfected cells, and thus disarm PAR₃ (Cumashi *et al.*, 2001).

The concept of biased agonism at GPCRs is not a new one [reviewed in Urban *et al.* (2007)]. Given that PARs are activated by many different peptidases, it is not surprising that the generation of different N-termini can result in peptidase-specific signalling. It is these unique N-termini that act as intramolecular ligands causing specific GPCR conformations to elicit biased signalling responses. The earliest indications that PARs could be selectively activated came from observations comparing activation by synthetic ligands to peptidases. For PAR₁, mutations in the extracellular regions resulted in differential signalling between synthetic peptide agonists and thrombin (Blackhart *et al.*, 2000). For example, a PAR₁ deletion mutant lacking amino acids 68–93 of the N-terminus, failed to mobilize intracellular calcium in response to the synthetic peptide, but retained the ability to respond to thrombin (Blackhart *et al.*, 2000). A subsequent study using human endothelial cells showed that thrombin favours G_{12/13} signalling and induction of endothelial barrier permeability rather than intracellular calcium mobilization, whereas synthetic peptides preferentially caused intracellular calcium mobilization by triggering G_q signalling (McLaughlin *et al.*, 2005). Studies have also shown that activation of PAR₁ by different peptidases can have different functional consequences for endothelial barrier permeability (Feistritzer and

Riewald, 2005; Finigan *et al.*, 2005). Whereas thrombin-dependent cleavage of PAR₁ disrupts endothelial barrier integrity, activated protein C-dependent activation of PAR₁ enhances endothelial barrier function (Feistritzer and Riewald, 2005; Finigan *et al.*, 2005). Challenging human lung epithelial cells with elastase-2 leads to PAR₁-mediated apoptosis, similar to that observed with thrombin and a synthetic-activating peptide, but modified kinetics (Suzuki *et al.*, 2005; 2009). Matrix metalloproteinase-1 cleaves PAR₁ at a distinct site to thrombin and activates Rho-GTP and mitogenic pathways in a biased mechanism, to promote cell shape change and motility (Trivedi *et al.*, 2009).

Agonist-biased signalling has also been observed for PAR₂. Certain synthetic peptide agonists fail to cause PAR₂-mediated calcium signalling, while still activating ERK1/2, whereas SLIGRL-NH₂, which mimics the naturally unmasked tethered ligand, triggers both calcium and MAP kinase signalling (Ramachandran *et al.*, 2009). Further evidence came from the observation that a novel peptidomimetic PAR₂ antagonist, K-14585 activated ERK1/2 signalling, but failed to elicit calcium responses (Goh *et al.*, 2009). It was initially thought that elastase-2 also disarmed PAR₂ (Dulon *et al.*, 2003). However, recent evidence suggests that elastase-2 is in fact a biased agonist of PAR₂ (Ramachandran *et al.*, 2011) (Figure 2C). Although elastase-2 does not cause internalization of PAR₂ or mobilization of intracellular calcium, it does initiate activation of the ERK1/2 pathway (Ramachandran *et al.*, 2011). The exact mechanism by which peptidases can act as biased agonists has not yet been fully established. However, in the case of PAR₂ it is known that trypsin and elastase-2 cleave PAR₂ at different positions within the N-terminus. Therefore, it is highly probable that the different cleavages result in different conformations of PAR₂, which leads to biased activation of signalling pathways. With regard to PAR₁, although the peptidases may cleave at the same sites, the fact that thrombin actually binds to PAR₁ may lead to slight conformation changes, which also lead to biased activation of signalling cascades. However, until the crystal structures of the activated GPCRs are solved, this is pure conjecture. It is not yet known if biased agonism of PAR₃ and PAR₄ occurs. However, with numerous peptidases able to cleave members of this subfamily of GPCRs and perhaps each with different functional consequences, there is still much to discover about how peptidases signal through GPCRs.

Lysosomal peptidases degrade GPCRs to irrevocably terminate GPCR signalling

Once activation of a GPCR has occurred, many GPCRs are removed from the cell-surface to prevent uncontrolled signalling. The receptors are then either recycled back to the cell-surface mediating resensitization to allow cells to respond to the same stimulus again, or they are trafficked to intracellular compartments, such as lysosomes for degradation, which results in permanent signal arrest. The peptidases present in lysosomes mainly belong to the aspartic, cysteine and serine peptidase classes, with few metallo- and threonine peptidases residing in lysosomes. They are also mainly endopeptidases,

cleaving within polypeptide chains and not at the end or beginning of chains. Examples of peptidases localized to lysosomes include the serine peptidases, serine carboxypeptidase A (EC 3.4.16.5) and cathepsin G, aspartic peptidases, cathepsin D (EC 3.4.23.5) and E (EC 3.4.23.24) and cysteine peptidases, cathepsin B (EC 3.4.22.1) and L (EC 3.4.22.15). One mechanism by which many GPCRs are targeted for degradation is by the post-translational modification of ubiquitination, although this is not an absolute requirement for all GPCRs. Once activated, ubiquitin moieties are added to intracellular facing lysine residues by a process requiring three enzymes, an ubiquitin-activating enzyme (E1), an ubiquitin-conjugating enzyme (E2) and an ubiquitin ligase (E3). Before the GPCR is degraded by peptidases present in the lysosomes, the ubiquitin molecules are removed by metallo- or cysteine peptidases, referred to as de-ubiquitinating enzymes or ubiquitin-specific peptidases (USPs). In fact it is an absolute requirement that the ubiquitin molecules are removed before the GPCR can enter the lysosome. The first identified GPCR to be regulated by ubiquitination was the yeast GPCR, Ste2p. Ubiquitination of Ste2p is a requirement for internalization (Hicke and Riezman, 1996). Prolonged exposure of β_2 -adrenoceptors to agonists promotes sorting of receptors to lysosomes and degradation by lysosomal peptidases (Moore *et al.*, 1999). A combination of the aspartic peptidase inhibitor, pepstatin A and the serine peptidase inhibitor, leupeptin prevented degradation (Moore *et al.*, 1999). The β_2 -adrenoceptor was the first identified mammalian GPCR shown to be directed to lysosomes for degradation by ubiquitination (Shenoy *et al.*, 2001), an observation closely followed by that for the C-X-C chemokine receptor type 4 (CXCR₄) (Marchese and Benovic, 2001). PARs are sometimes termed one-shot receptors, because of the proteolytic nature of their activation. PAR₂ is also targeted for degradation by lysosomal peptidases by ubiquitination (Jacob *et al.*, 2005). N-CBZ-L-phenylalanyl-L-alanine-diazomethylketone, an inhibitor of lysosomal cysteine peptidases, cathepsins B and L prevents degradation of PAR₂. However, there are GPCRs that are trafficked to lysosomes without the need for modification by ubiquitin moieties. Although, agonists induce ubiquitination of the δ -opioid receptor, this ubiquitination is not an absolute requirement for lysosomal targeting, as a lysine-less mutant of the δ -opioid receptor is still efficiently trafficked to lysosomes and degraded by lysosomal serine peptidases (Tsao and von Zastrow, 2000; Tanowitz and Von Zastrow, 2002). However, further studies have indicated that ubiquitination does have a subtle role in the regulation of δ -opioid receptors. Ubiquitination, although not affecting trafficking to lysosomes does alter the rate at which the δ -opioid receptor is degraded in lysosomes (Hislop *et al.*, 2009). A similar role for ubiquitination is observed for the μ -opioid receptor. It is sequences in the C-terminal tail of the μ -opioid receptor that direct its trafficking to lysosomes, whereas ubiquitination of lysine residues in the first intracellular loop promote transfer of internalized μ -opioid receptors from the limiting endosome membrane to lumen thereby facilitating degradation (Hislop *et al.*, 2011). The receptor for calcitonin gene-related peptide (CGRP) is an unusual heterodimeric GPCR, comprising the GPCR, calcitonin receptor-like receptor (CLR) and a single transmembrane protein, receptor activity-modifying protein 1 (RAMP1) (McLatchie *et al.*, 1998). Following sus-

tained activation with CGRP, CLR•RAMP1 traffics to lysosomes and is degraded by lysosomal peptidases (Kuwasako *et al.*, 2000; Cottrell *et al.*, 2007). The identity of the lysosomal peptidases degrading CLR and RAMP1 is less clear, as an inhibitor cocktail blocking serine, aspartic and cysteine peptidases was used. However, CGRP does not induce ubiquitination of CLR or RAMP1 (Cottrell *et al.*, 2007). Although there are known roles for lysosomal peptidases in the immune system, in generating or destroying antigenic peptides and in trafficking of growth factor receptors [reviewed in Muller *et al.* (2012)], the functional consequences of preventing the degradation of GPCRs in lysosomes remain to be determined.

Proteolytic removal of ubiquitin regulates GPCR-dependent trafficking and signalling

The identity of many of the E3 ligases responsible for facilitating the addition of ubiquitin moieties to GPCRs are now known and are summarized elsewhere (Hislop and von Zastrow, 2011). However, unlike the E3 ligases, fewer USPs responsible for the de-ubiquitination of GPCRs have been identified. USPs play an important role in the regulation of GPCRs by ubiquitination, opposing the action of E3 ligases by removing ubiquitin molecules. This proteolysis is not only important for regulating GPCR signalling, but is also critical for maintaining the cellular pools of ubiquitin, which is critical for many other processes including regulation of transcription, cell-cycle control, DNA damage responses, apoptosis and the immune response [reviewed in Malynn and Ma (2010); Ramaekers and Wouters (2011); Vucic *et al.* (2011); Hammond-Martel *et al.* (2012); Starostina and Kipreos (2012)]. Studies have shown that under basal conditions PAR₁ is ubiquitinated at the cell-surface and that this ubiquitination promotes cell-surface retention (Wolfe *et al.*, 2007). Following proteolytic activation by thrombin, PAR₁ is de-ubiquitinated by unidentified USPs, a process that facilitates internalization to endosomes (Wolfe *et al.*, 2007). CXCR₇, a recycling GPCR is also ubiquitinated under basal conditions. After activation induced by the stromal-derived factor CXCL12, CXCR₇ is reversibly de-ubiquitinated by an unidentified USP, before trafficking back to the cell-surface (Canals *et al.*, 2012). Unusually, κ -opioid receptors are polyubiquitinated following agonist stimulation. These polyubiquitin chains are subsequently removed by CylD protein, a USP that specifically cleaves Lys-63 ubiquitin chains. After removal of the ubiquitin chains, κ -opioid receptors are degraded in lysosomes (Li *et al.*, 2008). However, other than down-regulation of the κ -opioid receptor, the functional consequences of this polyubiquitination have yet to be elucidated. β_2 -Adrenoceptors are regulated by USP33 and USP20, which act in a coordinated fashion to regulate the β_2 -adrenoceptor recycling and resensitization (Berthouze *et al.*, 2009). Similarly, ubiquitinated adenosine A_{2A} receptors are de-ubiquitinated by USP4, promoting trafficking back to the cell-surface (Milojevic *et al.*, 2006). In contrast, the proteolytic activities of associated molecule with the SH3 domain of signal-transducing adaptor molecule de-

ubiquitinating peptidase (AMSH) and USP8 are required to promote entry of PAR₂ to lysosomes (Hasdemir *et al.*, 2009). AMSH and USP8 also regulate the proteolytic down-regulation of the δ -opioid receptor, but in contrast to PAR₂, their activities do not affect the trafficking δ -opioid receptors to lysosomes (Hislop *et al.*, 2009). Expression of dominant negative mutants of AMSH and USP8 caused accumulation of PAR₂ in endosomes, but had no effect on endosomal mitogenic signalling, indicating that de-ubiquitination does not regulate association with β -arrestins (Hasdemir *et al.*, 2009). The effect of deubiquitination on the mitogenic signalling of other GPCRs has not been examined. However, ubiquitination of CXCR₄ by atrophin-interacting protein 4 is reported to enhance CXCR₄-mediated ERK1/2 activation (Malik *et al.*, 2012). Although only a few USPs that regulate ubiquitin dynamics for GPCRs have been identified to date, it is clear that de-ubiquitination by peptidases plays a different role for each GPCR, in regulating trafficking and signalling.

The cytosolic proteins, β -arrestin 1 and 2 are key regulators of GPCR signalling and internalization [reviewed in Moore *et al.* (2007); Shenoy and Lefkowitz (2011)] and may be ubiquitinated following agonist stimulation. The first observation of β -arrestin ubiquitination occurred following activation of the β_2 -adrenoceptor and is regulated by the E3 ligase, murine double minute 2 (Mdm2) (Shenoy *et al.*, 2001). A later study showed that de-ubiquitination of β -arrestin triggers its release from β_2 -adrenoceptors and the V₂ vasopressin receptors (Shenoy and Lefkowitz, 2003). GPCRs may be classed on their association with β -arrestins. Class A receptors bind only β -arrestin 2 and with low affinity, whereas class B receptors bind both β -arrestin 1 and 2 with similar affinities and for prolonged periods (Oakley *et al.*, 2000). When a β -arrestin-ubiquitin fusion protein was overexpressed in cells expressing the β_2 -adrenoceptor, the receptor associated with the β -arrestin-ubiquitin fusion protein for prolonged periods, changing the β_2 -adrenoceptor from a class A to a class B GPCR (Shenoy and Lefkowitz, 2003). Consistent with this observation and the fact that β -arrestins can mediate GPCR-induced ERK1/2 activation (Luttrell *et al.*, 1999), sustained ubiquitination of β -arrestins results in prolonged activation of ERK1/2 by agonists of β_2 -adrenoceptors and V₂ vasopressin receptors (Shenoy *et al.*, 2007) (Figure 3, panel B). The ubiquitination state of β -arrestins is reciprocally regulated by Mdm2 and USP33, with the actions of USP33 promoting the disassembly of endosomal signalling complexes terminating ERK signalling (Shenoy *et al.*, 2009).

There are additional ubiquitin-like modifying proteins, such as small ubiquitin-like modifier 1 (SUMO-1). Upon activation of the β_2 -adrenoceptor, β -arrestins are modified by the E2-modifying enzyme Ubc9, in a process termed sumoylation (Wyatt *et al.*, 2011). Although it is known that this sumoylation promotes the internalization of the β_2 -adrenoceptor, the identity of the SUMO-specific peptidase and the functional consequences of the desumoylation are not yet known.

Endosomal proteolysis regulates the recycling and resensitization of GPCRs

SP and CGRP are neuropeptides expressed by primary sensory neurons (Lundberg *et al.*, 1985). Noxious stimuli and trauma




Figure 3

USPs promote lysosomal degradation and recycling of GPCRs. (A) (1) USPs cleave ubiquitin molecules from GPCRs to (2) promote entry in lysosomes and degradation by lysosomal peptidases or (3) recycling of GPCRs to the cell-surface mediating resensitization of signalling. (B) Ubiquitination of β -arrestins promotes GPCR-mediated phosphorylation of extracellular-regulated PK 1 and 2. (2) De-ubiquitination of β -arrestins by USPs destabilizes the GPCR• β -arrestin complex to terminate ERK1/2 activation.

can lead to release of CGRP and SP from primary sensory neurons. The central release of CGRP and SP facilitates nociceptive transmission (Kuraishi *et al.*, 1988; Kawamura *et al.*, 1989; Pedersen-Bjergaard *et al.*, 1991), whereas peripheral release mediates neurogenic inflammation, which is characterized by neutrophil infiltration, oedema and vasodilatation (McDonald, 1988; McDonald *et al.*, 1988). CGRP, a potent vasodilator (Brain *et al.*, 1985) is also implicated in the pathogenesis of migraine [reviewed in Raddant and Russo, (2011); Moore and Salvatore (2012)].

Following activation, many GPCRs internalize with β -arrestins to early endosomes. In order for a GPCR to recycle back to the cell-surface, β -arrestins must be released. Until recently, the molecular mechanisms that initiate the release of β -arrestins from neuropeptide GPCRs, such as the NK₁ receptor and CLR•RAMP1 were poorly defined. NK₁ receptors and CLR•RAMP1 are both class B GPCRs, having sustained interactions with β -arrestins (Oakley *et al.*, 2000; Hilai

et al., 2001). Similar to the mechanisms that operate at the cell-surface, NK₁ receptors and CLR•RAMP1 are regulated by the membrane-anchored peptidase, ECE-1 present in endosomes (Figure 4) (Padilla *et al.*, 2007; Roosterman *et al.*, 2007). SP and CGRP are substrates for the endosomal peptidase, ECE-1, but only at endosomal pH (Johnson *et al.*, 1999; Padilla *et al.*, 2007; Roosterman *et al.*, 2007). As NK₁ receptors and CLR•RAMP1 are trafficked through the endosomal system, vacuolar type H⁺-ATPases pump protons inside the vesicles lowering the pH of endosomes (Forgac *et al.*, 1983; Cain *et al.*, 1989). Acidification has two effects: firstly, the affinity of SP and CGRP for their respective receptors is lowered, and secondly, SP and CGRP become substrates for ECE-1 (Padilla *et al.*, 2007; Roosterman *et al.*, 2007). ECE-1 cleaves SP and CGRP into inactive metabolites that can no longer interact with their GPCRs. β -Arrestins are then released and NK₁ receptors and CLR•RAMP1 are free to recycle back to the cell-surface (Padilla *et al.*, 2007; Roosterman *et al.*, 2007). Initial experiments were carried out using transfected cell lines, but further studies have shown that ECE-1-dependent cleavage of these neuropeptides regulates GPCR trafficking both in cells that endogenously express these receptors and in *in vivo* models. For example, ECE-1 regulates the resensitization of SP-induced plasma extravasation both in mice and rats, indicating the ECE-1 regulates NK₁ receptors in endothelial cells (Roosterman *et al.*, 2007; Cattaruzza *et al.*, 2009). ECE-1 also regulates the trafficking of NK₁ receptors in primary myenteric neurons (Pelayo *et al.*, 2011). More recently, we have shown that ECE-1 regulates resensitization of CGRP-induced cAMP generation in primary mesenteric artery smooth muscle cells (McNeish *et al.*, 2012). We also demonstrated that ECE-1 inhibition prevents the resensitization of CGRP-induced relaxation in rat mesenteric resistance arteries (McNeish *et al.*, 2012). This ECE-1-dependent regulation is not confined to NK₁ receptors and CLR•RAMP1, as other GPCRs are also regulated by this mechanism. Somatostatin-14 and -28 are inhibitory peptides exhibiting broad endocrine, exocrine and neuronal functions, such as the suppression of growth hormone secretion and the inhibition of pancreatic and gastrointestinal hormone release [reviewed in Olias *et al.* (2004)]. Somatostatin-14 and -28 exert their biological effects via activation of somatostatin receptors (sst receptors), which are expressed throughout the CNS and endocrine and immune systems. Sst receptors are also found at particularly high densities in many neuroendocrine tumours (Reubi *et al.*, 1987a; 1987b; 1987c). This high density of sst receptors allows imaging of tumours using a radiolabelled analogue of somatostatin called octreotide, a process termed sst scintigraphy (Lamberts *et al.*, 1990). The sst₂ receptor is regulated by ECE-1 following stimulation with somatostatin-14, but not by octreotide, reflecting the ability of ECE-1 to cleave somatostatin-14, but not octreotide (Roosterman *et al.*, 2008). A similar agonist-dependent trafficking was observed in studies with the corticotropin-releasing factor receptor 1 (CRF₁). CRF₁ receptors have two known agonists, corticotropin-releasing factor (CRF) and urocortin 1 (Ucn1). ECE-1 cleaves both peptides at endosomal pH, but only cleaves Ucn1 at a residue critical for receptor binding (Hasdemir *et al.*, 2012). At low agonist concentrations (30 nM), both Ucn1- and CRF-mediated intracellular calcium mobilization are dependent




Figure 4

Endosomal peptidases promote GPCR recycling and resensitization. (1) Vacuolar-type H⁺-ATPases pump protons (H⁺) into vesicles, acidifying early endosomes. (2) Peptide agonists such as SP and CGRP have reduced affinity for their respective GPCRs. SP and CGRP become substrates for the endosomal peptidase, ECE-1 at low pH and are hydrolysed to inactive metabolites. (3) β-Arrestins dissociate from the GPCR, returning to the cytosol. (4) The GPCR, free from β-arrestins then recycles back to the cell-surface to mediate resensitization. (5) Certain GPCRs (e.g. neurokinin-1 receptor) signal from endosomes in a β-arrestin-dependent mechanism, phosphorylating extracellular-regulated PKs 1 and 2 (pERK1/2). ECE-1 promoted dissociation of β-arrestins terminates ERK1/2 activation. (6) β-Arrestins can recruit protein phosphatases such as protein phosphatase 2A (PP2A) to desensitized GPCRs at the cell-surface. (7) PP2A activity dephosphorylates cell-surface located GPCRs promoting resensitization.

on ECE-1 activity; however, at high concentrations (100 nM), CRF-mediated intracellular calcium mobilization and CRF₁ receptor recycling and resensitization cease to be ECE-1-dependent. This loss of ECE-1-dependent trafficking perhaps reflects a mechanism to mediate distinct CRF₁ receptor trafficking and signalling, at higher concentrations of agonist (Hasdemir *et al.*, 2012). Neurotensin is also a substrate for ECE-1 at endosomal pH (Johnson *et al.*, 1999) and mediates intestinal inflammation and cell proliferation through activation of the neurotensin 1 receptor (NTS₁) (Castagliuolo *et al.*, 1999; Brun *et al.*, 2005). Endosomal ECE-1 activity promotes degradation of neurotensin and recycling of NTS₁ receptors (Law *et al.*, 2012).

Not all peptide-activated GPCRs are regulated by ECE-1. Studies have shown that although ECE-1 degrades bradykinin, ECE-1 does not regulate the recycling and resensitization of B₂ receptors (Padilla *et al.*, 2007). This is because of the nature of the interaction of B₂ receptors with β-arrestins, B₂ receptors only exhibit a transient interact with β-arrestins (Simaan *et al.*, 2005). Thus, it is not only whether ECE-1 cleaves the agonist at endosomal pH that determines if ECE-1 regulates the GPCR, but also the duration of the association of the GPCR with β-arrestins. Moreover, agonists must be substrates for ECE-1 at endosomal pH in order for the GPCR to be regulated by ECE-1. Although angiotensin I is a substrate for ECE-1, angiotensin II is not (Johnson *et al.*, 1999;

Padilla *et al.*, 2007) and ECE-1 does not regulate the recycling and resensitization of the AT₂ receptors (Padilla *et al.*, 2007).

An additional role for ECE-1-mediated cleavage of SP emerged following the observation that resensitization of SP-induced intracellular calcium mobilization precedes recycling of the NK₁ receptor (Bennett *et al.*, 2002; 2005; Murphy *et al.*, 2011). It is known that agonist-unoccupied GPCRs are desensitized by phosphorylation of serine and threonine residues by the second messenger kinases, PKA and PKC (Roth *et al.*, 1991; Pitcher *et al.*, 1992; Dery *et al.*, 2001). This second messenger kinase-dependent phosphorylation desensitizes the GPCR without causing internalization. In the case of the NK₁ receptor, it is the dephosphorylation of cell-surface located, desensitized receptors that mediates resensitization of SP-induced signalling (Figure 4) (Murphy *et al.*, 2011). ECE-1-dependent cleavage of SP releases β-arrestins from the endosomal GPCR complex. β-Arrestins then facilitate the recruitment of protein phosphatase 2A (PP2A), a known regulator of GPCRs (Pitcher *et al.*, 1995) to the cell-surface, where PP2A dephosphorylates NK₁ receptors to promote resensitization (Murphy *et al.*, 2011). It is not currently known if the same or a similar mechanism controls the resensitization of other ECE-1-regulated GPCRs. Further, it is not yet known whether other endosomal peptidases regulate the trafficking of other peptide-activated GPCRs. Peptidases may represent a therapeutic target, whereby inhibitors of peptidases would

prevent recycling and resensitization of GPCRs to prevent uncontrolled GPCR-mediated signalling that contributes to disease.

Proteolysis regulates endosomal signalling of GPCRs

In contrast to their role at the cell-surface in terminating G protein-dependent signalling, an additional function of β -arrestins is the recruitment of the signalling molecules to GPCRs in early endosomes. The signalling apparatus recruited to GPCRs by β -arrestins, serves to initiate a second wave of signalling that is distinct from that initiated at the cell-surface (Luttrell *et al.*, 1999). This endosome-based signalling is a relatively new area of investigation and has been recently reviewed (von Zastrow and Sorkin, 2007; Murphy *et al.*, 2009). For certain GPCRs, proteolysis of ligands by peptidases located in endosomes, serves as the molecular switch that determines the duration of this β -arrestin-dependent signalling. At the endosome surface, β -arrestins act as a scaffold to aid formation of mitogenic signalling complexes that include signalling proteins, such as Raf-1, MAPK kinases and ERK (Daaka *et al.*, 1998; DeFea *et al.*, 2000). The stability of these so-called signalosomes is dependent on the sustained interaction of β -arrestins with GPCRs in endosomes. Thus, for NK₁ receptors, ECE-1-dependent hydrolysis of SP regulates the stability of the NK₁ receptor• β -arrestin interaction and thus, the duration of SP-induced β -arrestin-dependent ERK activity (Cottrell *et al.*, 2009). This endosome-derived ERK1/2 activation up-regulates and phosphorylates the nuclear death receptor, Nur77 via a mechanism that requires β -arrestins, Raf-1, MAPK kinase 2 and ERK2 (Castro-Obregon *et al.*, 2004). Inhibition of ECE-1, which prevents the dissociation of NK₁ receptors and β -arrestins (Roosterman *et al.*, 2007), causes a sustained activation of ERK1/2, increased phosphorylation of Nur77, promoting cell death (Cottrell *et al.*, 2009). This mechanism also operates in cultured primary myenteric neurons and in intact animals (Cottrell *et al.*, 2009; Pelayo *et al.*, 2011). In mice, intraplantar injection of capsaicin, an activator of transient receptor potential vanilloid ion channel 1, promotes the release of SP in the dorsal horn to cause internalization of NK₁ receptors and activation of ERK1/2 (Mantyh *et al.*, 1995; Kawasaki *et al.*, 2004). An intrathecal injection of the highly selective ECE-1 inhibitor SM-19712 (Umekawa *et al.*, 2000) promoted a more sustained ERK1/2 activation following injection of capsaicin (Cottrell *et al.*, 2009). In contrast to SP-induced ERK1/2 activation, neurotensin-induced ERK1/2 activation is attenuated by ECE-1 inhibition, indicating that the recycling of NTS₁ receptors promotes ERK1/2 activation (Law *et al.*, 2012). ECE-1 inhibition similarly attenuated JNK activation, but promoted NF- κ B activation and IL-8 secretion (Law *et al.*, 2012).

Concluding remarks

It has long been known that peptidases present on the cell-surface regulate GPCR activation by generating or destroying

bioactive peptides, and that many GPCRs are ultimately degraded by peptidases present in lysosomes. However, it is now apparent that proteolysis also regulates other aspects of GPCRs, including trafficking through the endocytic system and signalling from endosomes. Intracellular proteolysis of peptidic agonists by peptidases present in endosomes, not only controls the recycling and resensitization of GPCRs, but also regulates the signalling from internalized GPCRs in endosomes. Ubiquitination of GPCRs targets certain GPCRs to lysosomes, but de-ubiquitination performed by USPs is required for efficient delivery of GPCRs to lysosomes, promoting GPCR down-regulation. For other GPCRs, USPs regulate the recycling to the cell-surface and thus, resensitization of signalling. Finally, USPs act to regulate the ubiquitination of β -arrestins and downstream mitogenic signalling cascades. ACE inhibitors, such as captopril have been successfully used to treat hypertension for many years and offer hope that other peptidase inhibitors could also be used to treat disease. Inhibitors of endosomal peptidases, such as ECE-1 could be used to regulate the trafficking of CGRP and SP receptors. ECE-1 inhibitors, by preventing the recycling of CGRP receptors, could prevent sustained CGRP signalling implicated in migraine. Inhibitors of ADAM peptidases, responsible for the generation of ligands causing transactivation of ErbB receptors, may represent a new therapy to prevent proliferation of cells and thereby attenuate cancer growth. The signalling pathways arising from internalized GPCRs are distinct from pathways initiated at the cell-surface and have unique cellular consequences. Inhibitors of USPs and endosomal peptidases have been shown to regulate this signalling and thus inhibitors of these peptidases may be useful tools to regulate the signalling of internalized GPCRs. Thus, the discovery of these new peptidase-driven mechanisms of GPCR regulation opens up a new line of potential peptidase inhibitor therapies to treat GPCR-mediated diseases. A key factor in the design of these peptidase inhibitors will be targeting them to the cellular compartments where they will have the required effects.

Acknowledgements

Supported by British Heart Foundation Fellowship to G.S.C (FS/08/017/25027).

Conflict of interest

The author declares no conflict of interest.

References

- Alexander SPH, Mathie A, Peters JA (2011). Guide to Receptors and Channels (GRAC), 5th edition. Br J Pharmacol 164 (Suppl. 1): S1–S324.
- Asakura M, Kitakaze M, Takashima S, Liao Y, Ishikura F, Yoshinaka T *et al.* (2002). Cardiac hypertrophy is inhibited by antagonism of ADAM12 processing of HB-EGF: metalloproteinase inhibitors as a new therapy. Nat Med 8: 35–40.

- Bennett VJ, Perrine SA, Simmons MA (2002). A novel mechanism of neurokinin-1 receptor resensitization. *J Pharmacol Exp Ther* 303: 1155–1162.
- Bennett VJ, Perrine SA, Simmons MA (2005). Neurokinin-1 receptor resensitization precedes receptor recycling. *J Pharmacol Exp Ther* 313: 1347–1354.
- Berthouze M, Venkataramanan V, Li Y, Shenoy SK (2009). The deubiquitinases USP33 and USP20 coordinate beta2 adrenergic receptor recycling and resensitization. *EMBO J* 28: 1684–1696.
- Blackhart BD, Ruslim-Litrus L, Lu CC, Alves VL, Teng W, Scarborough RM *et al.* (2000). Extracellular mutations of protease-activated receptor-1 result in differential activation by thrombin and thrombin receptor agonist peptide. *Mol Pharmacol* 58: 1178–1187.
- Bohm SK, Kong W, Bromme D, Smeekens SP, Anderson DC, Connolly A *et al.* (1996). Molecular cloning, expression and potential functions of the human proteinase-activated receptor-2. *Biochem J* 314 (Pt 3): 1009–1016.
- Boire A, Covic L, Agarwal A, Jacques S, Sherifi S, Kuliopulos A (2005). PAR1 is a matrix metalloprotease-1 receptor that promotes invasion and tumorigenesis of breast cancer cells. *Cell* 120: 303–313.
- Brain SD, Williams TJ, Tippins JR, Morris HR, MacIntyre I (1985). Calcitonin gene-related peptide is a potent vasodilator. *Nature* 313: 54–56.
- Bralet J, Marie C, Mossiat C, Lecomte JM, Gros C, Schwartz JC (1994). Effects of alatriopril, a mixed inhibitor of atriopeptidase and angiotensin I-converting enzyme, on cardiac hypertrophy and hormonal responses in rats with myocardial infarction. Comparison with captopril. *J Pharmacol Exp Ther* 270: 8–14.
- Brun P, Mastrotto C, Beggiao E, Stefani A, Barzon L, Sturniolo GC *et al.* (2005). Neuropeptide neurotensin stimulates intestinal wound healing following chronic intestinal inflammation. *Am J Physiol Gastrointest Liver Physiol* 288: G621–G629.
- Burger MM (1970). Proteolytic enzymes initiating cell division and escape from contact inhibition of growth. *Nature* 227: 170–171.
- Buteau J, Foisy S, Joly E, Prentki M (2003). Glucagon-like peptide 1 induces pancreatic beta-cell proliferation via transactivation of the epidermal growth factor receptor. *Diabetes* 52: 124–132.
- Cain CC, Sipe DM, Murphy RF (1989). Regulation of endocytic pH by the Na⁺,K⁺-ATPase in living cells. *Proc Natl Acad Sci U S A* 86: 544–548.
- Camerer E, Huang W, Coughlin SR (2000). Tissue factor- and factor X-dependent activation of protease-activated receptor 2 by factor VIIa. *Proc Natl Acad Sci U S A* 97: 5255–5260.
- Canals M, Scholten DJ, de Munnik S, Han MK, Smit MJ, Leurs R (2012). Ubiquitination of CXCR7 controls receptor trafficking. *PLoS ONE* 7: e34192.
- Carney DH, Cunningham DD (1977). Initiation of chick cell division by trypsin action at the cell surface. *Nature* 268: 602–606.
- Carney DH, Cunningham DD (1978). Transmembrane action of thrombin initiates chick cell division. *J Supramol Struct* 9: 337–350.
- Castagliuolo I, Wang CC, Valenick L, Pasha A, Nikulasson S, Carraway RE *et al.* (1999). Neurotensin is a proinflammatory neuropeptide in colonic inflammation. *J Clin Invest* 103: 843–849.
- Castro-Obregon S, Rao RV, del Rio G, Chen SF, Poksay KS, Rabizadeh S *et al.* (2004). Alternative, nonapoptotic programmed cell death: mediation by arrestin 2, ERK2, and Nur77. *J Biol Chem* 279: 17543–17553.
- Cattaruzza F, Cottrell GS, Vaksman N, Bunnett NW (2009). Endothelin-converting enzyme 1 promotes re-sensitization of neurokinin 1 receptor-dependent neurogenic inflammation. *Br J Pharmacol* 156: 730–739.
- Chen LB, Buchanan JM (1975). Mitogenic activity of blood components. I. Thrombin and prothrombin. *Proc Natl Acad Sci U S A* 72: 131–135.
- Cheng ZJ, Vapaatalo H, Mervaala E (2005). Angiotensin II and vascular inflammation. *Med Sci Monit* 11: RA194–RA205.
- CONSENSUS (1987). Effects of enalapril on mortality in severe congestive heart failure. Results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). The CONSENSUS Trial Study Group. *N Engl J Med* 316: 1429–1435.
- Cottrell GS, Amadesi S, Grady EF, Bunnett NW (2004). Trypsin IV, a novel agonist of protease-activated receptors 2 and 4. *J Biol Chem* 279: 13532–13539.
- Cottrell GS, Padilla B, Pikios S, Roosterman D, Steinhoff M, Grady EF *et al.* (2007). Post-endocytic sorting of calcitonin receptor-like receptor and receptor activity-modifying protein 1. *J Biol Chem* 282: 12260–12271.
- Cottrell GS, Padilla BE, Amadesi S, Poole DP, Murphy JE, Hardt M *et al.* (2009). Endosomal endothelin-converting enzyme-1: a regulator of beta-arrestin-dependent ERK signaling. *J Biol Chem* 284: 22411–22425.
- Cumashi A, Ansuini H, Celli N, De Blasi A, O'Brien PJ, Brass LF *et al.* (2001). Neutrophil proteases can inactivate human PAR3 and abolish the co-receptor function of PAR3 on murine platelets. *Thromb Haemost* 85: 533–538.
- Daaka Y, Luttrell LM, Ahn S, Della Rocca GJ, Ferguson SS, Caron MG *et al.* (1998). Essential role for G protein-coupled receptor endocytosis in the activation of mitogen-activated protein kinase. *J Biol Chem* 273: 685–688.
- Daub H, Weiss FU, Wallasch C, Ullrich A (1996). Role of transactivation of the EGF receptor in signalling by G-protein-coupled receptors. *Nature* 379: 557–560.
- Daull P, Benrezzak O, Arsenault D, Pheng LH, Blouin A, Cayer J *et al.* (2005). Triple vasopectidase inhibition normalizes blood pressure in conscious, unrestrained, and spontaneously hypertensive rats. *Am J Hypertens* 18 (12 Pt 1): 1606–1613.
- Daull P, Blouin A, Belleville K, Beaudoin M, Arsenault D, Leonard H *et al.* (2006a). Triple VPI CGS 35601 reduces high blood pressure in low-renin, high-salt Dahl salt-sensitive rats. *Exp Biol Med* (Maywood) 231: 830–833.
- Daull P, Lepage R, Benrezzak O, Cayer J, Beaudoin M, Belleville K *et al.* (2006b). The first preclinical pharmacotoxicological safety assessment of CGS 35601, a triple vasopectidase inhibitor, in chronically instrumented, conscious, and unrestrained spontaneously hypertensive rats. *Drug Chem Toxicol* 29: 183–202.
- DeFea KA, Zalevsky J, Thoma MS, Dery O, Mullins RD, Bunnett NW (2000). beta-arrestin-dependent endocytosis of proteinase-activated receptor 2 is required for intracellular targeting of activated ERK1/2. *J Cell Biol* 148: 1267–1281.
- Dery O, DeFea KA, Bunnett NW (2001). Protein kinase C-mediated desensitization of the neurokinin 1 receptor. *Am J Physiol Cell Physiol* 280: C1097–C1106.
- Dive V, Chang CF, Yiotakis A, Sturrock ED (2009). Inhibition of zinc metallopeptidases in cardiovascular disease – from unity to trinity, or duality? *Curr Pharm Des* 15: 3606–3621.
- Dulon S, Cande C, Bunnett NW, Hollenberg MD, Chignard M, Pidard D (2003). Proteinase-activated receptor-2 and human lung epithelial cells: disarming by neutrophil serine proteinases. *Am J Respir Cell Mol Biol* 28: 339–346.

- Dulon S, Leduc D, Cottrell GS, D'Alayer J, Hansen KK, Bunnett NW *et al.* (2005). *Pseudomonas aeruginosa* elastase disables proteinase-activated receptor 2 in respiratory epithelial cells. *Am J Respir Cell Mol Biol* 32: 411–419.
- Eguchi S, Inagami T (2000). Signal transduction of angiotensin II type 1 receptor through receptor tyrosine kinase. *Regul Pept* 91: 13–20.
- Eguchi S, Frank GD, Mifune M, Inagami T (2003). Metalloprotease-dependent ErbB ligand shedding in mediating EGFR transactivation and vascular remodelling. *Biochem Soc Trans* 31 (Pt 6): 1198–1202.
- Feistritzer C, Riewald M (2005). Endothelial barrier protection by activated protein C through PAR1-dependent sphingosine 1-phosphate receptor-1 crossactivation. *Blood* 105: 3178–3184.
- Finigan JH, Dudek SM, Singleton PA, Chiang ET, Jacobson JR, Camp SM *et al.* (2005). Activated protein C mediates novel lung endothelial barrier enhancement: role of sphingosine 1-phosphate receptor transactivation. *J Biol Chem* 280: 17286–17293.
- Forgac M, Cantley L, Wiedenmann B, Altstiel L, Branton D (1983). Clathrin-coated vesicles contain an ATP-dependent proton pump. *Proc Natl Acad Sci U S A* 80: 1300–1303.
- Goh FG, Ng PY, Nilsson M, Kanke T, Plevin R (2009). Dual effect of the novel peptide antagonist K-14585 on proteinase-activated receptor-2-mediated signalling. *Br J Pharmacol* 158: 1695–1704.
- Gooz P, Dang Y, Higashiyama S, Twal WO, Haycraft CJ, Gooz M (2012). A disintegrin and metalloenzyme (ADAM) 17 activation is regulated by $\alpha 5\beta 1$ integrin in kidney mesangial cells. *PLoS ONE* 7: e33350.
- Gros C, Noel N, Souque A, Schwartz JC, Danvy D, Plaquevent JC *et al.* (1991). Mixed inhibitors of angiotensin-converting enzyme (EC 3.4.15.1) and enkephalinase (EC 3.4.24.11): rational design, properties, and potential cardiovascular applications of glycopril and alatriopril. *Proc Natl Acad Sci U S A* 88: 4210–4214.
- Gross DM, Sweet CS, Ulm EH, Backlund EP, Morris AA, Weitz D *et al.* (1981). Effect of N-[(S)-1-carboxy-3-phenylpropyl]-L-Ala-L-Pro and its ethyl ester (MK-421) on angiotensin converting enzyme in vitro and angiotensin I pressor responses in vivo. *J Pharmacol Exp Ther* 216: 552–557.
- Gschwind A, Hart S, Fischer OM, Ullrich A (2003). TACE cleavage of proamphiregulin regulates GPCR-induced proliferation and motility of cancer cells. *EMBO J* 22: 2411–2421.
- Hammond-Martel I, Yu H, Affar B (2012). Roles of ubiquitin signaling in transcription regulation. *Cell Signal* 24: 410–421.
- Hart S, Fischer OM, Prenzel N, Zwick-Wallasch E, Schneider M, Hennighausen L *et al.* (2005). GPCR-induced migration of breast carcinoma cells depends on both EGFR signal transactivation and EGFR-independent pathways. *Biol Chem* 386: 845–855.
- Hasdemir B, Murphy JE, Cottrell GS, Bunnett NW (2009). Endosomal deubiquitinating enzymes control ubiquitination and down-regulation of protease-activated receptor 2. *J Biol Chem* 284: 28453–28466.
- Hasdemir B, Mahajan S, Bunnett NW, Liao M, Bhargava A (2012). Endothelin-converting enzyme-1 actions determine differential trafficking and signaling of corticotropin-releasing factor receptor 1 at high agonist concentrations. *Mol Endocrinol* 26: 681–695.
- Hicke L, Riezman H (1996). Ubiquitination of a yeast plasma membrane receptor signals its ligand-stimulated endocytosis. *Cell* 84: 277–287.
- Hilairt S, Belanger C, Bertrand J, Laperriere A, Foord SM, Bouvier M (2001). Agonist-promoted internalization of a ternary complex between calcitonin receptor-like receptor, receptor activity-modifying protein 1 (RAMP1), and beta-arrestin. *J Biol Chem* 276: 42182–42190.
- Hislop JN, von Zastrow M (2011). Role of ubiquitination in endocytic trafficking of G-protein-coupled receptors. *Traffic* 12: 137–148.
- Hislop JN, Henry AG, Marchese A, von Zastrow M (2009). Ubiquitination regulates proteolytic processing of G protein-coupled receptors after their sorting to lysosomes. *J Biol Chem* 284: 19361–19370.
- Hislop JN, Henry AG, von Zastrow M (2011). Ubiquitination in the first cytoplasmic loop of mu-opioid receptors reveals a hierarchical mechanism of lysosomal down-regulation. *J Biol Chem* 286: 40193–40204.
- Huovila AP, Turner AJ, Pelto-Huikko M, Karkkainen I, Ortiz RM (2005). Shedding light on ADAM metalloproteinases. *Trends Biochem Sci* 30: 413–422.
- Ikeo T, Yabana H, Kurosawa H, Kaburaki M, Kikkawa K, Narita H *et al.* (1992). Acute hemodynamic effects of the active metabolite of imidapril, (4S)-3-((2S)-2-[N-((1S)-1-carboxy-3-phenyl-propyl)amino]propionyl)-1-methyl-2-oxoimidazolidine-4-carboxylic acid, and enalaprilat in anesthetized dogs. *Arzneimittelforschung* 42: 1109–1114.
- Ishihara H, Connolly AJ, Zeng D, Kahn ML, Zheng YW, Timmons C *et al.* (1997). Protease-activated receptor 3 is a second thrombin receptor in humans. *Nature* 386: 502–506.
- Ishihara H, Zeng D, Connolly AJ, Tam C, Coughlin SR (1998). Antibodies to protease-activated receptor 3 inhibit activation of mouse platelets by thrombin. *Blood* 91: 4152–4157.
- Itoh Y, Joh T, Tanida S, Sasaki M, Kataoka H, Itoh K *et al.* (2005). IL-8 promotes cell proliferation and migration through metalloproteinase-cleavage proHB-EGF in human colon carcinoma cells. *Cytokine* 29: 275–282.
- Jacob C, Cottrell GS, Gehringer D, Schmidlin F, Grady EF, Bunnett NW (2005). c-Cbl mediates ubiquitination, degradation, and down-regulation of human protease-activated receptor 2. *J Biol Chem* 280: 16076–16087.
- Johnson GD, Stevenson T, Ahn K (1999). Hydrolysis of peptide hormones by endothelin-converting enzyme-1. A comparison with neprilysin. *J Biol Chem* 274: 4053–4058.
- Kahn ML, Zheng YW, Huang W, Bigornia V, Zeng D, Moff S *et al.* (1998). A dual thrombin receptor system for platelet activation. *Nature* 394: 690–694.
- Kawamura M, Kuraishi Y, Minami M, Satoh M (1989). Antinociceptive effect of intrathecally administered antiserum against calcitonin gene-related peptide on thermal and mechanical noxious stimuli in experimental hyperalgesic rats. *Brain Res* 497: 199–203.
- Kawasaki Y, Kohno T, Zhuang ZY, Brenner GJ, Wang H, Van Der Meer C *et al.* (2004). Ionotropic and metabotropic receptors, protein kinase A, protein kinase C, and Src contribute to C-fiber-induced ERK activation and cAMP response element-binding protein phosphorylation in dorsal horn neurons, leading to central sensitization. *J Neurosci* 24: 8310–8321.
- Khimji AK, Rockey DC (2010). Endothelin – biology and disease. *Cell Signal* 22: 1615–1625.
- Kuraishi Y, Nanayama T, Ohno H, Minami M, Satoh M (1988). Antinociception induced in rats by intrathecal administration of antiserum against calcitonin gene-related peptide. *Neurosci Lett* 92: 325–329.

- Kuwasako K, Shimekake Y, Masuda M, Nakahara K, Yoshida T, Kitaura M *et al.* (2000). Visualization of the calcitonin receptor-like receptor and its receptor activity-modifying proteins during internalization and recycling. *J Biol Chem* 275: 29602–29609.
- Lamberts SW, Bakker WH, Reubi JC, Krenning EP (1990). Somatostatin-receptor imaging in the localization of endocrine tumors. *N Engl J Med* 323: 1246–1249.
- Law IK, Murphy JE, Bakirtzi K, Bunnett NW, Pothoulakis C (2012). Neurotensin-induced proinflammatory signaling in human colonocytes is regulated by beta-arrestins and endothelin-converting enzyme-1-dependent endocytosis and resensitization of neurotensin receptor 1. *J Biol Chem* 287: 15066–15075.
- Lee FS, Chao MV (2001). Activation of Trk neurotrophin receptors in the absence of neurotrophins. *Proc Natl Acad Sci U S A* 98: 3555–3560.
- Li JG, Haines DS, Liu-Chen LY (2008). Agonist-promoted Lys63-linked polyubiquitination of the human kappa-opioid receptor is involved in receptor down-regulation. *Mol Pharmacol* 73: 1319–1330.
- Linseman DA, Benjamin CW, Jones DA (1995). Convergence of angiotensin II and platelet-derived growth factor receptor signaling cascades in vascular smooth muscle cells. *J Biol Chem* 270: 12563–12568.
- Lundberg JM, Franco-Cereceda A, Hua X, Hokfelt T, Fischer JA (1985). Co-existence of substance P and calcitonin gene-related peptide-like immunoreactivities in sensory nerves in relation to cardiovascular and bronchoconstrictor effects of capsaicin. *Eur J Pharmacol* 108: 315–319.
- Luttrell LM, Ferguson SS, Daaka Y, Miller WE, Maudsley S, Della Rocca GJ *et al.* (1999). Beta-arrestin-dependent formation of beta2 adrenergic receptor-Src protein kinase complexes. *Science* 283: 655–661.
- McDonald DM (1988). Neurogenic inflammation in the rat trachea. I. Changes in venules, leucocytes and epithelial cells. *J Neurocytol* 17: 583–603.
- McDonald DM, Mitchell RA, Gabella G, Haskell A (1988). Neurogenic inflammation in the rat trachea. II. Identity and distribution of nerves mediating the increase in vascular permeability. *J Neurocytol* 17: 605–628.
- McLatchie LM, Fraser NJ, Main MJ, Wise A, Brown J, Thompson N *et al.* (1998). RAMPs regulate the transport and ligand specificity of the calcitonin-receptor-like receptor. *Nature* 393: 333–339.
- McLaughlin JN, Shen L, Holinstat M, Brooks JD, Dibenedetto E, Hamm HE (2005). Functional selectivity of G protein signaling by agonist peptides and thrombin for the protease-activated receptor-1. *J Biol Chem* 280: 25048–25059.
- McNeish AJ, Roux BT, Aylett S-B, Maassen Van Den Brink A, Cottrell GS (2012). Endosomal proteolysis regulates calcitonin gene-related peptide responses in mesenteric arteries. *Br J Pharmacol* 167: 1679–1690.
- Malik R, Soh UJ, Trejo J, Marchese A (2012). Novel roles for the E3 ubiquitin ligase atrophin-interacting protein 4 and signal transduction adaptor molecule 1 in G protein-coupled receptor signaling. *J Biol Chem* 287: 9013–9027.
- Malynn BA, Ma A (2010). Ubiquitin makes its mark on immune regulation. *Immunity* 33: 843–852.
- Mannaioni G, Orr AG, Hamill CE, Yuan H, Pedone KH, McCoy KL *et al.* (2008). Plasmin potentiates synaptic N-methyl-D-aspartate receptor function in hippocampal neurons through activation of protease-activated receptor-1. *J Biol Chem* 283: 20600–20611.
- Mantyh PW, DeMaster E, Malhotra A, Ghilardi JR, Rogers SD, Mantyh CR *et al.* (1995). Receptor endocytosis and dendrite reshaping in spinal neurons after somatosensory stimulation. *Science* 268: 1629–1632.
- Marchese A, Benovic JL (2001). Agonist-promoted ubiquitination of the G protein-coupled receptor CXCR4 mediates lysosomal sorting. *J Biol Chem* 276: 45509–45512.
- Matsas R, Kenny AJ, Turner AJ (1984). The metabolism of neuropeptides. The hydrolysis of peptides, including enkephalins, tachykinins and their analogues, by endopeptidase-24.11. *Biochem J* 223: 433–440.
- Maudsley S, Pierce KL, Zamah AM, Miller WE, Ahn S, Daaka Y *et al.* (2000). The beta(2)-adrenergic receptor mediates extracellular signal-regulated kinase activation via assembly of a multi-receptor complex with the epidermal growth factor receptor. *J Biol Chem* 275: 9572–9580.
- Mifune M, Ohtsu H, Suzuki H, Nakashima H, Brailoiu E, Dun NJ *et al.* (2005). G protein coupling and second messenger generation are indispensable for metalloprotease-dependent, heparin-binding epidermal growth factor shedding through angiotensin II type-1 receptor. *J Biol Chem* 280: 26592–26599.
- Milojevic T, Reiterer V, Stefan E, Korkhov VM, Dorostkar MM, Ducza E *et al.* (2006). The ubiquitin-specific protease Usp4 regulates the cell surface level of the A2A receptor. *Mol Pharmacol* 69: 1083–1094.
- Mogi M, Iwai M, Horiuchi M (2009). New insights into the regulation of angiotensin receptors. *Curr Opin Nephrol Hypertens* 18: 138–143.
- Molino M, Barnathan ES, Numerof R, Clark J, Dreyer M, Cumashi A *et al.* (1997). Interactions of mast cell tryptase with thrombin receptors and PAR-2. *J Biol Chem* 272: 4043–4049.
- Moore CA, Milano SK, Benovic JL (2007). Regulation of receptor trafficking by GRKs and arrestins. *Annu Rev Physiol* 69: 451–482.
- Moore EL, Salvatore CA (2012). Targeting a family B GPCR/RAMP receptor complex: CGRP receptor antagonists and migraine. *Br J Pharmacol* 166: 66–78.
- Moore RH, Tuffaha A, Millman EE, Dai W, Hall HS, Dickey BF *et al.* (1999). Agonist-induced sorting of human beta2-adrenergic receptors to lysosomes during downregulation. *J Cell Sci* 112 (Pt 3): 329–338.
- Muller S, Dennemarker J, Reinheckel T (2012). Specific functions of lysosomal proteases in endocytic and autophagic pathways. *Biochim Biophys Acta* 1824: 34–43.
- Murphy JE, Padilla BE, Hasdemir B, Cottrell GS, Bunnett NW (2009). Endosomes: a legitimate platform for the signaling train. *Proc Natl Acad Sci U S A* 106: 17615–17622.
- Murphy JE, Roosterman D, Cottrell GS, Padilla BE, Feld M, Brand E *et al.* (2011). Protein phosphatase 2A mediates resensitization of the neurokinin 1 receptor. *Am J Physiol Cell Physiol* 301: C780–C791.
- Myers TJ, Brennaman LH, Stevenson M, Higashiyama S, Russell WE, Lee DC *et al.* (2009). Mitochondrial reactive oxygen species mediate GPCR-induced TACE/ADAM17-dependent transforming growth factor-alpha shedding. *Mol Biol Cell* 20: 5236–5249.
- Neal B, MacMahon S, Ohkubo T, Brnabic A, Tonkin A (2002). Effects of the vasopeptidase inhibitor, omapatrilat, in 723 patients with coronary heart disease. *J Renin Angiotensin Aldosterone Syst* 3: 270–276.
- Nystedt S, Emilsson K, Wahlestedt C, Sundelin J (1994). Molecular cloning of a potential proteinase activated receptor. *Proc Natl Acad Sci U S A* 91: 9208–9212.

- Nystedt S, Emilsson K, Larsson AK, Strombeck B, Sundelin J (1995a). Molecular cloning and functional expression of the gene encoding the human proteinase-activated receptor 2. *Eur J Biochem* 232: 84–89.
- Nystedt S, Larsson AK, Aberg H, Sundelin J (1995b). The mouse proteinase-activated receptor-2 cDNA and gene. Molecular cloning and functional expression. *J Biol Chem* 270: 5950–5955.
- Oakley RH, Laporte SA, Holt JA, Caron MG, Barak LS (2000). Differential affinities of visual arrestin, beta arrestin1, and beta arrestin2 for G protein-coupled receptors delineate two major classes of receptors. *J Biol Chem* 275: 17201–17210.
- Ohtsu H, Dempsey PJ, Eguchi S (2006). ADAMs as mediators of EGF receptor transactivation by G protein-coupled receptors. *Am J Physiol Cell Physiol* 291: C1–10.
- Olias G, Viollet C, Kusserow H, Epelbaum J, Meyerhof W (2004). Regulation and function of somatostatin receptors. *J Neurochem* 89: 1057–1091.
- Ondetti MA, Rubin B, Cushman DW (1977). Design of specific inhibitors of angiotensin-converting enzyme: new class of orally active antihypertensive agents. *Science* 196: 441–444.
- Padilla BE, Cottrell GS, Roosterman D, Pikios S, Muller L, Steinhoff M *et al.* (2007). Endothelin-converting enzyme-1 regulates endosomal sorting of calcitonin receptor-like receptor and beta-arrestins. *J Cell Biol* 179: 981–997.
- Paolillo M, Schinelli S (2008). Therapeutic targeting of g-protein coupled receptor-mediated epidermal growth factor receptor transactivation in human glioma brain tumors. *Mini Rev Med Chem* 8: 1418–1428.
- Pedersen-Bjergaard U, Nielsen LB, Jensen K, Edvinsson L, Jansen I, Olesen J (1991). Calcitonin gene-related peptide, neurokinin A and substance P: effects on nociception and neurogenic inflammation in human skin and temporal muscle. *Peptides* 12: 333–337.
- Pelayo JC, Poole DP, Steinhoff M, Cottrell GS, Bunnett NW (2011). Endothelin-converting enzyme-1 regulates trafficking and signalling of the neurokinin 1 receptor in endosomes of myenteric neurones. *J Physiol* 589 (Pt 21): 5213–5230.
- Pierce KL, Tohgo A, Ahn S, Field ME, Luttrell LM, Lefkowitz RJ (2001). Epidermal growth factor (EGF) receptor-dependent ERK activation by G protein-coupled receptors: a co-culture system for identifying intermediates upstream and downstream of heparin-binding EGF shedding. *J Biol Chem* 276: 23155–23160.
- Pietri M, Schneider B, Mouillet-Richard S, Ermonval M, Mutel V, Launay JM *et al.* (2005). Reactive oxygen species-dependent TNF-alpha converting enzyme activation through stimulation of 5-HT2B and alpha1D autoreceptors in neuronal cells. *FASEB J* 19: 1078–1087.
- Pitcher J, Lohse MJ, Codina J, Caron MG, Lefkowitz RJ (1992). Desensitization of the isolated beta 2-adrenergic receptor by beta-adrenergic receptor kinase, cAMP-dependent protein kinase, and protein kinase C occurs via distinct molecular mechanisms. *Biochemistry* 31: 3193–3197.
- Pitcher JA, Payne ES, Csontos C, DePaoli-Roach AA, Lefkowitz RJ (1995). The G-protein-coupled receptor phosphatase: a protein phosphatase type 2A with a distinct subcellular distribution and substrate specificity. *Proc Natl Acad Sci U S A* 92: 8343–8347.
- Porrello ER, Delbridge LM, Thomas WG (2009). The angiotensin II type 2 (AT2) receptor: an enigmatic seven transmembrane receptor. *Front Biosci* 14: 958–972.
- Prenzel N, Zwick E, Daub H, Leserer M, Abraham R, Wallasch C *et al.* (1999). EGF receptor transactivation by G-protein-coupled receptors requires metalloproteinase cleavage of proHB-EGF. *Nature* 402: 884–888.
- Quinton TM, Kim S, Derian CK, Jin J, Kunapuli SP (2004). Plasmin-mediated activation of platelets occurs by cleavage of protease-activated receptor 4. *J Biol Chem* 279: 18434–18439.
- Raddant AC, Russo AF (2011). Calcitonin gene-related peptide in migraine: intersection of peripheral inflammation and central modulation. *Expert Rev Mol Med* 13: e36.
- Ramachandran R, Mihara K, Mathur M, Rochdi MD, Bouvier M, Defea K *et al.* (2009). Agonist-biased signaling via proteinase activated receptor-2: differential activation of calcium and mitogen-activated protein kinase pathways. *Mol Pharmacol* 76: 791–801.
- Ramachandran R, Mihara K, Chung H, Renaux B, Lau CS, Muruve DA *et al.* (2011). Neutrophil elastase acts as a biased agonist for proteinase-activated receptor-2 (PAR2). *J Biol Chem* 286: 24638–24648.
- Ramaekers CH, Wouters BG (2011). Regulatory functions of ubiquitin in diverse DNA damage responses. *Curr Mol Med* 11: 152–169.
- Rao GN, Delafontaine P, Runge MS (1995). Thrombin stimulates phosphorylation of insulin-like growth factor-1 receptor, insulin receptor substrate-1, and phospholipase C-gamma 1 in rat aortic smooth muscle cells. *J Biol Chem* 270: 27871–27875.
- Rawlings ND, Barrett AJ, Bateman A (2012). MEROPS: the database of proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Res* 40 (Database issue): D343–D350.
- Regamey F, Maillard M, Nussberger J, Brunner HR, Burnier M (2002). Renal hemodynamic and natriuretic effects of concomitant Angiotensin-converting enzyme and neutral endopeptidase inhibition in men. *Hypertension* 40: 266–272.
- Reubi JC, Hacki WH, Lamberts SW (1987a). Hormone-producing gastrointestinal tumors contain a high density of somatostatin receptors. *J Clin Endocrinol Metab* 65: 1127–1134.
- Reubi JC, Lang W, Maurer R, Koper JW, Lamberts SW (1987b). Distribution and biochemical characterization of somatostatin receptors in tumors of the human central nervous system. *Cancer Res* 47: 5758–5764.
- Reubi JC, Maurer R, von Werder K, Torhorst J, Klijn JG, Lamberts SW (1987c). Somatostatin receptors in human endocrine tumors. *Cancer Res* 47: 551–558.
- Riewald M, Petrovan RJ, Donner A, Mueller BM, Ruf W (2002). Activation of endothelial cell protease activated receptor 1 by the protein C pathway. *Science* 296: 1880–1882.
- Roosterman D, Cottrell GS, Padilla BE, Muller L, Eckman CB, Bunnett NW *et al.* (2007). Endothelin-converting enzyme 1 degrades neuropeptides in endosomes to control receptor recycling. *Proc Natl Acad Sci U S A* 104: 11838–11843.
- Roosterman D, Kempkes C, Cottrell GS, Padilla BE, Bunnett NW, Turck CW *et al.* (2008). Endothelin-converting enzyme-1 degrades internalized somatostatin-14. *Endocrinology* 149: 2200–2207.
- Roth NS, Campbell PT, Caron MG, Lefkowitz RJ, Lohse MJ (1991). Comparative rates of desensitization of beta-adrenergic receptors by the beta-adrenergic receptor kinase and the cyclic AMP-dependent protein kinase. *Proc Natl Acad Sci U S A* 88: 6201–6204.
- Rothmeier AS, Ruf W (2012). Protease-activated receptor 2 signaling in inflammation. *Semin Immunopathol* 34: 133–149.
- Sambrano GR, Huang W, Faruqi T, Mahrus S, Craik C, Coughlin SR (2000). Cathepsin G activates protease-activated receptor-4 in human platelets. *J Biol Chem* 275: 6819–6823.
- Sambrano GR, Weiss EJ, Zheng YW, Huang W, Coughlin SR (2001). Role of thrombin signalling in platelets in haemostasis and thrombosis. *Nature* 413: 74–78.

- Scase TJ, Heath MF, Evans RJ (1997). Cloning of PAR3 cDNA from human platelets, and human erythroleukemic and human promonocytic cell lines. *Blood* 90: 2113–2114.
- Schafer B, Marg B, Gschwind A, Ullrich A (2004). Distinct ADAM metalloproteinases regulate G protein-coupled receptor-induced cell proliferation and survival. *J Biol Chem* 279: 47929–47938.
- Sefton BM, Rubin H (1970). Release from density dependent growth inhibition by proteolytic enzymes. *Nature* 227: 843–845.
- Shenoy SK, Lefkowitz RJ (2003). Trafficking patterns of beta-arrestin and G protein-coupled receptors determined by the kinetics of beta-arrestin deubiquitination. *J Biol Chem* 278: 14498–14506.
- Shenoy SK, Lefkowitz RJ (2011). beta-Arrestin-mediated receptor trafficking and signal transduction. *Trends Pharmacol Sci* 32: 521–533.
- Shenoy SK, McDonald PH, Kohout TA, Lefkowitz RJ (2001). Regulation of receptor fate by ubiquitination of activated beta 2-adrenergic receptor and beta-arrestin. *Science* 294: 1307–1313.
- Shenoy SK, Barak LS, Xiao K, Ahn S, Berthouze M, Shukla AK *et al.* (2007). Ubiquitination of beta-arrestin links seven-transmembrane receptor endocytosis and ERK activation. *J Biol Chem* 282: 29549–29562.
- Shenoy SK, Modi AS, Shukla AK, Xiao K, Berthouze M, Ahn S *et al.* (2009). Beta-arrestin-dependent signaling and trafficking of 7-transmembrane receptors is reciprocally regulated by the deubiquitinase USP33 and the E3 ligase Mdm2. *Proc Natl Acad Sci U S A* 106: 6650–6655.
- Showkathali R, Natarajan A (2012). Antiplatelet and antithrombin strategies in acute coronary syndrome: state-of-the-art review. *Curr Cardiol Rev* PMID: 22935021 [Epub ahead of print].
- Simaan M, Bedard-Goulet S, Fessart D, Gratton JP, Laporte SA (2005). Dissociation of beta-arrestin from internalized bradykinin B2 receptor is necessary for receptor recycling and resensitization. *Cell Signal* 17: 1074–1083.
- SOLVD (1991). Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. The SOLVD Investigators. *N Engl J Med* 325: 293–302.
- Starostina NG, Kipreos ET (2012). Multiple degradation pathways regulate versatile CIP/KIP CDK inhibitors. *Trends Cell Biol* 22: 33–41.
- Suzuki T, Moraes TJ, Vachon E, Ginzberg HH, Huang TT, Matthay MA *et al.* (2005). Proteinase-activated receptor-1 mediates elastase-induced apoptosis of human lung epithelial cells. *Am J Respir Cell Mol Biol* 33: 231–247.
- Suzuki T, Yamashita C, Zemans RL, Briones N, Van Linden A, Downey GP (2009). Leukocyte elastase induces lung epithelial apoptosis via a PAR-1-, NF-kappaB-, and p53-dependent pathway. *Am J Respir Cell Mol Biol* 41: 742–755.
- Takeuchi T, Harris JL, Huang W, Yan KW, Coughlin SR, Craik CS (2000). Cellular localization of membrane-type serine protease 1 and identification of protease-activated receptor-2 and single-chain urokinase-type plasminogen activator as substrates. *J Biol Chem* 275: 26333–26342.
- Tanida S, Kataoka H, Mizoshita T, Shimura T, Kamiya T, Joh T (2010). Intracellular translocation signaling of HB-EGF carboxy-terminal fragment and mucosal defense through cell proliferation and migration in digestive tracts. *Digestion* 82: 145–149.
- Tanowitz M, Von Zastrow M (2002). Ubiquitination-independent trafficking of G protein-coupled receptors to lysosomes. *J Biol Chem* 277: 50219–50222.
- Thanawala V, Kadam VJ, Ghosh R (2008). Enkephalinase inhibitors: potential agents for the management of pain. *Curr Drug Targets* 9: 887–894.
- Trapani AJ, Beil ME, Bruseo CW, Savage P, Firooznia F, Jeng AY (2004). CGS 35601 and its orally active prodrug CGS 37808 as triple inhibitors of endothelin-converting enzyme-1, neutral endopeptidase 24.11, and angiotensin-converting enzyme. *J Cardiovasc Pharmacol* 44 (Suppl. 1): S211–S215.
- Trivedi V, Boire A, Tchernychev B, Kaneider NC, Leger AJ, O'Callaghan K *et al.* (2009). Platelet matrix metalloproteinase-1 mediates thrombogenesis by activating PAR1 at a cryptic ligand site. *Cell* 137: 332–343.
- Tsao PI, von Zastrow M (2000). Type-specific sorting of G protein-coupled receptors after endocytosis. *J Biol Chem* 275: 11130–11140.
- Umekawa K, Hasegawa H, Tsutsumi Y, Sato K, Matsumura Y, Ohashi N (2000). Pharmacological characterization of a novel sulfonylureid-pyrazole derivative, SM-19712, a potent nonpeptidic inhibitor of endothelin converting enzyme. *Jpn J Pharmacol* 84: 7–15.
- Urban JD, Clarke WP, von Zastrow M, Nichols DE, Kobilka B, Weinstein H *et al.* (2007). Functional selectivity and classical concepts of quantitative pharmacology. *J Pharmacol Exp Ther* 320: 1–13.
- Vu TK, Hung DT, Wheaton VI, Coughlin SR (1991). Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* 64: 1057–1068.
- Vucic D, Dixit VM, Wertz IE (2011). Ubiquitylation in apoptosis: a post-translational modification at the edge of life and death. *Nat Rev Mol Cell Biol* 12: 439–452.
- Wolfe BL, Marchese A, Trejo J (2007). Ubiquitination differentially regulates clathrin-dependent internalization of protease-activated receptor-1. *J Cell Biol* 177: 905–916.
- Wyatt D, Malik R, Vesecky AC, Marchese A (2011). Small ubiquitin-like modifier modification of arrestin-3 regulates receptor trafficking. *J Biol Chem* 286: 3884–3893.
- Xu WF, Andersen H, Whitmore TE, Presnell SR, Yee DP, Ching A *et al.* (1998). Cloning and characterization of human protease-activated receptor 4. *Proc Natl Acad Sci U S A* 95: 6642–6646.
- Yan Y, Shirakabe K, Werb Z (2002). The metalloprotease Kuzbanian (ADAM10) mediates the transactivation of EGF receptor by G protein-coupled receptors. *J Cell Biol* 158: 221–226.
- Yoshida N, Yoshikawa T (2008). Basic and translational research on proteinase-activated receptors: implication of proteinase/proteinase-activated receptor in gastrointestinal inflammation. *J Pharmacol Sci* 108: 415–421.
- Yusuf S, Sleight P, Pogue J, Bosch J, Davies R, Dagenais G (2000). Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 342: 145–153.
- von Zastrow M, Sorkin A (2007). Signaling on the endocytic pathway. *Curr Opin Cell Biol* 19: 436–445.
- Zhang Z, Oliver P, Lancaster JR Jr, Schwarzenberger PO, Joshi MS, Cork J *et al.* (2001). Reactive oxygen species mediate tumor necrosis factor alpha-converting, enzyme-dependent ectodomain shedding induced by phorbol myristate acetate. *FASEB J* 15: 303–305.
- Zwick E, Wallasch C, Daub H, Ullrich A (1999). Distinct calcium-dependent pathways of epidermal growth factor receptor transactivation and PYK2 tyrosine phosphorylation in PC12 cells. *J Biol Chem* 274: 20989–20996.