

RESEARCH PAPER

Potent vasorelaxant activity of the TMEM16A inhibitor T16A_{inh}-A01

Alison J Davis^{1*}, Jian Shi^{1*}, Harry AT Pritchard^{1*}, Preet S Chadha¹, Normand Leblanc², Georgios Vasilikostas³, Zhen Yao⁴, AS Verkman⁴, Anthony P Albert¹ and Iain A Greenwood¹

¹Pharmacology and Cell Physiology Research Group, Division of Biomedical Sciences, St George's, University of London, London, UK, ²Department of Pharmacology, University of Nevada, Reno, NV, USA, ³Department of Surgery, St George's Hospital, London, UK, and ⁴Departments of Medicine and Physiology, University of California, San Francisco, CA, USA

Correspondence

Iain Greenwood, Pharmacology and Cell Physiology Research Group, Division of Biomedical Sciences, St George's, University of London, London SW17 0RE, UK. E-mail: i.greenwood@sgul.ac.uk

*The authors Alison J Davis, Jian Shi and Harry AT Pritchard contributed equally.

Keywords

calcium-activated chloride channels; TMEM16A; anoctamins; Ano1; vascular smooth muscle; anti-hypertensive; patch clamp

Received

1 February 2012

Revised

25 July 2012

Accepted

15 August 2012

BACKGROUND AND PURPOSE

T16A_{inh}-A01 is a recently identified inhibitor of the calcium-activated chloride channel TMEM16A. The aim of this study was to test the efficacy of T16A_{inh}-A01 for inhibition of calcium-activated chloride channels in vascular smooth muscle and consequent effects on vascular tone.

EXPERIMENTAL APPROACH

Single channel and whole cell patch clamp was performed on single smooth muscle cells from rabbit pulmonary artery and mouse thoracic aorta. Isometric tension studies were performed on mouse thoracic aorta and mesenteric artery as well as human abdominal visceral adipose artery.

KEY RESULTS

In rabbit pulmonary artery myocytes T16A_{inh}-A01 (1–30 µM) inhibited single calcium (Ca²⁺)-activated chloride (Cl⁻) channels and whole cell currents activated by 500 nM free Ca²⁺. Similar effects were observed for single Ca²⁺-activated Cl⁻ channels in mouse thoracic aorta, and in both cell types, channel activity was abolished by two antisera raised against TMEM16A but not by a bestrophin antibody. The TMEM16A potentiator, F_{act} (10 µM), increased single channel and whole cell Ca²⁺-activated Cl⁻ currents in rabbit pulmonary arteries. In isometric tension studies, T16A_{inh}-A01 relaxed mouse thoracic aorta pre-contracted with methoxamine with an IC₅₀ of 1.6 µM and suppressed the methoxamine concentration–effect curve. T16A_{inh}-A01 did not affect the maximal contraction produced by 60 mM KCl and the relaxant effect of 10 µM T16A_{inh}-A01 was not altered by incubation of mouse thoracic aorta in a cocktail of potassium (K⁺) channel blockers. T16A_{inh}-A01 (10 µM) also relaxed human visceral adipose arteries by 88 ± 3%.

CONCLUSIONS AND IMPLICATIONS

T16A_{inh}-A01 blocks calcium-activated chloride channels in vascular smooth muscle cells and relaxes murine and human blood vessels.

Abbreviations

CACCs, calcium-activated chloride currents; I_{ClCa}, calcium-activated chloride currents; MA, mesenteric artery; PA, pulmonary artery; TA, thoracic aorta; VDCCs, voltage-dependent calcium channels

Introduction

Calcium (Ca^{2+})-activated chloride (Cl^-) currents (I_{ClCa}) have been recorded from a number of different cell types, including epithelial cells, secretory glands, cardiomyocytes, neurones and smooth muscle cells. Research into the physiological and pathophysiological roles of the underlying Ca^{2+} -activated Cl^- channels (CACCs) has been hampered by the lack of molecular identity and the poor selectivity of existing Cl^- channel blockers such as niflumic acid (see Greenwood and Leblanc, 2007). However, the identification that expression products of the gene *TMEM16A* (Ano1) generate a CACC with biophysical properties identical to native CACCs (Caputo *et al.*, 2008; Schroeder *et al.*, 2008; Yang *et al.*, 2008) has allowed the development through small-molecule screening of a new generation of reagents for investigating CACC (Namkung *et al.*, 2010; 2011a). These efforts led to the identification of T16A_{inh}-A01, a compound shown to inhibit CACC activity in TMEM16A-transfected FRT cells with an IC_{50} of $\sim 1 \mu\text{M}$ (Namkung *et al.*, 2011a). In addition, Namkung *et al.* (2011b) identified two agents that either activate TMEM16A independent of Ca^{2+} (E_{act}) or potentiate activated TMEM16A channels (F_{act}).

In vascular smooth muscle cells, Cl^- ions are accumulated so that opening of CACCs leads to membrane depolarization sufficient to increase the open probability of voltage-dependent Ca^{2+} channels (VDCCs; Large and Wang, 1996; Kitamura and Yamazaki, 2001; Leblanc *et al.*, 2005). Consequently, CACCs are generally considered as a mechanism for vasoconstrictors to promote membrane contraction, but the lack of reliable pharmacological tools has hampered the research on this aspect. However, TMEM16A expression was recently revealed in vascular smooth muscle that exhibits robust I_{ClCa} (Davis *et al.*, 2010; Manoury *et al.*, 2010; Sones *et al.*, 2010; Thomas-Gatewood *et al.*, 2011). Moreover, TMEM16A down-regulation of TMEM16A transcripts by interference RNA strategies inhibited I_{ClCa} in pulmonary artery (PA) smooth muscle cells in short-term culture (Manoury *et al.*, 2010). However, there is no information on the contribution of TMEM16A to CACCs in freshly dispersed smooth muscle cells or whether TMEM16A is involved in vascular contractions. The aims of the present study were twofold. Firstly, to investigate the effect of T16A_{inh}-A01 and F_{act} on whole cell and single channel I_{ClCa} in freshly dispersed rabbit PA smooth muscle cells, which have been studied extensively by our group (Greenwood *et al.*, 2001; 2004; Piper and Large, 2003), as well as mouse thoracic aortic (TA) smooth muscle cells that express TMEM16A. Secondly, to ascertain the effectiveness of T16A_{inh}-A01 on the contractility of mouse and human blood vessels in comparison to the conventional CACC blocker niflumic acid.

Methods

Initial electrophysiological experiments were performed on single smooth muscle cells isolated from the main PA from New Zealand White rabbits, killed by pentobarbitone sodium overdose in accordance with Schedule 1 of the United Kingdom Animals Act (1986). Additional electrophysiological and functional experiments were performed using TA and

mesenteric arteries (MAs) from BALB/c mice (6–8 weeks old) killed by cervical dislocation (Schedule 1). All studies involving animals are reported in accordance with the ARRIVE guidelines for reporting experiments involving animals (Kilkenny *et al.*, 2010; McGrath *et al.*, 2010). In addition, human visceral adipose tissue samples were taken from elective bariatric surgery at St George's Hospital, London (local ethics approval, 08/H0803/84). Arteries used in the study were dissected free from any damaged or pathologically altered tissue by the operating surgeon and micro-dissected in the laboratory within 1 h. Arteries were identified as the muscular vessel running as a pair with thin walled veins (Mulvany and Halpern, 1977).

Single cell dispersal

Initial electrophysiological experiments to characterize the effects of T16A_{inh}-A01 and F_{act} were performed in PA smooth muscle cells where Cl^- currents due to the activation of CACCs (I_{ClCa}) have been studied extensively at both whole cell and single channel levels (e.g. Greenwood *et al.*, 2001; 2004; Piper *et al.*, 2002; Piper and Large, 2003; Angermann *et al.*, 2006). Single channel experiments were also performed on mouse TA smooth muscle cells where expression of TMEM16A protein has been identified and whole cell I_{ClCa} have been characterized (Davis *et al.*, 2010). Rabbit PA myocytes were isolated by incubating segments of artery in Ca^{2+} -free physiological salt solution (PSS) containing 0.5 $\text{mg}\cdot\text{mL}^{-1}$ papain, 1 mM DTT and 2 $\text{mg}\cdot\text{mL}^{-1}$ BSA overnight at 4°C, followed by trituration through a wide bore glass pipette. TA myocytes were prepared by incubating segments of tissue in PSS containing 50 μM CaCl_2 , 1 $\text{mg}\cdot\text{mL}^{-1}$ collagenase type X and 1 $\text{mg}\cdot\text{mL}^{-1}$ protease type 2A for 25 min.

Whole cell currents

Similar to previous studies (e.g. Greenwood *et al.*, 2001; 2004; Piper *et al.*, 2002; Angermann *et al.*, 2006), whole cell I_{ClCa} were evoked immediately upon membrane rupture when the pipette solution contained a fixed concentration of $[\text{Ca}^{2+}]$. The pipette solution had the following composition (in mM): TEA (20), CsCl (106), HEPES-CsOH (10, pH 7.2), BAPTA (10), ATP.Mg (5) and GTP.diNa (0.2). To this solution, 4.2, 6.8 or 7.08 mM CaCl_2 were added to achieve a free $[\text{Ca}^{2+}]$ of 100, 250 and 500 nM respectively. The bathing solution had the following composition (in mM): NaCl (126), HEPES-NaOH (10, pH 7.35), TEA (8.4), glucose (20), MgCl_2 (1.2) and CaCl_2 (1.8). Voltage protocols were the same as used previously (Greenwood *et al.*, 2001; 2004; Angermann *et al.*, 2006). The effect of T16A_{inh}-A01 (1–30 μM) was studied on I_{ClCa} evoked by 500 nM free Ca^{2+} only, whereas the effect of F_{act} was studied on I_{ClCa} evoked by a range of $[\text{Ca}^{2+}]$ (20–500 nM) to allow comparison with the previous work (Namkung *et al.*, 2011b).

Single channel recordings

Single channel I_{ClCa} have been defined previously in rabbit PA myocytes by Piper and Large (2003). With 500 nM Ca^{2+} activating $[\text{Ca}^{2+}]$, channel activity is characterized by the existence of two sub-conductance states (1.8 and 1.2 pS), but a single 3.5 pS state dominates at lower $[\text{Ca}^{2+}]$ (Piper and Large, 2003). The effects of T16A_{inh}-A01, the less potent TMEM16A blocker tannic acid (Namkung *et al.*, 2010) and F_{act} on single channel I_{ClCa} were studied in outside-out patches using a patch

pipette solution containing (mM) NMDG-Cl [126, equimolar addition of *N*-methyl-D-glucamine (NMDG) and HCl], HEPES (10), MgCl_2 (1.2), EGTA (0.1) and MgATP (1), pH adjusted to 7.2 with NMDG or HCl. The bathing solution for outside-out patches contained (mM) NMDG-Cl (126), CaCl_2 (1.5), MgCl_2 1.2 and HEPES (10), pH adjusted to 7.2 with HCl. The effect of TMEM16A-specific antibodies on I_{ClCa} was studied using the inside-out configuration with a patch pipette solution containing (mM) NMDG-Cl, (126), MgCl_2 (1.2) CaCl_2 (10) and HEPES (10), pH set at 7.2 with NMDG or HCl. The bathing (intracellular) solution had the same composition as the patch pipette solution used in outside-out patches. Single Cl^- channel currents were recorded and analysed according to protocols detailed in Piper and Large (2003).

Immunopharmacological studies

Previously, Manoury *et al.* (2010) showed that siRNA targeted against TMEM16A decreased whole cell I_{ClCa} in rat PA cells in short-term culture. We used an antibody-based approach to ascertain if TMEM16A contributes to native I_{ClCa} in freshly dispersed cells. Consequently, we used an antibody-based approach that has been used previously to characterize TRPC channels in our laboratory (e.g. Albert *et al.*, 2009) to ascertain if TMEM16A contributes to native I_{ClCa} in freshly dispersed cells. In an attempt to circumvent off target effects, we used two different antibodies raised in different hosts (rabbit and goat) against different epitopes and we controlled for non-specific binding by comparing the effect of serum from non-immunized rabbits or goats (see Johnstone and Thorpe, 1996). In addition, an antibody against bestrophin-3, which underlies the cGMP-dependent CACC (Matchkov *et al.*, 2008), was also applied to excised patches as a control. Single channels were recorded in the inside-out configuration with bath $[\text{Ca}^{2+}]$ set to 500 nM in both PA and mouse TA cells. In single channel recordings using rabbit PA cells either sc-69343, a goat polyclonal antibody targeted to an internal epitope of human TMEM16A (Santa Cruz Biotechnology, Insight Biotech, Wembley, UK) or the bestrophin-3 antibody (ab94904, Abcam Ltd., Cambridge, UK) was applied. In the case of single channel electrophysiology using mouse cells, both sc-69343 and ab72984, a rabbit polyclonal raised against a peptide derived from the N-terminal region of human TMEM16A (Abcam) was applied. We have characterized ab72984 by Western blot in our previous work (Davis *et al.*, 2010).

Isometric tension recordings

The functional effects of T16A_{inh}-A01, tannic acid and niflumic acid were investigated in segments of mouse TA and MA as well as human visceral adipose arteries. All segments were transferred to Krebs' solution and then mounted on a wire myograph (DMT, Aarhus, Denmark). All resistance vessels were normalized to 90% of the diameter of the vessel at 100 mmHg using a passive stretch/tension relationship (Mulvany and Halpern, 1977), whereas the mouse TA segments were set to an initial tension of 12 mN (Yeung *et al.*, 2007). After a 30 min period of equilibration, segments were challenged twice with 60 mM KCl for 5 min. After washout and stabilization of tension, the vessels were then pre-contracted with either 3 μM methoxamine (TA) or 100 nM U46619 (MA/human arteries; Tocris, Bristol, UK) to obtain robust, stable contractions. After

the response had stabilized either dimethyl sulfoxide (DMSO) or the following Cl^- channel blockers – niflumic acid, T16A_{inh}-A01 or tannic acid – were applied at increasing concentrations to generate dose–response relationships. A second protocol was also employed in the mouse TA where increasing concentrations of methoxamine (100 nM–10 μM) were applied cumulatively. Tissues were then washed with Krebs' solution and incubated in either DMSO, T16A_{inh}-A01 (5 μM), F_{act} (10 μM), tannic acid (30 μM) or niflumic acid (10 and 100 μM) for 30 min before a second methoxamine concentration–effect curve was constructed in the presence of these agents. The composition of the Krebs' solution (mM) is as follows: NaCl (125), KCl (4.6), CaCl_2 (2.5), NaHCO_3 (25.4), Na_2HPO_4 (1), MgSO_4 (0.6) and glucose (10) bubbled with 95% O_2 /5% CO_2 at 37°C. PSS for dispersal (mM) is as follows: NaCl (126), KCl (6), glucose (10), HEPES (11), MgCl_2 (1.2) and CaCl_2 (1.5), with pH adjusted to 7.2 with NaOH.

Endpoint PCR

Total RNA was extracted from arteries, as well as pools of single smooth muscle cells from either rabbit PA or mouse TA, and reverse transcribed to cDNA as described previously (Davis *et al.*, 2010). Primer sequences and expected amplicon size are found in the Online Supplement (Table S1).

Immunohistochemistry

Protein detection in slices of artery was performed as described previously (Davis *et al.*, 2010; Ng *et al.*, 2011) and further details are included in the Online Supplement.

Immunocytochemistry

TMEM16A protein was detected in single smooth muscle cells from mouse TA using methodology described previously (Davis *et al.*, 2010). Cells were imaged using a Zeiss confocal microscope (Zeiss, Welwyn, UK) and membrane localization of TMEM16A protein in any Z-section was determined by overlap of fluorescence from an $\alpha 1$ sodium (Na^+)–potassium (K^+) ATPase antibody (Abcam; ab7671). As no control blocking peptide was supplied with the TMEM16A antibody, we used a boiled version of this sample in accordance with the previous work in cerebral arteries (Thomas-Gatewood *et al.*, 2011).

Materials

All chemicals were purchased from Sigma (Poole, Dorset, UK), unless otherwise stated.

Statistical analysis

All data are shown as the mean of *n* animals/patients (where appropriate) $\pm$ SEM. Statistical significance was determined by unpaired Student's *t*-test using GraphPad Prism software (GraphPad Software Inc., San Diego, CA, USA). All nomenclature is consistent with Alexander *et al.* (2011).

Results

In agreement with previous studies (Davis *et al.*, 2010; Sones *et al.*, 2010), expression of TMEM16A was identified in whole mouse TA and MA by endpoint PCR (Figure 1A). Amplicons for TMEM16A were also identified using RNA extracted from




Figure 1

Identification of TMEM16A expression in mouse and human arteries. (A) Endpoint PCR analysis of TMEM16A amplicons from whole mouse TA (mTA) and mesenteric artery (mMA) tissue (Ai), enzymatically isolated smooth muscle cells from rabbit pulmonary artery (rbPA) and mTA (Aii) and human visceral adipose arteries (hVAA; Aiii) confirms expression of *TMEM16A* gene in all tissues tested. The # denotes a different RNA extraction from new material. NTC = no template control and M = 100 base pair (bp) ladder. (B) Immunohistochemical analysis of transverse sections of mTA, mMA and hVAA show specific TMEM16A expression (red staining/Alexa Fluor 546) in the smooth muscle α -actin layer (green staining/Alexa Fluor 488) of all tissues tested. All images presented were exposure time-matched and are representative of four separate experiments, with four different animal/patient samples. Staining surrounding the vessels represents extraneous fat and (in the case of mMA) the paired vein can also be seen and were used to aid in the identification of the arteries. (C) Immunocytochemical analysis of enzymatically isolated mTA smooth muscle cells shows distinct overlap of TMEM16A expression (red staining/Alexa Fluor 546) with $\alpha 1$ sodium potassium ATPase (green staining/Alexa Fluor 488; a plasma membrane marker).

a pool of smooth muscle cells from both rabbit PA and mouse TA, as well as in all samples of human visceral adipose artery studied (confirmed by sequencing of product). Translation of *TMEM16A* gene to protein was confirmed by immunohistochemistry (Figure 1B) using an antibody that has been characterized thoroughly in overexpression systems and native protein by Davis *et al.* (2010). Figure 1B shows that TMEM16A localized to the smooth muscle region in human blood vessels as defined by the smooth muscle α -actin staining. In immunocytochemical experiments of enzymatically isolated smooth muscle cells, TMEM16A expression products were observed in the cytoplasm but predominantly co-localized with $\alpha 1$ sodium potassium ATPase, suggesting membrane localization (Figure 1C).

Electrophysiology

Consistent with previous studies (Greenwood *et al.*, 2001; 2004; Angermann *et al.*, 2006), distinctive whole cell I_{ClCa} with characteristic voltage-dependent kinetics were evoked immediately upon membrane rupture with a pipette solution containing 500 nM free Ca^{2+} . These currents exhibited considerable rundown within the first 2 min of recording (see Angermann *et al.*, 2006) but then remained stable (data not shown). Application of 10 μ M T16A_{inh}-A01 consistently inhibited a component of the evoked current after an initial lag period of a few minutes (Figure 2A,B) and this was readily reversed upon washout (Figure 2A). Inhibition of the evoked current was also observed with 1 μ M T16A_{inh}-A01 (Figure 2D),




Figure 2

Effect of T16A_{inh}-A01 on whole cell I_{ClCa} in rabbit PA smooth muscle cells. (A) shows a representative time course for the effect of 10 μM T16A_{inh}-A01 and subsequent washout on whole cell I_{ClCa} recorded at +70 mV. The gap represents the application of a current-voltage protocol. (B) shows representative currents recorded at voltages between -100 and +100 mV in the absence (Bi) and presence (Bii) of 10 μM T16A_{inh}-A01. (C) shows the T16A_{inh}-A01-sensitive current determined as (Bi) and (Bii). The mean T16A_{inh}-A01-sensitive current at different concentrations is shown in (D). Each point is the mean of 4–6 cells $\pm$ SEM. * $P < 0.05$; ** $P < 0.01$.

while 30 μM T16A_{inh}-A01 had no greater effect than 10 μM T16A_{inh}-A01 (Figure 2D). Previously, we found that niflumic acid had inhibitory and stimulatory effects on whole cell I_{ClCa} in rabbit PA smooth muscle cells that were manifested as an increase in currents at negative test potentials and enhancement of the current to approximately 200% of pre-drug levels upon washout (Piper *et al.*, 2002; see also Supporting Information Figure S1). However, no stimulatory effect was observed with T16A_{inh}-A01 either in the presence of the agent or upon washout (Figure 2, current after 5 min washout of 10 μM T16A_{inh}-A01 was $105 \pm 5\%$ of pre-drug levels, $n = 4$). In contrast to T16A_{inh}-A01, the TMEM16A potentiator F_{act} (10 μM) had no significant effect on I_{ClCa} evoked by 500 nM

free Ca^{2+} (Figure 3A) with the current at +90 mV being $15. \pm 7$ and 14.5 ± 6 $\text{pA} \cdot \text{pF}^{-1}$ in the absence and presence of F_{act} respectively. However, when I_{ClCa} was evoked by pipette solutions containing 100 or 250 nM free Ca^{2+} , the currents were enhanced considerably (Figure 3B,C).

Single channel studies

In outside-out patches from PA smooth muscle cells, no channel activity was present when the pipette solution was Ca^{2+} free ($n = 5$, Figure 4A). However, inclusion of 500 nM free Ca^{2+} in the patch pipette solution activated single I_{ClCa} channel currents (Figure 4B), which had a mean open probability (NP_o) of 0.68 ± 0.07 ($n = 12$) at -100 mV. I_{ClCa} channel




Figure 3

Effect of F_{act} on whole cell I_{CaCa} in rabbit PA smooth muscle cells. (A) shows currents evoked by 500 nM Ca^{2+} before (Ai) and after application of 10 μM F_{act} for 5 min (Aii). (B) shows currents evoked by 100 nM Ca^{2+} before (Bi) and after application of 10 μM F_{act} for 5 min (Bii). (C) shows the mean increase in current produced by 10 μM F_{act} at different $[\text{Ca}^{2+}]$. Current in the presence of the drug was normalized to the pre-drug amplitude. Each bar represents the mean of 4–7 cells $\pm$ SEM; ** $P < 0.01$, paired Student's t -test.

currents evoked by 500 nM free Ca^{2+}_i had two amplitudes of -0.16 ± 0.02 pA ($n = 12$) and -0.29 ± 0.02 pA ($n = 12$) at -100 mV. These two channel amplitudes are consistent with the two sub-conductance levels of 1.8 and 3.5 pS reported for I_{CaCa} in these cells previously (Piper and Large, 2003). Similar channel activity was evoked in inside-out patches by a bath solution containing 500 nM Ca^{2+} . Figure 4C shows that co-application of an anti-TMEM16A antibody (1:100 sc-69343), but not an anti-Bestrophin-3 antibody (1:200; ab94904), significantly reduced I_{CaCa} channel activity evoked by 500 nM Ca^{2+}_i , with NP_o decreasing from 0.61 ± 0.07 to 0.03 ± 0.01 ($n = 7$). In control experiments on smooth muscle cells from rabbit MA, bath application of the anti-bestrophin-3 antibody blocked I_{ClcGMP} channel activity by $95 \pm 2\%$ ($n = 5$, Figure 4D).

Identical single channel activity was recorded in smooth muscle cells from mouse TA upon application of 500 nM Ca^{2+}_i (Figure 4E). Application of the TMEM16A antibody sc-69343 (raised in goat) inhibited the activity of I_{CaCa} channels by $90 \pm 3\%$ ($n = 5$), similar to the data in rabbit PA (Figure 4C). A second TMEM16A antibody (ab72984, raised in rabbit) also inhibited channel activity markedly ($97 \pm 1\%$; Figure 4G). In contrast, serum from non-immunized goat or rabbit had no effect on I_{CaCa} channel activity (Figure 4F,G). Consequently, I_{CaCa} channel activity in rabbit PA and mouse TA was decreased considerably by two TMEM16A antibodies raised in different hosts and against different epitopes.

Effects of pharmacological agents on single channel I_{CaCa}

Experiments were performed to determine the effect of the TMEM16A modulators on single channel I_{CaCa} in rabbit PA and mouse TA smooth muscles. In outside-out recordings from rabbit PA smooth muscle cells application of 10 μM T16A_{inh}-01 inhibited I_{CaCa} channel activity evoked by 500 nM free Ca^{2+}_i by $93 \pm 2\%$ ($n = 6$) at -100 mV, which was partially reversed upon washout (Figure 5A,B). I_{CaCa} channel activity was also blocked by bath application of 30 μM tannic acid, which is a less potent blocker of TMEM16A channels (Namkung *et al.*, 2010; 2011a). Figure 5C shows that T16A_{inh}-01 also inhibited the single channel activity in mouse TA smooth muscle cells that was readily reversed upon washout.

A final series of experiments investigated the effect of the TMEM16A potentiator F_{act} on single channel I_{CaCa} from rabbit PA smooth muscle cells activated with either 100 or 500 nM free Ca^{2+} . With the lower $[\text{Ca}^{2+}]$, single channel activity was considerably less (NP_o was 0.65 ± 0.1 vs. 0.14 ± 0.02 , $n = 5$) and was dominated by transitions to the larger sub-conductance state (cf. Figures 5A and 6A). In agreement with the whole cell data, F_{act} had no significant effect upon single channel activity evoked by 500 nM Ca^{2+} (NP_o was 0.65 ± 0.12 and 0.76 ± 0.08 before and after F_{act} , $n = 3$) but produced a marked increase in channel activity evoked by 100 nM Ca^{2+} (Figure 6A,Cii).




Figure 4

Single channel recordings in rabbit PA and mouse TA smooth muscle cells. I_{ClCa} Activity in outside-out patches rabbit PA smooth muscle cells is evoked following inclusion of 500 nM free Ca^{2+}_i in the patch pipette solution (A,B). In the insert, (C) denotes closed state, O_1 and O_2 are different open states. Subpanels in (B) show the different channel amplitudes of individual opening on a faster time scale and their amplitude histogram. (C) shows a representative effect of antisera raised against TMEM16A (sc-69343). Antisera against bestrophin-3 had no effect on this channel activity (D) but inhibited cGMP-stimulated I_{ClCa} channel activity in rabbit mesenteric artery cells. (E) shows a representative trace for the effect of a TMEM16A antibody (sc-69343) on single Cl^- channel activity in mouse TA smooth muscle cells. (F,G) show the mean effect of both TMEM16A antibodies, sc-69343 (1:100) and ab72984 (1:500) on I_{ClCa} channel activity in inside-out patches from mouse TA cells evoked by applying 500 nM free Ca^{2+} ($n = 5$). Note that the corresponding rabbit and goat non-immune serum had no effect on I_{ClCa} channel activity. * $P < 0.05$; *** $P < 0.001$.

Whole tissue functional studies

The simple working hypothesis is that TMEM16A functions as a CACC that when opened produces membrane depolarization sufficient to increase Ca^{2+} influx through VDCC. Consequently, blockers of TMEM16A should relax pre-contracted vessels and suppress vascular contractions. Pre-contraction of mouse TA by application of the α 1-adrenoceptor agonist methoxamine produced robust contractions that were inhibited in a concentration-dependent manner by T16A_{inh}-A01 with an apparent IC_{50} of 1.7 μ M derived from the averaged data (Figure 7A,D), whereas the equivalent volume of DMSO had negligible effect (Figure 7C; mean relaxation was $2 \pm 1\%$, $n = 5$). The effect of T16A_{inh}-A01 was relatively slow to develop, with a maximum effect occurring after 18 ± 4 min for 10 μ M ($n = 8$) but was well maintained for at least 60 min (data not shown). The prototypic CACC blocker niflumic acid (Figure 7B,D), currently considered the most effective blocker

of I_{ClCa} (see Greenwood and Leblanc, 2007), inhibited contractions in a concentration-dependent manner, although at higher concentrations compared with T16A_{inh}-A01 (Figure 7A,B). Tannic acid was a less effective relaxant of mouse TA segments (Figure 7D), although subsequent application of niflumic acid or T16A_{inh}-A01 produced marked relaxation of the tissues in the continued presence of tannic acid ($n = 4$, data not shown).

Incubation of TA segments with a submaximal concentration of T16A_{inh}-A01 (5 μ M) suppressed the subsequent methoxamine concentration–effect curve to a greater extent than 30 μ M tannic acid or 10 μ M niflumic acid (Figure 7E). In contrast, F_{act} had no significant effect on the methoxamine-induced contractions ($n = 5$, data not shown). Furthermore, 100 μ M niflumic acid abrogated the methoxamine concentration-dependent contractions (data not shown) but also abolished contractions induced by 60 mM KCl, suggesting a direct effect on VDCCs. T16A_{inh}-A01 had no significant




Figure 5

Effect of T16A_{inh}-A01 on single channel recordings in rabbit PA and mouse TA smooth muscle cells. (A) shows an example of the inhibitory effect of 10 μM T16A_{inh}-A01 on I_{Ca} activity in outside-out patches from rabbit PA cells evoked following inclusion of 500 nM free Ca²⁺_i in the patch pipette solution. In the insert, (C) denotes closed state, O₁ and O₂ are different open states. Subpanels show the different channel amplitudes of individual opening on a faster time scale and their amplitude histogram. (B) shows the mean change in NP_o produced by 10 μM T16A_{inh}-A01 and 30 μM tannic acid. (C) shows the concentration-dependent inhibition produced by 1 and 10 μM T16A_{inh}-A01 (*n* = 6) of channel activity evoked by 500 nM Ca²⁺ in mouse TA smooth muscle cells. **P* < 0.05; ****P* < 0.001.

effect on contractions produced by application of 60 mM KCl (1.93 ± 0.6 vs. 1.7 ± 0.5 mN, *n* = 5) and the inhibitory effect of 10 μM T16A_{inh}-A01 was not affected by incubation of the TA with a cocktail of K⁺ channel blockers. Thus, the relaxation to 10 μM T16A_{inh}-A01 in 10 μM glibenclamide (ATP-sensitive K⁺ channel blocker), 10 μM paxilline (BK_{Ca} blocker) and 1 μM linopirdine (Kv7 channel blocker, Yeung *et al.*, 2007) was $82 \pm 6\%$ (*n* = 4) compared with $74 \pm 5\%$ in the absence of these agents (*n* = 8). These studies suggest that T16A_{inh}-A01 does not relax mouse TA through direct block of VDCCs or activation of K⁺ channels.

Effect of T16A_{inh}-A01 in mouse and human mesenteric arteries

Having established that T16A_{inh}-A01 was an effective vasorelaxant of mouse TA, experiments were performed to ascertain the impact of this agent in arteries that have a crucial role in blood pressure regulation. Figure 8A shows that 10 μM T16A_{inh}-A01 was a highly effective relaxant of mouse MA

pre-contracted with the stable thromboxane mimetic, U46619, compared with the equivalent vehicle (*P* < 0.0001, *n* = 4). Figure 8B shows that 10 μM T16A_{inh}-A01 also relaxed U46619-induced contractions of human visceral adipose arteries. The mean data for the experiments on mouse MA and human visceral adipose arteries are shown in Figure 8C. These data provide evidence for a functional role of TMEM16A in the vasoconstrictor responses of murine TA and MA as well as human resistance vessels.

Discussion

In the present study, we have shown for the first time that a small molecule blocker of TMEM16A (T16A_{inh}-A01; Namkung *et al.*, 2011a) and the recently identified TMEM16A activator F_{act} (Namkung *et al.*, 2011b) have marked effects on native I_{Ca} recorded in vascular smooth muscle cells. These data add weight to the view that




Figure 6

Effect of F_{act} on single channel recordings in rabbit PA smooth muscle cells. (A) shows an example of the effect of $10 \mu\text{M } F_{\text{act}}$ on single channel evoked with $100 \text{ nM } \text{Ca}^{2+}$ in the pipette solution. (B) shows amplitude histograms in the absence (i) and presence of $10 \mu\text{M } F_{\text{act}}$ (ii). Mean channel activity $\pm F_{\text{act}}$ for 500 nM and $100 \text{ nM } \text{Ca}^{2+}$ is shown in (Ci) and (Cii) respectively ($n = 3-5$).

TMEM16A constitutes a major component of native I_{ClCa} in freshly dispersed smooth muscle cells and agree with Manoury *et al.* (2010) who showed that TMEM16A targeted siRNA reduced I_{ClCa} in rat PA smooth muscle cells maintained in short-term culture. Moreover, single channel I_{ClCa} in smooth muscle cells from rabbit PA and mouse TA were abolished by two TMEM16A-specific antibodies raised in different hosts and against different epitopes. These inhibitory effects were not seen with non-immune serum from the host animals or antisera against bestrophin-3, which comprises the cGMP-sensitive I_{ClCa} in rat MA (Matchkov *et al.*, 2008) and is expressed in rabbit PA (Leblanc *et al.*, 2005). While there are caveats to the use of antibodies as pharmacological tools and the precise mechanism by which these polyclonal antibodies produced an inhibitory effect is not known, antibodies have been used as blocking molecules in the characterization of a number of different ion channels (e.g. Albert *et al.*, 2009), including CACC studies (Thomas-Gatewood *et al.*, 2011). Consequently, the fact that two structurally different antibodies targeted against TMEM16A proteins, and a

TMEM16A blocker, exert an inhibitory effect on I_{ClCa} provides evidence that TMEM16A contributes to native CACCs. However, it is likely that the native CACC in vascular smooth muscle is not composed solely of TMEM16A. Firstly, the single channel conductance of native CACCs ($1-3 \text{ pS}$) determined in the rabbit PA (present study, Piper and Large, 2003) as well as other blood vessels (Klößner, 1993; Van Renterghem and Lazdunski, 1993; Hirakawa *et al.*, 1999) is smaller than that determined for TMEM16A channels in overexpression studies ($\sim 8 \text{ pS}$; Yang *et al.*, 2008), although it is similar to channels generated by heterologously expressed TMEM16B (Pifferi *et al.*, 2009). Secondly, in our whole cell recordings, only a component of the evoked current was inhibited by T16A_{inh}-A01 over the same concentration range where full blockade of TMEM16A-generated currents is seen (Namkung *et al.*, 2011a). These differences may represent variable contribution from the different TMEM16A splice variants (Ferrera *et al.*, 2009) that are expressed in vascular smooth muscle (Davis *et al.*, 2010; Manoury *et al.*, 2010) or the native CACC comprises TMEM16A proteins in




Figure 7

Effect of T16A_{inh}-A01 in isometric tension recordings from murine TA. Representative effects of 1 μM and 10 μM T16A_{inh}-A01 (A), niflumic acid (10–100 μM, B) and equivalent DMSO on TA pre-contracted by 3 μM methoxamine (applied at •). (D) shows relaxant concentration–effect curves for niflumic acid, T16A_{inh}-A01 and tannic acid on pre-contracted mouse TA. All data are the mean of 5–8 experiments ± SEM. (E) shows the mean effect of cumulative application of methoxamine in the presence of DMSO, 5 μM T16A_{inh}-A01, 30 μM tannic acid and 10 μM niflumic acid. All points are the mean of 5 experiments ± SEM.

association with other proteins. These aspects will be the focus of future studies.

The present study shows that T16A_{inh}-A01 relaxed pre-contracted mouse TA with an IC₅₀ of 1.7 μM, which is similar to the derived value for inhibition of epithelial short circuit current (Namkung *et al.*, 2011a) and for reduction of I_{ClCa} in the present study. T16A_{inh}-A01 at 10 μM was also a remarkably effective relaxant of pre-contracted resistance arteries and human visceral adipose arteries that we showed express TMEM16A and has recently been shown to relax 5HT-precontracted rat pulmonary arteries (Sun *et al.*, 2012). T16A_{inh}-A01 was considerably more potent vasorelaxant than conventional Cl channel blockers such as niflumic acid (see Criddle *et al.*, 1996 and present study). These findings are consistent with a simplified working model, whereby activation of a membrane receptor leads to a release of calcium sufficient to activate CACCs, which leads to Cl[−] efflux because smooth muscle cells accumulate Cl[−] ions (Chipperfield and Harper, 2000), and the membrane depolarization increases Ca²⁺ influx through voltage-dependent Ca²⁺ channels (VDCCs, Kitamura and Yamazaki, 2001; Leblanc *et al.*, 2005). A blocker of CACCs would therefore limit the agonist-induced membrane depolarization and impair voltage-dependent calcium influx. The ability of a TMEM16A-specific

blocker to impair vasoconstriction therefore suggests that TMEM16A expression products contribute to the native CACCs or some undefined part of the working model. However, using pharmacological agents to define a functional role requires a degree of caution. For instance, a similar degree of vasorelaxation would be produced by direct block of calcium channels or activation of potassium channels that would indirectly close the same channels. With respect to this point, T16A_{inh}-A01 had no significant effect on contractions produced by high KCl and the inhibitory effect of this agent was unaffected by paxilline, glibenclamide or linopirdine, effectively repudiating a role for ATP-sensitive K⁺ channels, KCa3.1 or KCNQ-encoded K⁺ channels that are present in this tissue. In addition, T16A_{inh}-A01 was equally as effective against methoxamine-induced or U46619-induced contractions, suggesting that this agent is not functioning as a receptor blocker. Having shown that the TMEM16A potentiator, F_{act}, increased single channel and whole cell I_{ClCa}, we hypothesized that this agent would augment the methoxamine-induced concentration–effect curve in line with the model described earlier. However, this was not the case and F_{act} did not produce a contraction by itself. It is possible that the activator was more effective at stimulating the current at positive than at negative potentials, which could have




Figure 8

Effect of T16A_{inh}-A01 in isometric tension recordings from mouse and human resistance arteries. Ai shows the rapid effect of 10 μ M T16A_{inh}-A01 on mouse MA pre-contracted with U46619. Aii shows the lack of effect of DMSO equivalent. Scale bars denote 8 min and 2 mN. Representative traces for the effect of 10 μ M T16A_{inh}-A01 (Bi) and the equivalent DMSO volume (Bii) on segments of human visceral adipose arteries pre-contracted with U46619. Scale bars denote 2 mN and 8 min for both (Bi and Bii). (C) shows the mean relaxant effect for 10 μ M T16A_{inh}-A01 in human visceral adipose arteries and mouse MA. Each column is the mean of 4–6 animals $\pm$ SEM. Open bars show the DMSO equivalent. Application of vasoconstrictor in all panels is shown by (•).

resulted in minimal effects of the drug in the physiological range of membrane potentials influenced by methoxamine and future studies will investigate the vascular effects of this agent and F_{act} (Namkung *et al.*, 2011b) further.

In summary, the identification of TMEM16A expression products as constituents of native CACCs has energized the CACC research field and has allowed for novel reagents to be developed, which can provide insight on the physiological role of these proteins in various tissues. We now provide further evidence that TMEM16A proteins comprise native CACCs in vascular smooth muscle cells and inhibiting TMEM16A has marked anti-contractile effects in various mouse and human arteries.

Acknowledgements

We thank all the patients and associated staff at St George's Hospital for providing human samples. We also thank Mr Thomas Jepps and the Image Resource Facility at SGUL for their support with the immunological experiments. This work was supported by the British Heart Foundation grants awarded to I. A. G. (PG/09/104 and PG/07/127/24235), APA (PG/08/042/2506, BB/J007226/1) and by grants to N. L. from

the National Institutes of Health (Grants 5 R01 HL 075477 and 3 R01 HL075477-04S1).

Conflict of interests

None of the authors have any conflict of interests.

References

- Albert AP, Saleh SN, Large WA (2009). Identification of canonical transient receptor potential (TRPC) channel proteins in native vascular smooth muscle cells. *Curr Med Chem* 16: 1158–1165.
- Alexander SPH, Mathie A, Peters JA (2011). Guide to receptors and channels (GRAC), 5th edition. *Br J Pharmacol* 164: S1–S324.
- Angermann JE, Sanguinetti AR, Kenyon JL, Leblanc N, Greenwood IA (2006). Mechanism of the inhibition of Ca²⁺-activated Cl⁻ currents by phosphorylation in pulmonary arterial smooth muscle cells. *J Gen Physiol* 128: 73–87.
- Caputo A, Caci E, Ferrera L, Pedemonte N, Barsanti C, Sondo E *et al.* (2008). TMEM16A, a membrane protein associated with calcium-dependent chloride channel activity. *Science* 322: 590–594.

- Chipperfield AR, Harper AA (2000). Chloride in smooth muscle. *Prog Biophys Mol Biol* 74: 175–221.
- Criddle DN, de Moura RS, Greenwood IA, Large WA (1996). Effect of niflumic acid on noradrenaline-induced contractions of the rat aorta. *Br J Pharmacol* 118: 1065–1071.
- Davis AJ, Forrest AS, Jepps TA, Valencik ML, Wiwchar M, Singer CA *et al.* (2010). Expression profile and protein translation of TMEM16A in murine smooth muscle. *Am J Physiol Cell Physiol* 299: C948–C959.
- Ferrera L, Caputo A, Ubby I, Bussani E, Zegarra-Moran O, Ravazzolo R *et al.* (2009). Regulation of TMEM16A chloride channel properties by alternative splicing. *J Biol Chem* 284: 33360–33368.
- Greenwood IA, Leblanc N (2007). Overlapping pharmacology of Ca^{2+} -activated Cl^- and K^+ channels. *Trends Pharmacol Sci* 28: 1–5.
- Greenwood IA, Ledoux J, Leblanc N (2001). Differential regulation of Ca^{2+} -activated Cl^- currents in rabbit arterial and portal vein smooth muscle cells by Ca^{2+} -calmodulin-dependent kinase. *J Physiol* 534: 395–408.
- Greenwood IA, Ledoux J, Sanguinetti A, Perrino BA, Leblanc N (2004). Calcineurin $\text{A}\alpha$ but not $\text{A}\beta$ augments $\text{I}_{\text{Cl}(\text{Ca})}$ in rabbit pulmonary artery smooth muscle cells. *J Biol Chem* 279: 38830–38837.
- Hirakawa Y, Gericke M, Cohen RA, Bolotina VM (1999). Ca^{2+} -dependent Cl^- channels in mouse and rabbit aortic smooth muscle cells: regulation by intracellular Ca^{2+} and NO. *Am J Physiol* 277: H1732–H1744.
- Johnstone AP, Thorpe RC (1996). *Immunohistochemistry in Practice*, 3rd edn. Wiley-Blackwell: Oxford.
- Kilkenny C, Browne W, Cuthill IC, Emerson M, Altman DG (2010). NC3Rs Reporting Guidelines Working Group. *Br J Pharmacol* 160: 1577–1579.
- Kitamura K, Yamazaki J (2001). Chloride channels and their functional roles in smooth muscle tone in the vasculature. *Jpn J Pharmacol* 85: 351–357.
- Klößner U (1993). Intracellular calcium ions activate a low-conductance chloride channel in smooth muscle cells isolated from human mesenteric artery. *Pflügers Arch* 424: 231–237.
- Large WA, Wang Q (1996). Characteristics and physiological role of the Ca^{2+} -activated Cl^- conductance in smooth muscle. *Am J Physiol* 271: C435–C454.
- Leblanc N, Ledoux J, Saleh S, Sanguinetti A, Angermann J, O'Driscoll K *et al.* (2005). Regulation of calcium-activated chloride channels in smooth muscle cells: a complex picture is emerging. *Can J Physiol Pharmacol* 83: 541–556.
- Manoury B, Tamuleviciute A, Tammaro P (2010). TMEM16A/anoctamin 1 protein mediates calcium-activated chloride currents in pulmonary arterial smooth muscle cells. *J Physiol* 588: 2305–2314.
- Matchkov VV, Larsen P, Bouzinova EV, Rojek A, Boedtker DM, Golubinskaya V *et al.* (2008). Bestrophin-3 (vitelliform macular dystrophy2-like 3 protein) is essential for the cGMP-dependent calcium-activated chloride conductance in vascular smooth muscle cells. *Circ Res* 103: 864–872.
- McGrath J, Drummond G, Kilkenny C, Wainwright C (2010). Guidelines for reporting experiments involving animals: the ARRIVE guidelines. *Br J Pharmacol* 160: 1573–1576.
- Mulvany MJ, Halpern W (1977). Contractile properties of small arterial resistance vessels in spontaneously hypertensive and normotensive rats. *Circ Res* 41: 19–26.
- Namkung W, Thiagarajah JR, Phuan PW, Verkman AS (2010). Inhibition of Ca^{2+} -activated Cl^- channels by gallotannins as a possible molecular basis for health benefits of red wine and green tea. *FASEB J* 24: 4178–4186.
- Namkung W, Phuan PW, Verkman AS (2011a). TMEM16A inhibitors reveal TMEM16A as a minor component of calcium-activated chloride channel conductance in airway and intestinal epithelial cells. *J Biol Chem* 286: 2365–2374.
- Namkung W, Yao Z, Finkbeiner WE, Verkman AS (2011b). Small-molecule activators of TMEM16A, a calcium-activated chloride channel, stimulate epithelial chloride secretion and intestinal contraction. *FASEB J* 25: 4048–4062.
- Ng FL, Davis AJ, Jepps TA, Harhun MI, Yeung SY, Wan A *et al.* (2011). Expression and function of the K^+ channel KCNQ genes in human arteries. *Br J Pharmacol* 162: 42–53.
- Pifferi S, Dibattista M, Menini A (2009). TMEM16B induces chloride currents activated by calcium in mammalian cells. *Pflügers Arch* 458: 1023–1038.
- Piper AS, Large WA (2003). Multiple conductance states of single Ca^{2+} -activated Cl^- channels in rabbit pulmonary artery smooth muscle cells. *J Physiol* 547: 181–196.
- Piper AS, Greenwood IA, Large WA (2002). Dual effect of blocking agents on Ca^{2+} -activated Cl^- currents in rabbit pulmonary artery smooth muscle cells. *J Physiol* 539: 119–131.
- Schroeder BC, Cheng T, Jan YN, Jan LY (2008). Expression cloning of TMEM16A as a calcium-activated chloride channel subunit. *Cell* 134: 1019–1029.
- Sones WR, Davis AJ, Leblanc N, Greenwood IA (2010). Cholesterol depletion alters amplitude and pharmacology of vascular calcium-activated chloride channels. *Cardiovasc Res* 87: 476–484.
- Sun H, Xia Y, Paudel O, Yang XR, Sham JS (2012). Chronic hypoxia-induced upregulation of Ca^{2+} -activated Cl^- channel in pulmonary arterial myocytes: a mechanism contributing to enhanced vasoreactivity. *J Physiol* 590: 3507–3521.
- Thomas-Gatewood C, Neeb ZP, Bulley S, Adebisi A, Bannister JP, Leo MD *et al.* (2011). TMEM16A channels generate Ca^{2+} -activated Cl^- currents in cerebral artery smooth muscle cells. *Am J Physiol Heart Circ Physiol* 301: H1819–H1827.
- Van Renterghem C, Lazdunski M (1993). Endothelin and vasopressin activate low conductance chloride channels in aortic smooth muscle cells. *Pflügers Arch* 425: 156–163.
- Yang YD, Cho H, Koo JY, Tak MH, Cho Y, Shim WS *et al.* (2008). TMEM16A confers receptor-activated calcium-dependent chloride conductance. *Nature* 455: 1210–1215.
- Yeung SY, Pucovsky V, Moffatt JD, Saldanha L, Schwake M, Ohya S *et al.* (2007). Molecular expression and pharmacological identification of a role for Kv7 channels in murine vascular reactivity. *Br J Pharmacol* 151: 758–770.

Supporting information

Additional Supporting Information may be found in the online version of this article:

Figure S1 Effect of niflumic acid on whole cell I_{ClCa} evoked in rabbit PA smooth muscle cells.

Table S1 PCR primer pair sequences used to probe for TMEM16A in various tissues.