

## RESEARCH PAPER

# Noradrenaline stimulates cell proliferation by suppressing potassium channels via $G_{i/o}$ -protein-coupled $\alpha_{1B}$ -adrenoceptors in human osteoblasts

D Kodama and A Togari

*Department of Pharmacology, School of Dentistry, Aichi-Gakuin University, Nagoya, Japan*

### Correspondence

Akifumi Togari, Department of Pharmacology, School of Dentistry, Aichi-Gakuin University, 1-100 Kusumoto-cho, Chikusa-ku, Nagoya 464-8650, Japan. E-mail: togariaf@dpc.aichi-gakuin.ac.jp

### Keywords

sympathetic nervous system; osteoblast; potassium channels; cell proliferation; heterotrimeric G-protein

### Received

19 March 2012

### Revised

20 September 2012

### Accepted

24 September 2012

## BACKGROUND AND PURPOSE

Recent studies demonstrated that the sympathetic nervous system regulates bone metabolism via  $\beta_2$ -adrenoceptors. Although  $\alpha$ -adrenoceptors are also expressed in osteogenic cells, their functions in bone metabolism have been less studied. We previously demonstrated that noradrenaline suppressed potassium currents via  $\alpha_{1B}$ -adrenoceptors in the human osteoblast SaM-1 cell line. The aim of this study was to investigate the signal transduction pathway and the physiological role of noradrenaline in human osteoblasts in more detail.

## EXPERIMENTAL APPROACH

To investigate signal transduction through  $\alpha_{1B}$ -adrenoceptors, we used whole-cell patch clamp recording and Ca fluorescence imaging. Potassium channels regulate membrane potential and cell proliferation activity in non-excitable cells, so we evaluated cell proliferation activity by BrdU incorporation and WST assay.

## KEY RESULTS

In SaM-1 cells, bath-applied noradrenaline elevated intracellular  $Ca^{2+}$  concentration and this effect was abolished by both chloroethylclonidine, an  $\alpha_{1B}$ -adrenoceptor antagonist, and U73122, a PLC inhibitor. However, the inhibitory effect of noradrenaline on whole-cell current was unaffected by U73122. In contrast, in cells pretreated with either *Pertussis* toxin, a  $G_{i/o}$ -protein-coupled receptor inhibitor, or gallein, a  $G\beta\gamma$ -protein inhibitor, the inhibitory effect of noradrenaline on whole-cell current was significantly suppressed. Noradrenaline-induced enhancement of cell proliferation was inhibited by CsCl, a non-selective potassium channel blocker, gallein and H89, a PKA inhibitor, but not by U73122.

## CONCLUSIONS AND IMPLICATIONS

Noradrenaline facilitated cell proliferation by regulation of potassium currents in human osteoblasts via  $G_{i/o}$ -protein-coupled  $\alpha_{1B}$ -adrenoceptors, not via coupling to Gq-proteins.

## Abbreviations

$\alpha$ -MEM,  $\alpha$ -modified minimum essential medium; BK channel, large-conductance calcium-activated potassium channel; BrdU, 5-bromo-2'-deoxyuridine; CCK-8, cell counting kit-8; CEC, chloroethylclonidine; CREB, cAMP response element binding protein; ERG channel, ether-a-go-go-related gene channel; GIRK, G-protein-gated inward rectifying potassium channel; MC3T3-E1, mouse calvaria-derived osteoblastic cells; PDL, population doubling level; PI-PLC, phosphoinositide-PLC; PTX, *Pertussis* toxin; RANKL, receptor activator of NF- $\kappa$ B ligand; SaM-1, human periosteum-derived osteoblastic cells; TEA, tetraethylammonium; TREK-1, TWIK-related potassium channel 1; WST, water-soluble tetrazolium

## Introduction

Bones are continuously resorbed by osteoclasts and formed by osteoblasts, and bone mass is maintained by the balance between their functions. In recent years, many studies have demonstrated that the sympathetic nervous system is involved in bone metabolism (Cherruau *et al.*, 1999; Eleftheriou *et al.*, 2005; Togari and Arai, 2008; He *et al.*, 2011). Bone loss can be induced by continuously high sympathetic tone and this is reversed by  $\beta$ -adrenoceptor blockade (Bonnet *et al.*, 2008; Sato *et al.*, 2010). Previous studies, including ours, showed that mRNAs of both  $\alpha$ - and  $\beta$ -adrenoceptors were expressed in human osteoblasts (Togari *et al.*, 1997; Togari, 2002; Huang *et al.*, 2009). Although many studies have suggested that up-regulation of osteoclastogenesis and osteoclastic activity via  $\beta$ -adrenoceptors enhanced bone resorption (Arai *et al.*, 2003; Eleftheriou *et al.*, 2005; Kondo and Togari, 2011), the physiological role of  $\alpha$ -adrenoceptors in bone metabolism has been less well studied. Adrenaline stimulated cell proliferation and differentiation via  $\alpha_1$ -adrenoceptors in MC3T3-E1 osteoblast-like cells (Suzuki *et al.*, 1998; 2001). The expression levels of osteoprotegerin and receptor activator of NF- $\kappa$ B ligand (RANKL) were increased by  $\alpha$ - and  $\beta$ -adrenoceptor agonists (Takeuchi *et al.*, 2001; Nishiura and Abe, 2007; Huang *et al.*, 2009; receptor nomenclature follows Alexander *et al.*, (2011).

In general, GPCRs exert their functions via the phosphoinositide–phospholipase C (PI–PLC) pathway or the cAMP/PKA pathway, but recent studies suggested that ion channels also play important roles downstream of the signal transduction pathway of several GPCRs, including adrenoceptors (Inoue *et al.*, 2001; Marx *et al.*, 2002; Kim *et al.*, 2005; O-uchi *et al.*, 2008). We previously demonstrated that noradrenaline suppressed Cs-sensitive and tetraethylammonium-insensitive (TEA-insensitive) potassium channels via  $\alpha_{1B}$ -adrenoceptors in human osteoblasts, SaM-1 cells, by using whole-cell patch clamp recordings (Kodama and Togari, 2010). Several types of potassium channel had been found to regulate membrane potential and cell proliferation (Pardo *et al.*, 2005; Liebau *et al.*, 2006; Henney *et al.*, 2009; Jang *et al.*, 2011). In osteoblasts, blockade of large conductance Ca-activated potassium (BK) channels, ERG channels and TREK-1 channels promoted cell proliferation (Hughes *et al.*, 2006; Hernandez *et al.*, 2007), while a high dose of TEA, a potassium channel inhibitor, impaired cell proliferation (Hernandez *et al.*, 2007; Henney *et al.*, 2009). The physiological roles of the inhibitory effects of noradrenaline on potassium channels in human osteoblasts have not yet been defined.

In the present study, we undertook more detailed investigation of the signal transduction pathway in terms of the effect of noradrenaline and whether it is involved in cell proliferation in human osteoblasts, SaM-1 cells. Pharmacological analysis suggested that both G<sub>q</sub>- and G<sub>i/o</sub>-protein-coupled  $\alpha_{1B}$ -adrenoceptors were expressed in SaM-1 cells, and the latter was involved in the inhibitory effect of noradrenaline on potassium channels. Additionally, G<sub>i/o</sub>-coupled  $\alpha_1$ -adrenoceptors, not those coupled to G<sub>q</sub>-proteins, participated in noradrenaline-induced enhancement of cell proliferation. These results suggested that noradrenaline regulates cell proliferation via G<sub>i/o</sub>-protein-coupled

$\alpha_{1B}$ -adrenoceptors by regulating potassium channel activity in human osteoblasts.

## Methods

### Cell culture

The human osteoblasts used in this study, SaM-1 cells, were provided by Dr Koshihara, who prepared them with informed consent from an explant of ulnar periosteum tissue from a 20-year-old male patient who underwent curative surgery (Koshihara *et al.*, 1987). These cells have a mitotic lifespan of 34 population doubling levels (PDLs), and we used them at PDL of 22–24 for our experiments. We confirmed the cells are capable of calcifying at this level (Komoto *et al.*, 2012). The cells were cultured in  $\alpha$ -modified minimum essential medium ( $\alpha$ -MEM; Invitrogen, Carlsbad, CA) containing 10% FBS (Moregate Biotech, Bulimba, Australia) and 60  $\mu$ g·mL<sup>-1</sup> kanamycin at 37°C in 95% humidified air containing 5% CO<sub>2</sub>. The growth media were renewed every 2 days. For optical measurements of intracellular calcium concentration ([Ca<sup>2+</sup>]<sub>i</sub>) and whole-cell patch clamp recording, they were seeded on a cover slip, 1–2 days before the experiments.

### Optical measurements of [Ca<sup>2+</sup>]<sub>i</sub>

SaM-1 cells were loaded with fluo-3AM (2.5  $\mu$ M) for 30 min and washed three times with extracellular solution, which contained 124 mM NaCl, 3 mM KCl, 1 mM MgCl<sub>2</sub>, 2 mM CaCl<sub>2</sub>, 14 mM D-glucose and 10 mM HEPES (pH adjusted to 7.4 with NaOH), just before use. Then, the glass coverslip was transferred to a superfusion chamber on the stage of a confocal laser scanning microscope (LSM710, Carl Zeiss, Jena, Germany). Cells were superfused with extracellular solution at a rate of 2 mL·min<sup>-1</sup>. The fluorescence was recorded every 2 s at room temperature at an excitation wavelength of 488 nm, and the data were analysed using ZEN 2009 software (Carl Zeiss). Stock solutions of drugs were prepared and diluted 1000-fold into extracellular solution just before use, and they were applied by bath perfusion.

### Whole-cell patch clamp recording

SaM-1 cells were seeded on a glass coverslip. After 1–2 days of culture, the glass coverslip was transferred to a superfusion chamber on the stage of an inverted microscope (Axiovert200, Carl Zeiss). Cells were superfused with extracellular solution at 1–1.5 mL·min<sup>-1</sup>. Whole-cell voltage clamping was performed using patch electrodes (3–5 M $\Omega$ ) filled with internal solution containing 140 mM KCl, 2 mM MgCl<sub>2</sub>, 10 mM HEPES, 0.1 mM EGTA, 3 mM Na<sub>2</sub>ATP and 0.3 mM Na<sub>3</sub>GTP (pH adjusted to 7.4 with KOH) and Axopatch200B patch clamp amplifier (Axon Instruments, Sunnyvale, CA). In the experiments to investigate the involvement of [Ca<sup>2+</sup>]<sub>i</sub>, we used internal solution containing 130 mM KCl, 2 mM MgCl<sub>2</sub>, 10 mM HEPES, 5 mM EGTA, 3 mM Na<sub>2</sub>ATP and 0.3 mM Na<sub>3</sub>GTP (pH adjusted to 7.4 with KOH).

The holding potential was –60 mV, and voltage–current relationships were recorded using voltage steps or ramps. In the voltage step protocol, 10 mV voltage steps from –100 to +40 mV from a holding potential of –60 mV for 500 ms were applied. In the voltage ramp protocol, 1 s voltage ramp from

–100 to +40 mV following 150 ms voltage steps of –100 mV was applied every 10 s. The signals were low-pass filtered at 10 kHz and digitized at 20 kHz for analysis with pClamp9.2 software (Molecular Devices, Silicon Valley, CA). Stock solutions of drugs were prepared and diluted 1000-fold into extracellular solution just before use, and they were applied by bath perfusion.

### Cell proliferation assay

**5-Bromo-2'-deoxyuridine (BrdU) incorporation.** Cell proliferation activity was assessed by BrdU incorporation using Cell Proliferation ELISA, BrdU kit (Roche Applied Science, Indianapolis, IN) according to the manufacturer's instructions. Cells were seeded in a 96-well plate at a density of 10 000 cells-per well in 100  $\mu$ L of culture medium. After 1 day, culture media were replaced with  $\alpha$ -MEM containing 2.5% FBS and kanamycin, and then the cells were treated with noradrenaline for 20 h at 37°C. BrdU labelling solution (100  $\mu$ M) was added at 10  $\mu$ L-per well, and the cells were incubated for an additional 4 h at 37°C. After this, the labelling media were removed, the cells were fixed and DNA was denatured with FixDenat solution. The cells were incubated with peroxidase-conjugated anti-BrdU antibody for 1.5 h at room temperature. The cells were then washed three times with PBS, followed by the addition of substrate solution (tetraethyl-benzidine) at 100  $\mu$ L-per well. After 15 min incubation, 1 M  $\text{H}_2\text{SO}_4$  was added at 25  $\mu$ L-per well to stop the peroxidase reaction, and then absorbance of wells was measured at 450 nm using Multiskan FC (Thermo Fisher Scientific, Waltham, MA).

**Water-soluble tetrazolium (WST) assay.** Cell number was assessed as dehydrogenase activity by using Cell Counting Kit-8 (CCK-8, Dojindo, Kumamoto, Japan). In this assay, WST-8 is reduced to water-soluble formazan outside the cells. Cells were seeded in a 96-well plate at a density of 10 000 cells-per well in 100  $\mu$ L of culture medium. After 1 day, culture media were replaced with  $\alpha$ -MEM containing 2.5% FBS and kanamycin, and then the cells were treated with noradrenaline for 22.5 h at 37°C. CCK-8 solution was added at 10  $\mu$ L-per well, and the cells were incubated for an additional 1.5 h at 37°C. Then absorbance of wells was measured at 450 nm using Multiskan FC (Thermo Fisher Scientific).

To confirm that the experimental conditions were adequate, the data from non-drug-treated cultures were also recorded in each experiment.

### Statistical analysis

All data are expressed as mean  $\pm$  SEM. In the voltage step protocol, the changes of whole-cell currents in a stable state were evaluated using the paired *t*-test. In the voltage ramp protocol, the effects of the drugs on whole-cell currents were evaluated by comparing average currents (six traces for 1 min) taken during the peak response to each drug with those before drug application. Cell proliferation activities were evaluated by comparing with the data from a control group. The two-tailed *t*-test combined with Bonferroni's correction following one-way ANOVA was used for multiple comparisons. Differences with *P*-values <0.05 were considered significant.

### Materials

L-Noradrenaline; chloroethylclonidine (CEC), an  $\alpha_{1B}$ -adrenoceptor-selective antagonist; U73122, a PLC inhibitor; H89, a PKA inhibitor; clonidine, prazosin and propranolol were purchased from Sigma Aldrich (St. Louis, MO, USA). Pertussis toxin (PTX) was purchased from Merck KGaA (Darmstadt, Germany). CsCl was purchased from Nacalai Tesque (Kyoto, Japan). A  $\text{G}\beta\gamma$ -protein inhibitor, gallein, was purchased from Tocris Biosciences (Bristol, UK). A calcium fluorophore, fluo-3AM, was purchased from Dojindo. U73122, gallein and H89 were dissolved in dimethyl sulfoxide. All other chemicals used were of reagent grade.


### Results

#### *Involvement of the PI-PLC pathway in the effects of noradrenaline*

In  $\text{Ca}^{2+}$  fluorescence imaging, elevation of fluo-3AM fluorescence intensity was induced by bath application of 1  $\mu$ M noradrenaline in  $50.4 \pm 6.4\%$  of cells examined and the responses were reproducible on the same cells with repeated application in SaM-1 cells (13 individual experiments; Figure 1A). In the cells pretreated with 100  $\mu$ M CEC for 45 min at 37°C, bath-applied noradrenaline had no effects on fluorescence (five individual experiments, data not shown). Additionally, the effect of noradrenaline on  $[\text{Ca}^{2+}]_i$  was eliminated by pretreatment with the PLC inhibitor U73122 for 10 min (five individual experiments; Figure 1B). On the other hand, in whole-cell patch clamp recording, the currents induced by voltage steps were significantly reduced and reversal potential was shifted rightwards by 1  $\mu$ M bath-applied noradrenaline, as shown in our previous study ( $n = 6$ ; Figure 1C; Kodama and Togari, 2010). The ratios between the current amplitude at the beginning (50 ms) of the pulse and that at the end of pulse (500 ms) were  $95.1 \pm 2.5\%$  and  $90.1 \pm 4.4\%$  in the absence and presence of noradrenaline respectively. There was no apparent effect of noradrenaline on the current kinetics. Similarly, in the voltage ramp protocol, whole-cell current was reduced, at 40 mV ( $n = 7$ ; Figure 1D). These inhibitory effects of noradrenaline on whole-cell current were shown in almost all cells tested (48 of 50 independent experiments, including preliminary data). Bath-applied U73122 slightly suppressed whole-cell current ( $n = 5$ ), and treatment with noradrenaline following U73122 exhibited a similar extent of inhibition as in the control cells ( $n = 5$ ; Figure 1D, E). Additionally, the inhibitory effect of noradrenaline was unaffected by using the internal solution containing 5 mM EGTA ( $n = 5$ ; Figure 1D).

#### *Involvement of $\text{G}_{i/o}$ - and $\text{G}\beta\gamma$ -protein in the effects of noradrenaline on whole-cell current*

We next examined whether  $\text{G}_{i/o}$ -protein is involved in the inhibitory effect of noradrenaline on whole-cell current. In the cells pretreated with  $\text{G}_{i/o}$ -coupled receptor inhibitor, PTX, at 150 ng·mL $^{-1}$  for 24 h, noradrenaline-induced suppression of whole-cell current was significantly attenuated ( $n = 11$ , 5, respectively; Figure 2A, D). Treatment with PTX inhibits not only the effect via  $\text{G}_i$ -protein but also that via  $\text{G}\beta\gamma$ -protein



**Figure 1**

Involvement of the G<sub>q</sub>/PI-PLC pathway in the inhibitory effect of noradrenaline (NA) on whole-cell current in human osteoblasts. (A and B) Representative traces of  $[Ca^{2+}]_i$  elevation induced by repeated application of noradrenaline in SaM-1 cells. (A) The representative recordings from cells that responded to noradrenaline (B) Representative responses to noradrenaline in the absence (the first application) and presence (the second application) of U73122, an inhibitor of PLC. (C) The averaged voltage-current density relationship recorded by voltage step from -100 to +40 mV. (D) Summary of the influence of the presence of EGTA, a Ca chelator, at 5 mM in the pipette solution and pretreatment with U73122, on the effects of noradrenaline (control,  $n = 7$ ; EGTA, U73122,  $n = 5$ ). (E) Representative averaged traces of six consecutive whole-cell currents during a -100 to +40 mV voltage ramp (a) and time course of whole-cell currents at +40 mV (b). The bars indicate the time points when the averaged traces were recorded. Values are shown as the mean  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$  compared with non-treated control. Each figure is representative data of five independent experiments in panels A, B and E.


(Katz *et al.*, 1992). Then we examined the effect of G $\beta\gamma$ -protein inhibitor, gallein and PKA inhibitor, H89. Pretreatment with gallein at 10  $\mu$ M for 30 min just before the experiment significantly reduced the inhibitory effect of noradrenaline ( $n = 5$ ; Figure 2B, D). Meanwhile, bath application of H89 at 2.5  $\mu$ M slightly suppressed whole-cell current ( $n = 5$ ), and the inhibitory effect of noradrenaline following H89 application tended to be attenuated ( $n = 5$ ; Figure 2C, D).

In our previous study,  $\alpha_{2B}$ -adrenoceptors were expressed at a low level in SaM-1 cells (Togari *et al.*, 1997; Togari, 2002). In

general,  $\alpha_2$ -adrenoceptors are known to be coupled with G<sub>i/o</sub>-protein. We then examined the effect of the  $\alpha_2$ -adrenoceptor agonist, clonidine, on whole-cell current. Bath application of clonidine had no obvious effect on the voltage-current relationship ( $n = 5$ ; Figure 3).

### Effects of adrenoceptor ligands on proliferation in SaM-1 cells


Cell proliferation activity was evaluated as DNA synthesis by BrdU test, and live cell number was evaluated as



**Figure 2**

Involvement of  $G_{i/o}$ - and  $G\beta\gamma$ -proteins in the inhibitory effects of noradrenaline (NA) on whole-cell current. (A–C) Representative averaged traces of six consecutive whole-cell currents during a  $-100$  to  $+40$  mV voltage ramp (a) and time course of whole-cell currents recorded at  $+40$  mV (b) in the cells pretreated with *Pertussis* toxin (PTX), a  $G_{i/o}$ -protein-coupled receptor inhibitor (A), gallein, a  $G\beta\gamma$ -protein inhibitor (B), or H89, a PKA inhibitor (C). The bars indicate the time points when the averaged traces were recorded. (D) Summary results of the pretreatment with PTX, gallein or H89 on the effects of noradrenaline (control,  $n = 11$ ; PTX, gallein, H89,  $n = 5$ ). Values are shown as the mean  $\pm$  SEM.  $**P < 0.01$  compared with control. Each figure shows representative data of five independent experiments in panels A, B and C.





**Figure 3**

The effects of clonidine, an  $\alpha_2$ -adrenoceptor agonist, on whole-cell current in human osteoblasts. (A and B) Representative averaged traces of six consecutive whole-cell currents during a  $-100$  to  $+40$  mV voltage ramp (A) and time course of whole-cell currents recorded at  $+40$  mV (B). These are representative data from five independent experiments. The bars indicate the time points when the averaged traces were recorded.

dehydrogenase activity by WST assay. In SaM-1 cells, treatment with noradrenaline for 24 h increased BrdU incorporation at a lower dose ( $0.03 - 0.3 \mu\text{M}$ ) and suppressed it at higher doses ( $1 \mu\text{M} - 30 \mu\text{M}$ ;  $n = 8$ ; Figure 4A). On the other hand, noradrenaline only increased formazan formation in WST assays, over the whole concentration range ( $n = 8$ ; Figure 4D). In the presence of prazosin, an  $\alpha_1$ -adrenoceptor antagonist, at  $30 \mu\text{M}$ , noradrenaline induced only decreased effects in both the BrdU test and the WST assay ( $n = 8$ ; Figure 4B, E). In contrast, in the presence of a  $\beta$ -adrenoceptor antagonist, propranolol, at  $10 \mu\text{M}$ , noradrenaline showed only positive effects in both assays ( $n = 8$ ; Figure 4C, F). These results suggested that cell proliferation was facilitated via  $\alpha_1$ -adrenoceptors and inhibited via  $\beta$ -adrenoceptors.

### Signal pathways involved in the effects of noradrenaline on cell proliferation


Previously, we demonstrated that Cs-sensitive potassium channels were involved in the inhibitory effects of noradrenaline on whole-cell current (Kodama and Togari, 2010). Here, we have examined whether CsCl affected the increasing effects of noradrenaline via  $\alpha_1$ -adrenoceptors. In the presence of  $1 \text{ mM}$  CsCl and  $10 \mu\text{M}$  propranolol, noradrenaline had no effect on BrdU incorporation and formazan formation ( $n = 8$ ; Figure 5).

Next, to study whether signal pathways were involved in the positive effects on cell proliferation of noradrenaline, we evaluated BrdU incorporation and formazan formation in the presence of propranolol and gallein, H89, PTX or U73122. The positive effects of noradrenaline in both assays were eliminated by pretreatment with PTX for 24 h ( $n = 8$ ; Figure 5). Likewise, in the presence of gallein or H89, noradrenaline had no effect in both assays ( $n = 8$ ; Figure 5). In contrast, in the cells treated with U73122, the positive effect via  $\alpha_1$ -adrenoceptors was shown in both assays ( $n = 8$ ; Figure 5). These results suggested that regulation of potassium current via G<sub>i/o</sub>-coupled  $\alpha_1$ -adrenoceptors was involved in the positive effects on cell proliferation.

### Discussion

Recent studies have demonstrated that a single subtype of receptor can be associated with different types of heterotrimeric G-proteins (Wenzel-Seifert and Seifert, 2000; Gazi *et al.*, 2003; Cordeaux *et al.*, 2004). It is also suggested that coupling with a particular type of G-protein can be regulated by the location, phosphorylation and expression of GPCRs (Davies *et al.*, 1999; Xiang *et al.*, 2002; Hasseldine *et al.*, 2003). The results presented in this study suggest that  $\alpha_{1B}$ -adrenoceptors can be coupled to both G<sub>q</sub>-protein and G<sub>i/o</sub>-protein in human osteoblasts. We previously reported that mRNAs of  $\alpha_{1B}$ -,  $\alpha_{2B}$ - and  $\beta_2$ -adrenoceptors were expressed in SaM-1 cells (Togari *et al.*, 1997; Togari, 2002). Bath application of noradrenaline induced  $[\text{Ca}^{2+}]_i$  elevation, and the effect was inhibited by the  $\alpha_{1B}$ -adrenoceptor-selective inhibitor, CEC, and the PLC inhibitor, U73122 (Figure 1A, B). These results indicated that noradrenaline activated the PI-PLC pathway via G<sub>q</sub>-protein-coupled  $\alpha_{1B}$ -adrenoceptors in SaM-1 cells. Meanwhile, noradrenaline-induced suppression of whole-cell current, which was inhibited by pretreatment with CEC in our previous study (Kodama and Togari, 2010), was affected by neither addition of  $\text{Ca}^{2+}$  chelator to internal solution nor treatment with U73122 (Figure 1D, E). In contrast, the inhibitory effect of noradrenaline on whole-cell current was significantly attenuated by pretreatment with PTX (Figure 2A). However, the  $\alpha_2$ -adrenoceptor agonist, clonidine, had no effect on the whole-cell current (Figure 3). These results indicated that noradrenaline inhibited whole-cell current via G<sub>i/o</sub>-protein-coupled  $\alpha_{1B}$ -adrenoceptors. Although noradrenaline-induced  $[\text{Ca}^{2+}]_i$  elevations were detected in about half of SaM-1 cells, the inhibitory effect of noradrenaline on whole-cell current appeared in almost all cells tested. This suggested the possibility that  $\alpha_{1B}$ -adrenoceptors were mainly, but not all, coupled to G<sub>i/o</sub>-proteins in human osteoblasts.

The effects of G<sub>i/o</sub>-protein-coupled GPCRs are exerted not only through the G $\alpha$  subunit but also through G $\beta\gamma$  subunits. Many studies reported that G-protein-gated inward rectifying




**Figure 4**

The effects of noradrenaline (NA) on cell proliferation of human osteoblasts through  $\alpha_1$ - and  $\beta$ -adrenoceptors. (A–C) Effects of noradrenaline on DNA synthesis measured by BrdU incorporation. (A) Dose–response relationship of the effect of noradrenaline ( $n = 8$ ). (B, C) Effects of noradrenaline in the presence of the  $\alpha_1$ -blocker prazosin at 30  $\mu$ M (B) and  $\beta$ -blocker propranolol at 10  $\mu$ M (C) ( $n = 8$ ). (D–F) Effects of noradrenaline on water-soluble tetrazolium salt (formazan) formation by dehydrogenase in live cells were assessed by WST assay. (D) Dose–response relationship of the effect of noradrenaline ( $n = 8$ ). (E, F) Effects of noradrenaline in the presence of prazosin at 30  $\mu$ M (E) and propranolol at 10  $\mu$ M (F) ( $n = 8$ ). Values are shown as the mean  $\pm$  SEM. \* $P$  < 0.05, \*\* $P$  < 0.01 compared with control. Each figure is representative data from three to five independent experiments.

potassium channels (GIRK channels) are regulated by G $\beta\gamma$  subunits (Filippov *et al.*, 2004; Lüscher and Slesinger, 2010). In addition, recent studies reported that G $\beta\gamma$  subunits could interact with several proteins, for example, L-type calcium channels, connexin 43 and GPCR kinase (O-uchi *et al.*, 2008; Sato *et al.*, 2009; Casey *et al.*, 2010). In this study, the inhibitory effect of noradrenaline on whole-cell current was significantly attenuated by pretreatment with the G $\beta\gamma$ -protein inhibitor, gallein (Figure 2B). The inhibitory effect of noradrenaline also tended to be attenuated following application of a PKA inhibitor, H89 (Figure 2C). In SaM-1 cells, the reversal potential of whole-cell current was considered to be less negative. This suggested the possibility that not only potassium channels but also chloride channels, contributed to the whole-cell current. However, noradrenaline-induced suppression was eliminated in the presence of CsCl, a non-selective potassium channel blocker, in our previous study (Kodama and Togari, 2010). Therefore, these results suggested that the potassium channels were inhibited by G $\beta\gamma$ -proteins and were partly regulated by PKA. Although the GIRK channels are a major target of G $\beta\gamma$ -proteins, the whole-cell current suppressed by noradrenaline did not show inward rectification. Zhou *et al.*, (2008) demonstrated that BK channels were

inhibited by G $\beta\gamma$ -protein and we previously reported that BK channels were expressed in SaM-1 cells (Hirukawa *et al.*, 2008). However, BK channels are inhibited by TEA and activated by [Ca<sup>2+</sup>]<sub>i</sub> elevation (Henney *et al.*, 2009; Lee and Cui, 2010). The inhibition of the current by noradrenaline was seen at cell potentials below –60 mV, and further inhibition was seen over 0 mV (Figure 1C). Additionally, there was no apparent effect of noradrenaline on the current kinetics. These results suggested that the channels inhibited by noradrenaline were non-inactivated, outward rectifying and open at the resting potential. Recent studies demonstrated that the two-pore domain K channel family is responsible for non-inactivated, leak or background, potassium current (Patel and Honoré, 2001) and TREK-1, a mechanosensitive member of the two-pore domain K channels, was expressed and regulated cell proliferation in human osteoblasts (Hughes *et al.*, 2006). The current through TREK-1 is outwardly rectifying and is insensitive to classical potassium channel blockers, including TEA (Patel and Honoré, 2001). However, TREK-1 activity was inhibited by G<sub>s</sub>-protein-coupled receptors and stimulated by G<sub>i/o</sub>-protein-coupled receptors (Mathie, 2007). Rivard *et al.* (2009) reported that overexpression of  $\alpha_{1B}$ -adrenoceptors decreased outwards potassium



**Figure 5**

Involvement of noradrenaline-induced potassium channel inhibition on its effect on cell proliferation. The effects of noradrenaline (NA) via  $\alpha_1$ -adrenoceptor on cell proliferation were evaluated as DNA synthesis by BrdU test (A) and as dehydrogenase activity by WST assay (B) in the presence of propranolol and CsCl, gallein, H89 or U73122 (BrdU test: control, CsCl;  $n = 8$ ; gallein, H89, U73122;  $n = 14$ ; WST assay:  $n = 8$ ). Open square indicates the data from cells pretreated with PTX for 24 h before noradrenaline application ( $n = 8$ ). Values are shown as the mean  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$  compared with each non-NA-treated group. Each figure shows representative data from three independent experiments.

current and several types of potassium channel were involved in its effects in heart. Further investigations are needed to identify the potassium channels suppressed by noradrenaline in SaM-1 cells.

A number of studies have demonstrated that membrane potential and cell proliferation capacity were regulated by activities of potassium channels in several types of cells including osteoblasts (Pardo *et al.*, 2005; Liebau *et al.*, 2006; Henney *et al.*, 2009; Jang *et al.*, 2011). Adequate inhibition of potassium channels is thought to cause membrane depolarization and activate voltage-dependent calcium channels, leading to cell proliferation (Burg *et al.*, 2008). Additionally, activation of potassium channels plays an important role at an early stage of apoptosis (Burg *et al.*, 2008; Valencia-Cruz *et al.*, 2009). In the present study, noradrenaline increased DNA synthesis at a low dose and decreased it at a high dose (Figure 4A). In WST assays, noradrenaline also increased formazan formation in a dose-dependent manner (Figure 4D). While the results of the BrdU tests indicated DNA synthesis

during 20–24 h after treatment with noradrenaline, those from the WST assays indicated cumulative effects on cell proliferation. These differences might account for the divergent results of the BrdU test and the WST assay, at high concentrations of noradrenaline. In the BrdU test, the positive and negative effects of noradrenaline were inhibited by prazosin and propranolol respectively (Figure 4B, C). Additionally, in the WST assay, positive effects of noradrenaline were inhibited by prazosin but not by propranolol (Figure 4E, F). These results suggested that cell proliferation in human osteoblasts is bi-directionally regulated by noradrenaline via  $\alpha_1$ - and  $\beta$ -adrenoceptors. This is consistent with a previous study in human osteoblasts (Huang *et al.*, 2009). A number of studies have demonstrated involvement of  $\beta_2$ -adrenoceptors in bone metabolism. In general, the  $\beta_2$ -adrenoceptor is coupled with G<sub>s</sub>-protein and is thought to activate the cAMP/PKA pathway and Fu *et al.*, (2005) demonstrated that  $\beta_2$ -adrenoceptors regulated osteoblast proliferation via the cAMP/PKA/CREB signalling pathway *in vivo*.

As observed with whole-cell current, the positive effects of noradrenaline via  $\alpha_1$ -adrenoceptors, shown in BrdU tests and WST assays, was inhibited in the presence of CsCl (Figure 5). This suggested that noradrenaline enhanced cell proliferation by blocking Cs-sensitive potassium channels. Additionally, the positive effects of noradrenaline via  $\alpha_1$ -adrenoceptors on cell proliferation in both assays were suppressed by inhibition of G<sub>i/o</sub>-protein-coupled receptors by PTX, G $\beta\gamma$ -protein by gallein or the cAMP/PKA pathway by H89, but not by inhibition of the PI-PLC pathway by U73122 (Figure 5). These results suggested the possibility that noradrenaline-induced suppressions of potassium channels via G $\beta\gamma$ -protein and the cAMP–PKA pathway were involved in its effect on cell proliferation mediated by  $\alpha_{1B}$ -adrenoceptors. Suzuki *et al.* (1998; 2001) reported that adrenaline stimulated cell proliferation and alkaline phosphatase activity and increased type III P<sub>i</sub> transporter expression via  $\alpha_1$ -adrenoceptors, and these effects were inhibited by PTX in MC3T3-E1 osteoblast-like cells. Activation of  $\alpha_1$ -adrenoceptors induced the expression of RANKL, and this effect was inhibited by a PKC inhibitor (Nishiura and Abe, 2007). Therefore, G<sub>q</sub>- and G<sub>i/o</sub>-coupled  $\alpha_1$ -adrenoceptors are thought to play different roles in osteoblasts.

Both experimental and clinical studies have demonstrated that  $\beta$ -adrenoceptor blockade is effective against osteoporosis accompanied by a highly active sympathetic nervous system (Pasco *et al.*, 2004; Bonnet *et al.*, 2007; 2008; Sato *et al.*, 2010; Yang *et al.*, 2011). There is also a report that  $\alpha_1$ -adrenoceptor blockade increased the risk of hip/femur fracture (Souverein *et al.*, 2003). Phenylephrine, an  $\alpha_1$ -adrenoceptor agonist, promotes osteogenesis in bone fracture healing (McDonald *et al.*, 2011). All these reports suggest the possibility that activation of  $\alpha_1$ -adrenoceptors could lead to a treatment for osteoporosis. The positive effects via  $\alpha_{1B}$ -adrenoceptors on osteoblast proliferation might contribute to bone fracture healing and treatment of osteoporosis. However, neither knockout nor overexpression of  $\alpha_{1B}$ -adrenoceptors has been reported to result in obvious skeletal phenotypes (Battaglia *et al.*, 2003; Rivard *et al.*, 2009; Docherty, 2010). Detailed analysis of bone metabolism in a pathophysiological state showing high sympathetic tone, for example, in leptin-injected mice, ovariectomized mice or



spontaneous hypertensive rats, is required to investigate the physiological role of  $\alpha_{1B}$ -adrenoceptor (Eleftheriou *et al.*, 2005; Sato *et al.*, 2010).

In conclusion, the present study suggested that noradrenaline suppressed potassium channels via  $G_{i/o}$ -coupled  $\alpha_{1B}$ -adrenoceptors and that the  $G\beta\gamma$ -proteins play an important role in this effect. Additionally, this effect was involved in the noradrenaline-induced increase in proliferation of human osteoblasts, SaM-1 cells. These results will aid understanding of the mechanism behind the regulation of bone metabolism by the sympathetic nervous system.

## Acknowledgements

This work was partly supported by a grant-in-aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (No. 21890276 to D.K.) and by a Grant-in-Aid from Strategic Research AGU-Platform Formation (2008-2012).

## Conflicts of interest

All authors have no conflicts of interest.

## References

- Alexander SPH, Mathie A, Peters JA (2011). Guide to Receptors and Channels (GRAC), 5th edition. Br J Pharmacol 164 (Suppl. 1): S1–S324.
- Arai M, Nagasawa T, Koshihara Y, Yamamoto S, Togari A (2003). Effects of beta-adrenergic agonists on bone-resorbing activity in human osteoclast-like cells. Biochim Biophys Acta 1640: 137–142.
- Battaglia G, Fornai F, Busceti CL, Lembo G, Nicoletti F, De Blasi A (2003). Alpha-1B adrenergic receptor knockout mice are protected against methamphetamine toxicity. J Neurochem 86: 413–421.
- Bonnet N, Gadois C, McCloskey E, Lemineur G, Lespessailles E, Courteix D *et al.* (2007). Protective effect of beta blockers in postmenopausal women: influence on fractures, bone density, micro and macroarchitecture. Bone 40: 1209–1216.
- Bonnet N, Benhamou CL, Malaval L, Goncalves C, Vico L, Eder V *et al.* (2008). Low dose beta-blocker prevents ovariectomy-induced bone loss in rats without affecting heart functions. J Cell Physiol 217: 819–827.
- Burg ED, Remillard CV, Yuan JX (2008). Potassium channels in the regulation of pulmonary artery smooth muscle cell proliferation and apoptosis: pharmacotherapeutic implications. Br J Pharmacol 153 (Suppl. 1): S99–S111.
- Casey LM, Pistner AR, Belmonte SL, Migdalovich D, Stolpnik O, Nwakanma FE *et al.* (2010). Small molecule disruption of G beta gamma signaling inhibits the progression of heart failure. Circ Res 107: 532–539.
- Cherruau M, Facchinetti P, Baroukh B, Saffar JL (1999). Chemical sympathectomy impairs bone resorption in rats: a role for the sympathetic system on bone metabolism. Bone 25: 545–551.
- Cordeaux Y, Ijzerman AP, Hill SJ (2004). Coupling of the human A1 adenosine receptor to different heterotrimeric G proteins: evidence for agonist-specific G protein activation. Br J Pharmacol 143: 705–714.
- Davies MG, Huynh TT, Hagen PO (1999). Functional characterization of alpha1-adrenergic receptors in experimental vein grafts. J Surg Res 84: 40–45.
- Docherty JR (2010). Subtypes of functional alpha1-adrenoceptor. Cell Mol Life Sci 67: 405–417.
- Eleftheriou F, Ahn JD, Takeda S, Starbuck M, Yang X, Liu X *et al.* (2005). Leptin regulation of bone resorption by the sympathetic nervous system and CART. Nature 434: 514–520.
- Filippov AK, Fernández-Fernández JM, Marsh SJ, Simon J, Barnard EA, Brown DA (2004). Activation and inhibition of neuronal G protein-gated inwardly rectifying K(+) channels by P2Y nucleotide receptors. Mol Pharmacol 66: 468–477.
- Fu L, Patel MS, Bradley A, Wagner EF, Karsenty G (2005). The molecular clock mediates leptin-regulated bone formation. Cell 122: 803–815.
- Gazi L, Nickolls SA, Strange PG (2003). Functional coupling of the human dopamine D2 receptor with G alpha i1, G alpha i2, G alpha i3 and G alpha o G proteins: evidence for agonist regulation of G protein selectivity. Br J Pharmacol 138: 775–786.
- Hasseldine AR, Harper EA, Black JW (2003). Cardiac-specific overexpression of human beta2 adrenoceptors in mice exposes coupling to both Gs and Gi proteins. Br J Pharmacol 138: 1358–1366.
- He JY, Jiang LS, Dai LY (2011). The roles of the sympathetic nervous system in osteoporotic diseases: a review of experimental and clinical studies. Ageing Res Rev 10: 253–263.
- Henney NC, Li B, Elford C, Reviriego P, Campbell AK, Wann KT *et al.* (2009). A large-conductance (BK) potassium channel subtype affects both growth and mineralization of human osteoblasts. Am J Physiol Cell Physiol 297: C1397–C1408.
- Hernandez L, Park KH, Cai SQ, Qin L, Partridge N, Sesti F (2007). The antiproliferative role of ERG K+ channels in rat osteoblastic cells. Cell Biochem Biophys 47: 199–208.
- Hirukawa K, Muraki K, Ohya S, Imaizumi Y, Togari A (2008). Electrophysiological properties of a novel Ca(2+)-activated K(+) channel expressed in human osteoblasts. Calcif Tissue Int 83: 222–229.
- Huang HH, Brennan TC, Muir MM, Mason RS (2009). Functional alpha1- and beta2-adrenergic receptors in human osteoblasts. J Cell Physiol 220: 267–275.
- Hughes S, Magnay J, Foreman M, Publicover SJ, Dobson JP, El Haj AJ (2006). Expression of the mechanosensitive 2PK+ channel TREK-1 in human osteoblasts. J Cell Physiol 206: 738–748.
- Inoue R, Okada T, Onoue H, Hara Y, Shimizu S, Naitoh S *et al.* (2001). The transient receptor potential protein homologue TRP6 is the essential component of vascular alpha(1)-adrenoceptor-activated Ca(2+)-permeable cation channel. Circ Res 88: 325–332.
- Jang SS, Park J, Hur SW, Hong YH, Hur J, Chae JH *et al.* (2011). Endothelial progenitor cells functionally express inward rectifier potassium channels. Am J Physiol Cell Physiol 301: C150–C161.
- Katz A, Wu D, Simon MI (1992). Subunits beta gamma of heterotrimeric G protein activate beta 2 isoform of phospholipase C. Nature 360: 686–689.
- Kim Y, Park MK, Uhm DY, Shin J, Chung S (2005). Modulation of delayed rectifier potassium channels by alpha1-adrenergic activation via protein kinase C zeta and p62 in PC12 cells. Neurosci Lett 387: 43–48.

- Kodama D, Togari A (2010). Modulation of potassium channels via the  $\alpha$ 1B-adrenergic receptor in human osteoblasts. *Neurosci Lett* 485: 102–106.
- Komoto S, Kondo H, Fukuta O, Togari A (2012). Comparison of  $\beta$ -adrenergic and glucocorticoid signaling on clock gene and osteoblast-related gene expressions in human osteoblast. *Chronobiol Int* 29: 66–74.
- Kondo H, Togari A (2011). Continuous treatment with a low-dose  $\beta$ -agonist reduces bone mass by increasing bone resorption without suppressing bone formation. *Calcif Tissue Int* 88: 23–32.
- Koshihara Y, Kawamura M, Oda H, Higaki S (1987). In vitro calcification in human osteoblastic cell line derived from periosteum. *Biochem Biophys Res Commun* 145: 651–657.
- Lee US, Cui J (2010). BK channel activation: structural and functional insights. *Trends Neurosci* 33: 415–423.
- Liebau S, Pröpper C, Böckers T, Lehmann-Horn F, Storch A, Grissmer S *et al.* (2006). Selective blockage of Kv1.3 and Kv3.1 channels increases neural progenitor cell proliferation. *J Neurochem* 99: 426–437.
- Lüscher C, Slesinger PA (2010). Emerging roles for G protein-gated inwardly rectifying potassium (GIRK) channels in health and disease. *Nat Rev Neurosci* 11: 301–315.
- McDonald SJ, Dooley PC, McDonald AC, Djouma E, Schuijers JA, Ward AR *et al.* (2011).  $\alpha$ (1) adrenergic receptor agonist, phenylephrine, actively contracts early rat rib fracture callus *ex vivo*. *J Orthop Res* 29: 740–745.
- Marx SO, Kurokawa J, Reiken S, Motoike H, D'Armiento J, Marks AR *et al.* (2002). Requirement of a macromolecular signaling complex for beta adrenergic receptor modulation of the KCNQ1-KCNE1 potassium channel. *Science* 295: 496–499.
- Mathie A (2007). Neuronal two-pore-domain potassium channels and their regulation by G protein-coupled receptors. *J Physiol* 578: 377–385.
- Nishiura T, Abe K (2007). Alpha1-adrenergic receptor stimulation induces the expression of receptor activator of nuclear factor kappaB ligand gene via protein kinase C and extracellular signal-regulated kinase pathways in MC3T3-E1 osteoblast-like cells. *Arch Oral Biol* 52: 778–785.
- O-Uchi J, Sasaki H, Morimoto S, Kusakari Y, Shinji H, Obata T *et al.* (2008). Interaction of alpha1-adrenoceptor subtypes with different G proteins induces opposite effects on cardiac L-type Ca<sup>2+</sup> channel. *Circ Res* 102: 1378–1388.
- Pardo LA, Contreras-Jurado C, Zientkowska M, Alves F, Stühmer W (2005). Role of voltage-gated potassium channels in cancer. *J Membr Biol* 205: 115–124.
- Pasco JA, Henry MJ, Sanders KM, Kotowicz MA, Seeman E, Nicholson GC (2004). Beta-adrenergic blockers reduce the risk of fracture partly by increasing bone mineral density: Geelong Osteoporosis Study. *J Bone Miner Res* 19: 19–24.
- Patel AJ, Honoré E (2001). Properties and modulation of mammalian 2P domain K<sup>+</sup> channels. *Trends Neurosci* 24: 339–346.
- Rivard K, Trépanier-Boulay V, Rindt H, Fiset C (2009). Electrical remodeling in a transgenic mouse model of alpha1B-adrenergic receptor overexpression. *Am J Physiol Heart Circ Physiol* 296: H704–H718.
- Sato M, Jiao Q, Honda T, Kurotani R, Toyota E, Okumura S *et al.* (2009). Activator of G protein signaling 8 (AGS8) is required for hypoxia-induced apoptosis of cardiomyocytes: role of G betagamma and connexin 43 (CX43). *J Biol Chem* 284: 31431–31440.
- Sato T, Arai M, Goto S, Togari A (2010). Effects of propranolol on bone metabolism in spontaneously hypertensive rats. *J Pharmacol Exp Ther* 334: 99–105.
- Souverain PC, Van Staa TP, Egberts AC, De la Rosette JJ, Cooper C, Leufkens HG (2003). Use of alpha-blockers and the risk of hip/femur fractures. *J Intern Med* 254: 548–554.
- Suzuki A, Palmer G, Bonjour JP, Caverzasio J (1998). Catecholamines stimulate the proliferation and alkaline phosphatase activity of MC3T3-E1 osteoblast-like cells. *Bone* 23: 197–203.
- Suzuki A, Palmer G, Bonjour JP, Caverzasio J (2001). Stimulation of sodium-dependent inorganic phosphate transport by activation of Gi/o-protein-coupled receptors by epinephrine in MC3T3-E1 osteoblast-like cells. *Bone* 28: 589–594.
- Takeuchi T, Tsuboi T, Arai M, Togari A (2001). Adrenergic stimulation of osteoclastogenesis mediated by expression of osteoclast differentiation factor in MC3T3-E1 osteoblast-like cells. *Biochem Pharmacol* 61: 579–586.
- Togari A (2002). Adrenergic regulation of bone metabolism: possible involvement of sympathetic innervation of osteoblastic and osteoclastic cells. *Microsc Res Tech* 58: 77–84.
- Togari A, Arai M (2008). Pharmacological topics of bone metabolism: the physiological function of the sympathetic nervous system in modulating bone resorption. *J Pharmacol Sci* 106: 542–546.
- Togari A, Arai M, Mizutani S, Mizutani S, Koshihara Y, Nagatsu T (1997). Expression of mRNAs for neuropeptide receptors and beta-adrenergic receptors in human osteoblasts and human osteogenic sarcoma cells. *Neurosci Lett* 233: 125–128.
- Valencia-Cruz G, Shabala L, Delgado-Enciso I, Shabala S, Bonales-Alatorre E, Pottosin II *et al.* (2009). K<sub>bg</sub> and Kv1.3 channels mediate potassium efflux in the early phase of apoptosis in Jurkat T lymphocytes. *Am J Physiol Cell Physiol* 297: C1544–C1533.
- Wenzel-Seifert K, Seifert R (2000). Molecular analysis of beta(2)-adrenoceptor coupling to G(s)-, G(i)-, and G(q)-proteins. *Mol Pharmacol* 58: 954–966.
- Xiang Y, Rybin VO, Steinberg SF, Kobilka B (2002). Caveolar localization dictates physiologic signaling of beta 2-adrenoceptors in neonatal cardiac myocytes. *J Biol Chem* 277: 34280–34286.
- Yang S, Nguyen ND, Center JR Jr, Eisman JA, Nguyen TV (2011). Association between beta-blocker use and fracture risk: the Dubbo Osteoporosis Epidemiology Study. *Bone* 48: 451–455.
- Zhou XB, Wulfsen I, Lutz S, Utku E, Sausbier U, Ruth P *et al.* (2008). M2 muscarinic receptors induce airway smooth muscle activation via a dual, Gbetagamma-mediated inhibition of large conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel activity. *J Biol Chem* 283: 21036–21044.