

THE CONCISE GUIDE TO PHARMACOLOGY 2013/14: NUCLEAR HORMONE RECEPTORS

Stephen P.H. Alexander^{*1}, Helen E. Benson², Elena Faccenda², Adam J. Pawson²,
Joanna L. Sharman², Michael Spedding³, John A. Peters⁴, Anthony J. Harmar² and
CGTP Collaborators

^{*}Author for correspondence; steve.alexander@guidetopharmacology.org

¹*School of Life Sciences, University of Nottingham Medical School, Nottingham NG7 2UH, UK*


²*The University/BHF Centre for Cardiovascular Science, University of Edinburgh, Edinburgh EH16 4TJ, UK*

³*Spedding Research Solutions SARL, Le Vésinet 78110, France*

⁴*Neuroscience Division, Medical Education Institute, Ninewells Hospital and Medical School, University of Dundee, Dundee DD1 9SY, UK*



BRITISH
PHARMACOLOGICAL
SOCIETY



Abstract

The Concise Guide to PHARMACOLOGY 2013/14 provides concise overviews of the key properties of over 2000 human drug targets with their pharmacology, plus links to an open access knowledgebase of drug targets and their ligands (www.guidetopharmacology.org), which provides more detailed views of target and ligand properties. The full contents can be found at <http://onlinelibrary.wiley.com/doi/10.1111/bph.12444/full>.

Nuclear hormone receptors are one of the seven major pharmacological targets into which the Guide is divided, with the others being G protein-coupled receptors, ligand-gated ion channels, ion channels, catalytic receptors, transporters and enzymes. These are presented with nomenclature guidance and summary information on the best available pharmacological tools, alongside key references and suggestions for further reading. A new landscape format has easy to use tables comparing related targets.

It is a condensed version of material contemporary to late 2013, which is presented in greater detail and constantly updated on the website www.guidetopharmacology.org, superseding data presented in previous Guides to Receptors and Channels. It is produced in conjunction with NC-IUPHAR and provides the official IUPHAR classification and nomenclature for human drug targets, where appropriate. It consolidates information previously curated and displayed separately in IUPHAR-DB and the Guide to Receptors and Channels, providing a permanent, citable, point-in-time record that will survive database updates.

An Introduction to Nuclear Hormone Receptors

Nuclear hormone receptors are specialised transcription factors with commonalities of sequence and structure, which bind as homo- or heterodimers to specific consensus sequences of DNA (response elements) in the promoter region of particular target genes. They regulate (either promoting or repressing) transcription of these target genes in response to a variety of endogenous ligands. Endogenous agonists are hydrophobic entities which, when bound to the receptor promote conformational changes in the receptor to allow recruitment (or dissociation) of protein partners, generating a large multiprotein complex.

Two major subclasses of nuclear hormone receptors with identified endogenous agonists can be identified: steroid and non-steroid hormone receptors. Steroid hormone receptors function typically as dimeric entities and are thought to be resident outside the nucleus in the unliganded state in a complex with chaperone proteins, which are liberated upon agonist binding. Migration to the nucleus and interaction with other regulators of gene transcription, including RNA polymerase, acetyltransferases and deacetylases, allows gene transcription to be regulated. Non-steroid hormone receptors typically exhibit a greater distribution

in the nucleus in the unliganded state and interact with other nuclear hormone receptors to form heterodimers, as well as with other regulators of gene transcription, leading to changes in gene transcription upon agonist binding.

Selectivity of gene regulation is brought about through interaction of nuclear hormone receptors with particular consensus sequences of DNA, which are arranged typically as repeats or inverted palindromes to allow accumulation of multiple transcription factors in the promoter regions of genes.

Searchable database: <http://www.guidetopharmacology.org/index.jsp>

Full Contents of Concise Guide: <http://onlinelibrary.wiley.com/doi/10.1111/bph.12444/full>



Nuclear hormone receptors 1652

Acknowledgements

We wish to acknowledge the tremendous help provided by the Consultants to the Guides past and present (see list in the Overview, p. 1452). We are also extremely grateful for the financial contributions from the British Pharmacological Society, the International Union of Basic and Clinical Pharmacology, the Wellcome Trust (099156/Z/12/Z), which support the website and the University of Edinburgh, who host the guidetopharmacology.org website.

Conflict of interest

The authors state that there is no conflict of interest to disclose.

Further reading

Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* 58: 685–704.

List of records presented

1654 1A. Thyroid Hormone Receptors
1655 1B. Retinoic acid receptors
1656 1C. Peroxisome proliferator-activated receptors
1657 1D. Rev-Erb receptors
1658 1F. Retinoic acid-related orphans
1659 1H. Liver X receptor-like receptors
1660 1I. Vitamin D receptor-like receptors
1661 2A. Hepatocyte nuclear factor-4 receptors
1662 2B. Retinoid X receptors
1663 2C. Testicular receptors
1664 2E. Tailless-like receptors
1665 2F. COUP-TF-like receptors
1666 3B. Estrogen-related receptors
1667 4A. Nerve growth factor IB-like receptors
1668 5A. Fushi tarazu F1-like receptors
1669 6A. Germ cell nuclear factor receptors
1670 0B. DAX-like receptors
1671 Steroid hormone receptors



1A. Thyroid Hormone Receptors

Overview: Thyroid hormone receptors (TRs, nomenclature as agreed by [NC-IUPHAR](#) Committee on Nuclear Receptors, [3]) are nuclear hormone receptors of the NR1A family, with diverse roles regulating macronutrient metabolism, cognition and

cardiovascular homeostasis. TRs are activated by thyroxine (T_4) and thyroid hormone (T_3). Once activated by a ligand, the receptor acts as a transcription factor either as a monomer, homodimer or heterodimer with members of the retinoid X receptor

family. NH-3 has been described as an antagonist at TRs with modest selectivity for TR β [4].

Nomenclature	Thyroid hormone receptor- α	Thyroid hormone receptor- β
Systematic nomenclature	NR1A1	NR1A2
HGNC, UniProt	THRA, P10827	THRB, P10828
Rank order of potency	$T_3 > T_4$	$T_3 > T_4$
Selective agonists (pK _i)	–	GC-1 (pK _d 10.17) [2,5]

Comments: An interaction with integrin $\alpha V\beta 3$ has been suggested to underlie plasma membrane localization of TRs and non-genomic signalling [1]. One splice variant, TR α_2 , lacks a functional DNA-binding domain and appears to act as a transcription suppressor.

Although radioligand binding assays have been described for these receptors, the radioligands are not commercially available.

Further reading

Bianco AC. (2011) Minireview: cracking the metabolic code for thyroid hormone signaling. *Endocrinology* **152**: 3306–3311. [PMID:21712363]
 Brent GA. (2012) Mechanisms of thyroid hormone action. *J Clin Invest* **122**: 3035–3043. [PMID:22945636]
 Flamant F, Baxter JD, Forrest D, Refetoff S, Samuels H, Scanlan TS, Vennström B, Samarut J. (2006) International Union of Pharmacology. LIX. The pharmacology and classification of the nuclear receptor superfamily: thyroid hormone receptors. *Pharmacol Rev* **58**: 705–711. [PMID:17132849]

Pramfalk C, Pedrelli M, Parini P. (2011) Role of thyroid receptor β in lipid metabolism. *Biochim Biophys Acta* **1812**: 929–937. [PMID:21194564]
 Sirakov M, Plateroti M. (2011) The thyroid hormones and their nuclear receptors in the gut: from developmental biology to cancer. *Biochim Biophys Acta* **1812**: 938–946. [PMID:21194566]
 Tancevski I, Rudling M, Eller P. (2011) Thyromimetics: a journey from bench to bed-side. *Pharmacol Ther* **131**: 33–39. [PMID:21504761]



1B. Retinoic acid receptors

Overview: Retinoic acid receptors (nomenclature as agreed by NC-IUPHAR Committee on Nuclear Receptors, [8]) are nuclear hormone receptors of the NR1B family activated by the vitamin A-derived agonists all-trans-retinoic acid (ATRA) and 9-cis-retinoic acid, and the RAR-selective synthetic agonists TTNPB and adapalene.

Nomenclature	Retinoic acid receptor- α	Retinoic acid receptor- β	Retinoic acid receptor- γ
Systematic nomenclature	NR1B1	NR1B2	NR1B3
HGNC, UniProt	RARA, P10276	RARB, P10826	RARG, P13631
Selective agonists (pK _i)	Ro 40-6055 [7,11,18], BMS753 (8.7) [10]	AC261066 (pEC ₅₀ 7.9 – 8.1) [15], AC55649 (pEC ₅₀ 6.5 – 7.3) [15]	AHPN [16]
Selective antagonists (pK _i)	Ro 41-5253 (pIC ₅₀ 6.3 – 7.2) [6,12]	–	MM 11253 [13]

Comments: Ro 41-5253 has been suggested to be a PPAR γ agonist [17]. LE135 is an antagonist with selectivity for RAR α and RAR β compared with RAR γ [14]. [9] is a family-selective antagonist.

Further reading

Bour G, Lalevée S, Rochette-Egly C. (2007) Protein kinases and the proteasome join in the combinatorial control of transcription by nuclear retinoic acid receptors. *Trends Cell Biol* **17**: 302–309. [PMID:17467991]

Duong V, Rochette-Egly C. (2011) The molecular physiology of nuclear retinoic acid receptors. From health to disease. *Biochim Biophys Acta* **1812**: 1023–1031. [PMID:20970498]

Germain P, Chambon P, Eichele G, Evans RM, Lazar MA, Leid M, De Lera AR, Lotan R, Mangelsdorf DJ, Gronemeyer H. (2006) International Union of Pharmacology. LX. Retinoic acid receptors. *Pharmacol Rev* **58**: 712–725. [PMID:17132850]

Maden M. (2007) Retinoic acid in the development, regeneration and maintenance of the nervous system. *Nat Rev Neurosci* **8**: 755–765. [PMID:17882253]

Mark M, Ghyselinck NB, Chambon P. (2006) Function of retinoid nuclear receptors: lessons from genetic and pharmacological dissections of the retinoic acid signaling pathway during mouse embryogenesis. *Annu Rev Pharmacol Toxicol* **46**: 451–480. [PMID:16402912]



1C. Peroxisome proliferator-activated receptors

Overview: Peroxisome proliferator-activated receptors (PPARs, nomenclature as agreed by [NC-IUPHAR](#) Committee on Nuclear Receptors, [33]) are nuclear hormone receptors of the NR1C family, with diverse roles regulating lipid homeostasis, cellular differentiation, proliferation and the immune response. PPARs have many potential endogenous agonists [21,33], including 15-deoxy- $\Delta^{12,14}$ -PGJ₂, prostacyclin (PGI₂), many fatty acids and

their oxidation products, lysophosphatidic acid (LPA) [32], 13-HODE, 15S-HETE, Paz-PC, azelaoyl-PAF and leukotriene B₄ (LTB₄). bezafibrate acts as a non-selective agonist for the PPAR family [41]. These receptors also bind hypolipidaemic drugs (PPAR α) and anti-diabetic thiazolidinediones (PPAR γ), as well as many non-steroidal anti-inflammatory drugs, such as sulindac and indomethacin. Once activated by a ligand, the receptor

forms a heterodimer with members of the retinoid X receptor family and can act as a transcription factor. Although radioligand binding assays have been described for all three receptors, the radioligands are not commercially available. Commonly, receptor occupancy studies are conducted using fluorescent ligands and truncated forms of the receptor limited to the ligand binding domain.


Nomenclature	Peroxisome proliferator-activated receptor- α	Peroxisome proliferator-activated receptor- β/δ	Peroxisome proliferator-activated receptor- γ
Systematic nomenclature	NR1C1	NR1C2	NR1C3
HGNC, UniProt	<i>PPARA</i> , Q07869	<i>PPARD</i> , Q03181	<i>PPARG</i> , P37231
Selective agonists (pK _i)	ciprofibrate, GW7647 (pEC ₅₀ 8.2) [22–23], CP-775146 (pEC ₅₀ 7.3) [28], pirinixic acid (pEC ₅₀ 5.3) [41]	GW501516 (pEC ₅₀ 9.0) [35], GW0742X (pIC ₅₀ 9.0) [25,39]	rosiglitazone (pK _d 7.4) [27,31,44], GW1929 (8.8) [22], CDDO (Partial agonist) (8.0) [40], troglitazone (5.8) [19], ciglitazone (pEC ₅₀ 4.6) [27], troglitazone (pIC ₅₀ 6.3) [27,44], pioglitazone (pIC ₅₀ 6.2) [27,37,44]
Selective antagonists (pK _i)	GW6471 (pIC ₅₀ 6.6) [42]	GSK0660 (pIC ₅₀ 6.5) [38]	T0070907 (9.0) [29], CDDO-Me (6.9) [40], GW9662 (Irreversible inhibition) (pIC ₅₀ 8.1) [30]

Comments: As with the estrogen receptor antagonists, many agents show tissue-selective efficacy (*e.g.* [20,34,36]). Agonists with mixed activity at PPAR α and PPAR γ have also been described (*e.g.* [24,26,43]).

Further reading

Huang JV, Greyson CR, Schwartz GG. (2012) PPAR- γ as a therapeutic target in cardiovascular disease: evidence and uncertainty. *J Lipid Res* 53: 1738–1754. [PMID:22685322]
 Michalik L, Auwerx J, Berger JP, Chatterjee VK, Glass CK, Gonzalez FJ, Grimaldi PA, Kadowaki T, Lazar MA, O'Rahilly S, Palmer CN, Plutzky J, Reddy JK, Spiegelman BM, Staels B, Wahli W. (2006) International Union of Pharmacology. LXI. Peroxisome proliferator-activated receptors. *Pharmacol Rev* 58: 726–741. [PMID:17132851]
 Michalik L, Wahli W. (2008) PPARs Mediate Lipid Signaling in Inflammation and Cancer. *PPAR Res* 2008: 134059. [PMID:19125181]
 Peters JM, Shah YM, Gonzalez FJ. (2012) The role of peroxisome proliferator-activated receptors in carcinogenesis and chemoprevention. *Nat Rev Cancer* 12: 181–195. [PMID:22318237]

Pirat C, Farce A, Lebègue N, Renault N, Furman C, Millet R, Yous S, Specia S, Berthelot P, Desreumaux P *et al.* (2012) Targeting peroxisome proliferator-activated receptors (PPARs): development of modulators. *J Med Chem* 55: 4027–4061. [PMID:22260081]
 Varga T, Czimmerer Z, Nagy L. (2011) PPARs are a unique set of fatty acid regulated transcription factors controlling both lipid metabolism and inflammation. *Biochim Biophys Acta* 1812: 1007–1022. [PMID:21382489]
 Youssef J, Badr M. (2011) Peroxisome proliferator-activated receptors and cancer: challenges and opportunities. *Br J Pharmacol* 164: 68–82. [PMID:21449912]



1D. Rev-Erb receptors

Overview: Rev-erb receptors (nomenclature as agreed by NC-IUPHAR committee on nuclear hormone receptors [45]) have yet to be officially paired with an endogenous ligand, but are thought to be activated by heme.

Nomenclature	Rev-Erb- α	Rev-Erb- β
Systematic nomenclature	NR1D1	NR1D2
HGNC, UniProt	NR1D1, P20393	NR1D2, Q14995
Endogenous agonists (pK _i)	heme (Selective) [48–49]	heme (Selective) [48–49]
Selective agonists (pK _i)	GSK4112 (pEC ₅₀ 6.4) [46], GSK4112 (pIC ₅₀ 5.6) [47]	–
Selective antagonists (pK _i)	SR8278 (pIC ₅₀ 6.5) [47]	–

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
- Duez H, Staels B. (2009) Rev-erb-alpha: an integrator of circadian rhythms and metabolism. *J Appl Physiol* **107**: 1972–1980. [PMID:19696364]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
- Phelan CA, Gampe RT, Lambert MH, Parks DJ, Montana V, Bynum J, Broderick TM, Hu X, Williams SP, Nolte RT, Lazar MA. (2010) Structure of Rev-erbalph bound to N-CoR reveals a unique mechanism of nuclear receptor-co-repressor interaction. *Nat Struct Mol Biol* **17**: 808–814. [PMID:20581824]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]
- Yin L, Wu N, Lazar MA. (2010) Nuclear receptor Rev-erbalph: a heme receptor that coordinates circadian rhythm and metabolism. *Nucl Recept Signal* **8**: e001. [PMID:20414452]



1F. Retinoic acid-related orphans

Overview: Retinoic acid receptor-related orphan receptors (ROR, nomenclature as agreed by [NC-IUPHAR](#) Committee on Nuclear Receptors, [50]) have yet to be assigned a definitive endogenous ligand, although ROR α may be synthesized with a ‘captured’ agonist such as cholesterol [52–53].

Nomenclature	RAR-related orphan receptor- α	RAR-related orphan receptor- β	RAR-related orphan receptor- γ
Systematic nomenclature	NR1F1	NR1F2	NR1F3
HGNC, UniProt	RORA, P35398	RORB, Q92753	RORC, P51449
Endogenous agonists (pK _i)	cholesterol (Selective) [53–54]	–	–
Selective agonists (pK _i)	7-hydroxycholesterol [51], cholesterol sulphate [51,53]	–	–

Comments: all-trans-retinoic acid shows selectivity for ROR β within the ROR family [55]. ROR α has been suggested to be a nuclear receptor responding to melatonin [56].

Further reading

Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* 58: 798–836. [PMID:17132856]



1H. Liver X receptor-like receptors

Overview: Liver X and farnesoid X receptors (LXR and FXR, nomenclature as agreed by NC-IUPHAR Committee on Nuclear Receptors, [62]) are members of a steroid analogue-activated nuclear receptor subfamily (ENFSM00500000269785), which form heterodimers with members of the retinoid X receptor family. Endogenous ligands for LXRs include hydroxycholesterols (OHC), while FXRs appear to be activated by bile acids.

Nomenclature	Farnesoid X receptor	Farnesoid X receptor-β	Liver X receptor-α	Liver X receptor-β
Systematic nomenclature	NR1H4	NR1H5	NR1H3	NR1H2
HGNC, UniProt	NR1H4, Q96RI1	NR1H5P, -	NR1H3, Q13133	NR1H2, P55055
Potency order	chenodeoxycholic acid > lithocholic acid, deoxycholic acid [60,64]	–	20S-hydroxycholesterol, 22R-hydroxycholesterol, 24(S)-hydroxycholesterol > 25-hydroxycholesterol, 27-hydroxycholesterol [59]	20S-hydroxycholesterol, 22R-hydroxycholesterol, 24(S)-hydroxycholesterol > 25-hydroxycholesterol, 27-hydroxycholesterol [59]
Selective agonists (pK _i)	GW4064 (pEC ₅₀ 7.8) [61], ECDCA (pEC ₅₀ 7.0) [65], fexaramine (pEC ₅₀ 6.6) [58]	–	–	–
Selective antagonists (pK _i)	guggulsterone (pIC ₅₀ 5.7 – 6.0) [67]	–	–	–
Endogenous agonists (pK _i)	–	lanosterol (pEC ₅₀ 6.0 - Mouse) [63]	–	–

Comments: T0901317 [66] and GW3965 [57] are synthetic agonists acting at both LXRα and LXRβ with less than 10-fold selectivity. NR1H5P (FXRβ) is a pseudogene in man, but active in the mouse.

Further reading

- A-González N, Castrillo A. (2011) Liver X receptors as regulators of macrophage inflammatory and metabolic pathways. *Biochim Biophys Acta* **1812**: 982–994. [PMID:21193033]
- Calkin AC, Tontonoz P. (2010) Liver x receptor signaling pathways and atherosclerosis. *Arterioscler Thromb Vasc Biol* **30**: 1513–1518. [PMID:20631351]
- Chen WD, Wang YD, Meng Z, Zhang L, Huang W. (2011) Nuclear bile acid receptor FXR in the hepatic regeneration. *Biochim Biophys Acta* **1812**: 888–892. [PMID:21167938]
- Claudel T, Zollner G, Wagner M, Trauner M. (2011) Role of nuclear receptors for bile acid metabolism, bile secretion, cholestasis, and gallstone disease. *Biochim Biophys Acta* **1812**: 867–878. [PMID:21194565]
- El-Hajjaji FZ, Oumeddour A, Pommier AJ, Ouvrier A, Viennois E, Dufour J, Caira F, Drevet JR, Volle DH, Baron S *et al.* (2011) Liver X receptors, lipids and their reproductive secrets in the male. *Biochim Biophys Acta* **1812**: 974–981. [PMID:21334438]
- Gadaleta RM, van Mil SW, Oldenburg B, Siersema PD, Klomp LW, van Erpecum KJ. (2010) Bile acids and their nuclear receptor FXR: Relevance for hepatobiliary and gastrointestinal disease. *Biochim Biophys Acta* **1801**: 683–692. [PMID:20399894]
- Gardmo C, Tamburro A, Modica S, Moschetta A. (2011) Proteomics for the discovery of nuclear bile acid receptor FXR targets. *Biochim Biophys Acta* **1812**: 836–841. [PMID:21439373]
- Hageman J, Herrema H, Groen AK, Kuipers F. (2010) A role of the bile salt receptor FXR in atherosclerosis. *Arterioscler Thromb Vasc Biol* **30**: 1519–1528. [PMID:20631352]
- Kemper JK. (2011) Regulation of FXR transcriptional activity in health and disease: Emerging roles of FXR cofactors and post-translational modifications. *Biochim Biophys Acta* **1812**: 842–850. [PMID:21130162]
- Matsubara T, Li F, Gonzalez FJ. (2013) FXR signaling in the enterohepatic system. *Mol Cell Endocrinol* **368**: 17–29. [PMID:22609541]
- Mencarelli A, Fiorucci S. (2010) FXR an emerging therapeutic target for the treatment of atherosclerosis. *J Cell Mol Med* **14**: 79–92. [PMID:20041971]
- Moore DD, Kato S, Xie W, Mangelsdorf DJ, Schmidt DR, Xiao R, Kliewer SA. (2006) International Union of Pharmacology. LXII. The NR1H and NR1I receptors: constitutive androstane receptor, pregnane X receptor, farnesoid X receptor alpha, farnesoid X receptor beta, liver X receptor alpha, liver X receptor beta, and vitamin D receptor. *Pharmacol Rev* **58**: 742–759. [PMID:17132852]



11. Vitamin D receptor-like receptors

Overview: Vitamin D (VDR), Pregnane X (PXR) and Constitutive Androstane (CAR) receptors (nomenclature as agreed by NC-IUPHAR Committee on Nuclear Receptors, [79]) are members

of the NR1I family of nuclear receptors, which form heterodimers with members of the retinoid X receptor family. PXR and CAR are activated by a range of exogenous compounds, with no

established endogenous physiological agonists, although high concentrations of bile acids and bile pigments activate PXR and CAR[79].

Nomenclature	Vitamin D receptor	Pregnane X receptor	Constitutive androstane receptor
Systematic nomenclature	NR1I1	NR1I2	NR1I3
HGNC, UniProt	VDR, P11473	NR1I2, O75469	NR1I3, Q14994
Endogenous agonists (pK _i)	1,25-dihydroxyvitamin D3 (pK _d 8.9 – 9.2) [68,71]	17β-estradiol (Selective) [74]	–
Selective agonists (pK _i)	EB1089 (pK _d 9.57) [70,84]	hyperforin (pEC ₅₀ 7.6) [80,83], rifampicin (pEC ₅₀ 5.5 – 6.0) [69,76], lovastatin (pEC ₅₀ 5.3 – 6.0) [76], pregnanedione (pIC ₅₀ 6.4) [74]	TCPOBOP (pEC ₅₀ 7.7 - Mouse) [82], CITCO (pEC ₅₀ 7.3) [77]
Selective antagonists (pK _i)	TEI-9647 [78], ZK159222 (pIC ₅₀ 7.5) [72–73]	–	–
Comment	–	–	clotrimazole [81] and T0901317 [75] although acting at other sites, function as antagonists of the constitutive androstane receptor

Further reading

Bikle DD. (2011) Vitamin D: an ancient hormone. *Exp Dermatol* 20: 7–13. [PMID:21197695]

Campbell FC, Xu H, El-Tanani M, Crowe P, Bingham V. (2010) The yin and yang of vitamin D receptor (VDR) signaling in neoplastic progression: operational networks and tissue-specific growth control. *Biochem Pharmacol* 79: 1–9. [PMID:19737544]

Chen Y, Tang Y, Guo C, Wang J, Boral D, Nie D. (2012) Nuclear receptors in the multidrug resistance through the regulation of drug-metabolizing enzymes and drug transporters. *Biochem Pharmacol* 83: 1112–1126. [PMID:22326308]

Cheng J, Shah YM, Gonzalez FJ. (2012) Pregnane X receptor as a target for treatment of inflammatory bowel disorders. *Trends Pharmacol Sci* 33: 323–330. [PMID:22609277]

Ihunnah CA, Jiang M, Xie W. (2011) Nuclear receptor PXR, transcriptional circuits and metabolic relevance. *Biochim Biophys Acta* 1812: 956–963. [PMID:21295138]

Kachaylo EM, Pustyl'nyak VO, Lyakhovich VV, Gulyaeva LF. (2011) Constitutive androstane receptor (CAR) is a xenosensor and target for therapy. *Biochemistry Mosc* 76: 1087–1097. [PMID:22098234]

Krishnan AV, Feldman D. (2011) Mechanisms of the anti-cancer and anti-inflammatory actions of vitamin D. *Annu Rev Pharmacol Toxicol* 51: 311–336. [PMID:20936945]

Moore DD, Kato S, Xie W, Mangelsdorf DJ, Schmidt DR, Xiao R, Kliewer SA. (2006) International Union of Pharmacology. LXII. The NR1H and NR1I receptors: constitutive androstane receptor, pregnane X receptor, farnesoid X receptor alpha, farnesoid X receptor beta, liver X receptor alpha, liver X receptor beta, and vitamin D receptor. *Pharmacol Rev* 58: 742–759. [PMID:17132852]

Plum LA, DeLuca HF. (2010) Vitamin D, disease and therapeutic opportunities. *Nat Rev Drug Discov* 9: 941–955. [PMID:21119732]

Staudinger JL, Woody S, Sun M, Cui W. (2013) Nuclear-receptor-mediated regulation of drug- and bile-acid-transporter proteins in gut and liver. *Drug Metab Rev* 45: 48–59. [PMID:23330541]



2A. Hepatocyte nuclear factor-4 receptors

Overview: Hepatocyte nuclear factor-4 receptors (nomenclature as agreed by NC-IUPHAR committee on nuclear hormone receptors [85]) have yet to be officially paired with an endogenous ligand, although linoleic acid has been described to activate HNF4 α receptors.

Nomenclature	Hepatocyte nuclear factor-4- α	Hepatocyte nuclear factor-4- γ
Systematic nomenclature	NR2A1	NR2A2
HGNC, UniProt	<i>HNF4A</i> , P41235	<i>HNF4G</i> , Q14541
Endogenous agonists (p <i>K</i> _i)	linoleic acid (Selective) [87]	–
Selective antagonists (p <i>K</i> _i)	BI6015 [86]	–

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
- Hwang-Verslues WW, Sladek FM. (2010) HNF4 α –role in drug metabolism and potential drug target?. *Curr Opin Pharmacol* **10**: 698–705. [PMID:20833107]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]



2B. Retinoid X receptors

Overview: Retinoid X receptors (nomenclature as agreed by NC-IUPHAR Committee on Nuclear Receptors, [90]) are NR2B family members activated by 9-cis-retinoic acid and the RXR-selective agonists bexarotene and LG100268, sometimes referred to as rexinoids. UVI3003 [93] and HX531 [89] have been described as a pan-RXR antagonists. These receptors form RXR–RAR heterodimers and RXR–RXR homodimers [88,92].

Nomenclature	Retinoid X receptor- α	Retinoid X receptor- β	Retinoid X receptor- γ
Systematic nomenclature	NR2B1	NR2B2	NR2B3
HGNC, UniProt	RXRA, P19793	RXRB, P28702	RXRG, P48443
Selective agonists (pK _i)	CD3254 (pIC ₅₀ 8.5) [91]	–	–

Further reading

- Bour G, Lalevée S, Rochette-Egly C. (2007) Protein kinases and the proteasome join in the combinatorial control of transcription by nuclear retinoic acid receptors. *Trends Cell Biol* **17**: 302–309. [PMID:17467991]
- Duong V, Rochette-Egly C. (2011) The molecular physiology of nuclear retinoic acid receptors. From health to disease. *Biochim Biophys Acta* **1812**: 1023–1031. [PMID:20970498]
- Germain P, Chambon P, Eichele G, Evans RM, Lazar MA, Leid M, De Lera AR, Lotan R, Mangelsdorf DJ, Gronemeyer H. (2006) International Union of Pharmacology. LXIII. Retinoid X receptors. *Pharmacol Rev* **58**: 760–772. [PMID:17132853]
- Lefebvre P, Benomar Y, Staels B. (2010) Retinoid X receptors: common heterodimerization partners with distinct functions. *Trends Endocrinol Metab* **21**: 676–683. [PMID:20674387]
- Maden M. (2007) Retinoic acid in the development, regeneration and maintenance of the nervous system. *Nat Rev Neurosci* **8**: 755–765. [PMID:17882253]
- Mark M, Ghyselinck NB, Chambon P. (2006) Function of retinoid nuclear receptors: lessons from genetic and pharmacological dissections of the retinoic acid signaling pathway during mouse embryogenesis. *Annu Rev Pharmacol Toxicol* **46**: 451–480. [PMID:16402912]
- Pérez E, Bourguet W, Gronemeyer H, de Lera AR. (2011) Modulation of RXR function through ligand design. *Biochim Biophys Acta* [Epub ahead of print]. [PMID:21515403]



2C. Testicular receptors

Overview: Testicular receptors (nomenclature as agreed by [NC-IUPHAR](#) committee on nuclear hormone receptors [94]) have yet to be officially paired with an endogenous ligand, although testicular receptor 4 has been reported to respond to retinoids.

Nomenclature	Testicular receptor 2	Testicular receptor 4
Systematic nomenclature	NR2C1	NR2C2
HGNC, UniProt	NR2C1, P13056	NR2C2, P49116
Endogenous agonists (pK _i)	–	all-trans-retinoic acid (Selective) [96], retinol (Selective) [96]
Comment	Forms a heterodimer with TR4; gene disruption appears without effect on testicular development or function [95]	Forms a heterodimer with TR2

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
- Duez H, Staels B. (2009) Rev-erb- α : an integrator of circadian rhythms and metabolism. *J Appl Physiol* **107**: 1972–1980. [PMID:19696364]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
- Phelan CA, Gampe RT, Lambert MH, Parks DJ, Montana V, Bynum J, Broderick TM, Hu X, Williams SP, Nolte RT, Lazar MA. (2010) Structure of Rev-erb α bound to N-CoR reveals a unique mechanism of nuclear receptor-co-repressor interaction. *Nat Struct Mol Biol* **17**: 808–814. [PMID:20581824]
- Schimmer BP, White PC. (2010) Minireview: steroidogenic factor 1: its roles in differentiation, development, and disease. *Mol Endocrinol* **24**: 1322–1337. [PMID:20203099]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]
- Yin L, Wu N, Lazar MA. (2010) Nuclear receptor Rev-erb α : a heme receptor that coordinates circadian rhythm and metabolism. *Nucl Recept Signal* **8**: e001. [PMID:20414452]
- Zhang Y, Hagedorn CH, Wang L. (2011) Role of nuclear receptor SHP in metabolism and cancer. *Biochim Biophys Acta* **1812**: 893–908. [PMID:20970497]
- Zhao Y, Bruemmer D. (2010) NR4A orphan nuclear receptors: transcriptional regulators of gene expression in metabolism and vascular biology. *Arterioscler Thromb Vasc Biol* **30**: 1535–1541. [PMID:20631354]



2E. Tailless-like receptors

Overview: Tailless-like receptors (nomenclature as agreed by [NC-IUPHAR](#) committee on nuclear hormone receptors [97]) have yet to be officially paired with an endogenous ligand.

Nomenclature	TLX	PNR
Systematic nomenclature	NR2E1	NR2E3
HGNC, UniProt	<i>NR2E1</i> , Q9Y466	<i>NR2E3</i> , Q9Y5X4
Comment	Gene disruption is associated with abnormal brain development [98–99]	–

Further reading

Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
 Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]

Gui H, Li ML, Tsai CC. (2011) A tale of tailless. *Dev Neurosci* **33**: 1–13. [PMID:21124006]
 Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
 Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]



2F. COUP-TF-like receptors

Overview: COUP-TF-like receptors (nomenclature as agreed by [NC-IUPHAR](#) committee on nuclear hormone receptors [100]) have yet to be officially paired with an endogenous ligand.

Nomenclature	COUP-TF1	COUP-TF2	V-erbA-related gene
Systematic nomenclature	NR2F1	NR2F2	NR2F6
HGNC, UniProt	NR2F1, P10589	NR2F2, P24468	NR2F6, P10588
Comment	Gene disruption is perinatally lethal [102]	Gene disruption is embryonically lethal [101]	Gene disruption impairs CNS development [103]

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* 58: 798–836. [PMID:17132856]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* 58: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* 72: 247–272. [PMID:20148675]
- Lin FJ, Qin J, Tang K, Tsai SY, Tsai MJ. (2011) Coup d'Etat: an orphan takes control. *Endocr Rev* 32: 404–421. [PMID:21257780]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* 334: 3–13. [PMID:20615454]



3B. Estrogen-related receptors

Overview: Estrogen-related receptors (nomenclature as agreed by [NC-IUPHAR](#) committee on nuclear hormone receptors [104]) have yet to be officially paired with an endogenous ligand.

Nomenclature	Estrogen-related receptor- α	Estrogen-related receptor- β	Estrogen-related receptor- γ
Systematic nomenclature	NR3B1	NR3B2	NR3B3
HGNC, UniProt	ESRRA, P11474	ESRRB, O95718	ESRRG, P62508
Comment	Activated by some dietary flavonoids [105]; activated by the synthetic agonist GSK4716 [108] and blocked by XCT790 [106]	May be activated by DY131 [107]	May be activated by DY131 [107]

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* 58: 798–836. [PMID:17132856]
- Deblois G, Giguère V. (2011) Functional and physiological genomics of estrogen-related receptors (ERRs) in health and disease. *Biochim Biophys Acta* 1812: 1032–1040. [PMID:21172432]
- Deblois G, Giguère V. (2013) Oestrogen-related receptors in breast cancer: control of cellular metabolism and beyond. *Nat Rev Cancer* 13: 27–36. [PMID:23192231]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* 58: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* 72: 247–272. [PMID:20148675]
- Hwang-Verslues WW, Sladek FM. (2010) HNF4 α –role in drug metabolism and potential drug target?. *Curr Opin Pharmacol* 10: 698–705. [PMID:20833107]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* 334: 3–13. [PMID:20615454]




4A. Nerve growth factor IB-like receptors

Overview: Nerve growth factor IB-like receptors (nomenclature as agreed by [NC-IUPHAR](#) committee on nuclear hormone receptors [110]) have yet to be officially paired with an endogenous ligand.

Nomenclature	Nerve Growth factor IB	Nuclear receptor related 1	Neuron-derived orphan receptor 1
Systematic nomenclature	NR4A1	NR4A2	NR4A3
HGNC, UniProt	NR4A1, P22736	NR4A2, P43354	NR4A3, Q92570
Comment	An endogenous agonist, cytosporone B, has been described [113], although structural analysis and molecular modelling has not identified a ligand binding site [109,111–112]	–	–

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
- McMorrow JP, Murphy EP. (2011) Inflammation: a role for NR4A orphan nuclear receptors?. *Biochem Soc Trans* **39**: 688–693. [PMID:21428963]
- Mohan HM, Aherne CM, Rogers AC, Baird AW, Winter DC, Murphy EP. (2012) Molecular pathways: the role of NR4A orphan nuclear receptors in cancer. *Clin Cancer Res* **18**: 3223–3228. [PMID:22566377]
- Pearen MA, Muscat GE. (2010) Minireview: Nuclear hormone receptor 4A signaling: implications for metabolic disease. *Mol Endocrinol* **24**: 1891–1903. [PMID:20392876]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]
- van Tiel CM, de Vries CJ. (2012) NR4A1 in the vessel wall. *J Steroid Biochem Mol Biol* **130**: 186–193. [PMID:21277978]
- Zhao Y, Bruemmer D. (2010) NR4A orphan nuclear receptors: transcriptional regulators of gene expression in metabolism and vascular biology. *Arterioscler Thromb Vasc Biol* **30**: 1535–1541. [PMID:20631354]




5A. Fushi tarazu F1-like receptors

Overview: Fushi tarazu F1-like receptors (nomenclature as agreed by [NC-IUPHAR](#) committee on nuclear hormone receptors [114]) have yet to be officially paired with an endogenous ligand.

Nomenclature	Steroidogenic factor 1	Liver receptor homolog-1
Systematic nomenclature	NR5A1	NR5A2
HGNC, UniProt	NR5A1, Q13285	NR5A2, O00482
Comment	Reported to be inhibited by AC45594 [115] and SID7969543 [116]	–

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
- Büdefeld T, Tobet SA, Majdic G. (2012) Steroidogenic factor 1 and the central nervous system. *J Neuroendocrinol* **24**: 225–235. [PMID:21668533]
- El-Khairi R, Martinez-Aguayo A, Ferraz-de-Souza B, Lin L, Achermann JC. (2011) Role of DAX-1 (NR0B1) and steroidogenic factor-1 (NR5A1) in human adrenal function. *Endocr Dev* **20**: 38–46. [PMID:21164257]
- Fernandez-Marcos PJ, Auwerx J, Schoonjans K. (2011) Emerging actions of the nuclear receptor LRH-1 in the gut. *Biochim Biophys Acta* **1812**: 947–955. [PMID:21194563]
- Ferraz-de-Souza B, Lin L, Achermann JC. (2011) Steroidogenic factor-1 (SF-1, NR5A1) and human disease. *Mol Cell Endocrinol* **336**: 198–205. [PMID:21078366]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
- Lazarus KA, Wijayakumara D, Chand AL, Simpson ER, Clyne CD. (2012) Therapeutic potential of Liver Receptor Homolog-1 modulators. *J Steroid Biochem Mol Biol* **130**: 138–146. [PMID:22266285]
- Młynarczuk J, Rękawiecki R. (2010) The role of the orphan receptor SF-1 in the development and function of the ovary. *Reprod Biol* **10**: 177–193. [PMID:21113200]
- Schimmer BP, White PC. (2010) Minireview: steroidogenic factor 1: its roles in differentiation, development, and disease. *Mol Endocrinol* **24**: 1322–1337. [PMID:20203099]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]



6A. Germ cell nuclear factor receptors

Overview: Germ cell nuclear factor receptors (nomenclature as agreed by [NC-IUPHAR](#) committee on nuclear hormone receptors [117]) have yet to be officially paired with an endogenous ligand.

Nomenclature

Germ cell nuclear factor

Systematic nomenclature

NR6A1

HGNC, UniProt

NR6A1, Q15406

Further reading

Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]

Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]



0B. DAX-like receptors

Overview: Dax-like receptors (nomenclature as agreed by NC-IUPHAR committee on nuclear hormone receptors [118]) have yet to be officially paired with an endogenous ligand.

Nomenclature	DAX1	SHP
Systematic nomenclature	NR0B1	NR0B2
HGNC, UniProt	NR0B1, P51843	NR0B2, Q15466

Further reading

- Benoit G, Cooney A, Giguere V, Ingraham H, Lazar M, Muscat G, Perlmann T, Renaud JP, Schwabe J, Sladek F, Tsai MJ, Laudet V. (2006) International Union of Pharmacology. LXVI. Orphan nuclear receptors. *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
- Chanda D, Park JH, Choi HS. (2008) Molecular basis of endocrine regulation by orphan nuclear receptor Small Heterodimer Partner. *Endocr J* **55**: 253–268. [PMID:17984569]
- Ehrlund A, Treuter E. (2012) Ligand-independent actions of the orphan receptors/corepressors DAX-1 and SHP in metabolism, reproduction and disease. *J Steroid Biochem Mol Biol* **130**: 169–179. [PMID:21550402]
- El-Khairi R, Martinez-Aguayo A, Ferraz-de-Souza B, Lin L, Achermann JC. (2011) Role of DAX-1 (NR0B1) and steroidogenic factor-1 (NR5A1) in human adrenal function. *Endocr Dev* **20**: 38–46. [PMID:21164257]
- Germain P, Staels B, Dacquet C, Spedding M, Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol Rev* **58**: 685–704. [PMID:17132848]
- Huang P, Chandra V, Rastinejad F. (2010) Structural overview of the nuclear receptor superfamily: insights into physiology and therapeutics. *Annu Rev Physiol* **72**: 247–272. [PMID:20148675]
- Młynarczyk J, Rękawiecki R. (2010) The role of the orphan receptor SF-1 in the development and function of the ovary. *Reprod Biol* **10**: 177–193. [PMID:21113200]
- Sladek FM. (2011) What are nuclear receptor ligands?. *Mol Cell Endocrinol* **334**: 3–13. [PMID:20615454]
- Zhang Y, Hagedorn CH, Wang L. (2011) Role of nuclear receptor SHP in metabolism and cancer. *Biochim Biophys Acta* **1812**: 893–908. [PMID:20970497]



Steroid hormone receptors

Overview: Steroid hormone receptors (nomenclature as agreed by NC-IUPHAR Committee on Nuclear Receptors, [120,132]) are nuclear hormone receptors of the NR3 class, with endogenous agonists that may be divided into 3-hydroxysteroids (estrone and 17 β -estradiol) and 3-ketosteroids (dihydrotestosterone [DHT], aldosterone, cortisol, corticosterone, progesterone and testosterone). These receptors exist as dimers coupled with chaperone molecules (such as hsp90 (*HSP90AB1*, P08238) and immunophilin FKBP52:*FKBP4*, Q02790), which are shed on binding the steroid hormone. Although rapid signalling phenomena are

observed [130,138], the principal signalling cascade appears to involve binding of the activated receptors to nuclear hormone response elements of the genome, with a 15-nucleotide consensus sequence AGAACAnnnTGTCT (*i.e.* an inverted palindrome) as homo- or heterodimers. They also affect transcription by protein–protein interactions with other transcription factors, such as activator protein 1 (AP-1) and nuclear factor κ B (NF- κ B). Splice variants of each of these receptors can form functional or non-functional monomers that can dimerize to form functional or non-functional receptors. For example, alternative splicing of

PR mRNA produces A and B monomers that combine to produce functional AA, AB and BB receptors with distinct characteristics [148].

A 7TM receptor responsive to estrogen (*GPER1*, Q99527, also known as GPR30, see [137]) has been described. Human orthologues of 7TM ‘membrane progesterin receptors’ (*PAQR7*, *PAQR8* and *PAQR5*), initially discovered in fish [151–152], appear to localize to intracellular membranes and respond to ‘non-genomic’ progesterone analogues independently of G proteins [142].

3A. Estrogen receptors

Nomenclature	Estrogen receptor- α	Estrogen receptor- β
Systematic nomenclature	NR3A1	NR3A2
HGNC, UniProt	<i>ESR1</i> , P03372	<i>ESR2</i> , Q92731
Selective agonists (p <i>K</i> _i)	PPT (9.64) [128,143]	ERB 041 [133], diarylpropionitril (8.6) [135,143], WAY200070 (pIC ₅₀ 8.52 – 9.0) [133]
Selective antagonists (p <i>K</i> _i)	methyl-piperidino-pyrazole (8.57) [145]	PHTPP [119], R,R-THC (8.44) [134,146]

Comments: R,R-THC exhibits partial agonist activity at ER α [134,146]. Estrogen receptors may be blocked non-selectively by tamoxifen and raloxifene and labelled by [³H]17 β -estradiol and [³H]tamoxifen. Many agents thought initially to be antagonists

at estrogen receptors appear to have tissue-specific efficacy (*e.g.* tamoxifen is an antagonist at estrogen receptors in the breast, but is an agonist at estrogen receptors in the uterus), hence the descriptor SERM (selective estrogen receptor modula-

tor) or SnuRM (selective nuclear receptor modulator). Y134 has been suggested to be an ER α -selective estrogen receptor modulator [136].



3C. 3-Ketosteroid receptors

Nomenclature	Androgen receptor	Glucocorticoid receptor	Mineralocorticoid receptor	Progesterone receptor
Systematic nomenclature	NR3C4	NR3C1	NR3C2	NR3C3
HGNC, UniProt	AR, P10275	NR3C1, P04150	NR3C2, P08235	PGR, P06401
Rank order of potency	dihydrotestosterone>testosterone	cortisol,corticosterone>>aldosterone, deoxycortisone [139]	corticosterone,cortisol,aldosterone, progesterone [139]	progesterone
Endogenous agonists (pK _i)	dihydrotestosterone (pK _d 9.3) [147]	–	aldosterone (Selective) (pIC ₅₀ 9.8 – 10.0) [126,139]	progesterone (Selective)
Selective agonists (pK _i)	methyltrienolone (pEC ₅₀ < 5.0) [149], mibolerone (pIC ₅₀ 8.96) [124]	fluticasone, RU26988, RU28362	–	levonorgestrel [140], ORG2058
Selective antagonists (pK _i)	hydroxyflutamide (pEC ₅₀ 6.6) [149], PF0998425 (pIC ₅₀ 7.1 – 7.5) [131], nilutamide (pIC ₅₀ 7.07 – 7.12) [141]	onapristone, ZK112993, mifepristone (pK _d 9.4) [125,139]	onapristone, RU28318, ZK112993, eplerenone (pIC ₅₀ 1.0) [121,127]	mifepristone, onapristone, ZK112993
Radioligands (K _d)	[³ H]dihydrotestosterone (Agonist), [³ H]mibolerone (Agonist), [³ H]R1881 (Agonist)	[³ H]dexamethasone (Agonist)	[³ H]aldosterone (Agonist) (3x10 ⁻¹⁰ – 4x10 ⁻¹⁰ M - Rat) [123,144]	[³ H]ORG2058 (Agonist)

Comments: [³H]dexamethasone also binds to MR *in vitro*. PR antagonists have been suggested to subdivide into Type I (e.g. onapristone) and Type II (e.g. ZK112993) groups. These groups appear to promote binding of PR to DNA with different efficacies and evoke distinct conformational changes in the receptor, leading to a transcription-neutral complex [122,129]. Mutations in AR underlie testicular feminization and androgen insensitivity syndromes, spinal and bulbar muscular atrophy (Kennedy's disease).

Further reading

- Blaustein JD. (2008) Progesterone and progestin receptors in the brain: the neglected ones. *Endocrinology* **149**: 2737–2738. [PMID:18436712]
- Briet M, Schiffrin EL. (2010) Aldosterone: effects on the kidney and cardiovascular system. *Nat Rev Nephrol* **6**: 261–273. [PMID:20234356]
- Brinton RD. (2009) Estrogen-induced plasticity from cells to circuits: predictions for cognitive function. *Trends Pharmacol Sci* **30**: 212–222. [PMID:19299024]
- Callewaert F, Boonen S, Vanderschueren D. (2010) Sex steroids and the male skeleton: a tale of two hormones. *Trends Endocrinol Metab* **21**: 89–95. [PMID:19837603]
- Chen Y, Sawyers CL, Scher HI. (2008) Targeting the androgen receptor pathway in prostate cancer. *Curr Opin Pharmacol* **8**: 440–448. [PMID:18674639]
- Dahlman-Wright K, Cavaillès V, Fuqua SA, Jordan VC, Katzenellenbogen JA, Korach KS, Maggi A, Muramatsu M, Parker MG, Gustafsson JA. (2006) International Union of Pharmacology. LXIV. Estrogen receptors. *Pharmacol Rev* **58**: 773–781. [PMID:17132854]
- Feldman RD, Gros R. (2011) Unraveling the mechanisms underlying the rapid vascular effects of steroids: sorting out the receptors and the pathways. *Br J Pharmacol* **163**: 1163–1169. [PMID:21545416]
- Gross KL, Cidlowski JA. (2008) Tissue-specific glucocorticoid action: a family affair. *Trends Endocrinol Metab* **19**: 331–339. [PMID:18805703]
- Hadoke PW, Iqbal J, Walker BR. (2009) Therapeutic manipulation of glucocorticoid metabolism in cardiovascular disease. *Br J Pharmacol* **156**: 689–712. [PMID:19239478]
- Handelsman DJ. (2008) Indirect androgen doping by oestrogen blockade in sports. *Br J Pharmacol* **154**: 598–605. [PMID:18500381]
- Joëls M, Karst H, DeRijk R, de Kloet ER. (2008) The coming out of the brain mineralocorticoid receptor. *Trends Neurosci* **31**: 1–7. [PMID:18063498]
- Kellner M, Wiedemann K. (2008) Mineralocorticoid receptors in brain, in health and disease: possibilities for new pharmacotherapy. *Eur J Pharmacol* **583**: 372–378. [PMID:18272147]
- Kerkhofs S, Denayer S, Haelens A, Claessens F. (2009) Androgen receptor knockout and knock-in mouse models. *J Mol Endocrinol* **42**: 11–17. [PMID:18923000]
- Lange CA, Gioeli D, Hammes SR, Marker PC. (2007) Integration of rapid signaling events with steroid hormone receptor action in breast and prostate cancer. *Annu Rev Physiol* **69**: 171–199. [PMID:17037979]
- Levin ER. (2008) Rapid signaling by steroid receptors. *Am J Physiol Regul Integr Comp Physiol* **295**: R1425–R1430. [PMID:18784332]
- Lu NZ, Wardell SE, Burnstein KL, Defranco D, Fuller PJ, Giguere V, Hochberg RB, McKay L, Renoir JM, Weigel NL, Wilson EM, McDonnell DP, Cidlowski JA. (2006) International Union of Pharmacology. LXV. The pharmacology and classification of the nuclear receptor superfamily:



- glucocorticoid, mineralocorticoid, progesterone, and androgen receptors. *Pharmacol Rev* **58**: 782–797. [PMID:17132855]
- Matsumoto T, Shiina H, Kawano H, Sato T, Kato S. (2008) Androgen receptor functions in male and female physiology. *J Steroid Biochem Mol Biol* **109**: 236–241. [PMID:18434134]
- Mohler ML, Bohl CE, Jones A, Coss CC, Narayanan R, He Y, Hwang DJ, Dalton JT, Miller DD. (2009) Nonsteroidal selective androgen receptor modulators (SARMs): dissociating the anabolic and androgenic activities of the androgen receptor for therapeutic benefit. *J Med Chem* **52**: 3597–3617. [PMID:19432422]
- Pippal JB, Fuller PJ. (2008) Structure-function relationships in the mineralocorticoid receptor. *J Mol Endocrinol* **41**: 405–413. [PMID:18805937]
- Prossnitz ER, Arterburn JB, Smith HO, Oprea TI, Sklar LA, Hathaway HJ. (2008) Estrogen signaling through the transmembrane G protein-coupled receptor GPR30. *Annu Rev Physiol* **70**: 165–190. [PMID:18271749]
- Schumacher M, Sitruk-Ware R, De Nicola AF. (2008) Progesterone and progestins: neuroprotection and myelin repair. *Curr Opin Pharmacol* **8**: 740–746. [PMID:18929681]
- Simons SS. (2010) Glucocorticoid receptor cofactors as therapeutic targets. *Curr Opin Pharmacol* **10**: 613–619. [PMID:20801081]
- Taplin ME. (2007) Drug insight: role of the androgen receptor in the development and progression of prostate cancer. *Nat Clin Pract Oncol* **4**: 236–244. [PMID:17392714]
- Weiser MJ, Foradori CD, Handa RJ. (2008) Estrogen receptor beta in the brain: from form to function. *Brain Res Rev* **57**: 309–320. [PMID:17662459]
- Woolley CS. (2007) Acute effects of estrogen on neuronal physiology. *Annu Rev Pharmacol Toxicol* **47**: 657–680. [PMID:16918306]



References

1. Bergh JJ et al. (2005) *Endocrinology* **146**: 2864–2871. [PMID:15802494]
2. Chiellini G et al. (1998) *Chem Biol* **5**: 299–306. [PMID:9653548]
3. Flamant F et al. (2006) *Pharmacol Rev* **58**: 705–711. [PMID:17132849]
4. Nguyen NH et al. (2002) *J Med Chem* **45**: 3310–3320. [PMID:12109914]
5. Sharma SK et al. (1977) *Proc Natl Acad Sci USA* **74**: 3365–3369. [PMID:269396]
6. Apfel C et al. (1992) *Proc Natl Acad Sci USA* **89**: 7129–7133. [PMID:1323127]
7. Delescluse C et al. (1991) *Mol Pharmacol* **40**: 556–562. [PMID:1656191]
8. Germain P et al. (2006) *Pharmacol Rev* **58**: 712–725. [PMID:17132850]
9. Germain P et al. (2009) *Chem Biol* **16**: 479–489. [PMID:19477412]
10. Géhin M et al. (1999) *Chem Biol* **6**: 519–529. [PMID:10421757]
11. Kagechika H et al. (1988) *J Med Chem* **31**: 2182–2192. [PMID:3184125]
12. Keidel S et al. (1994) *Mol Cell Biol* **14**: 287–298. [PMID:8264595]
13. Le Q et al. (2000) *Oncogene* **19**: 1457–1465. [PMID:10723137]
14. Li E. (1999) *Mol Cell Biochem* **192**: 105–108. [PMID:10331664]
15. Lund BW et al. (2005) *J Med Chem* **48**: 7517–7519. [PMID:16302793]
16. Martin B et al. (1992) *Skin Pharmacol* **5**: 57–65. [PMID:1315557]
17. Schupp M et al. (2007) *Mol Pharmacol* **71**: 1251–1257. [PMID:17290005]
18. Thacher SM et al. (2000) *Curr Pharm Des* **6**: 25–58. [PMID:10637371]
19. Berger J et al. (1996) *Endocrinology* **137**: 4189–4195. [PMID:8828476]
20. Bishop-Bailey D et al. (2000) *Br J Pharmacol* **131**: 651–654. [PMID:11030710]
21. Bishop-Bailey D, Wray J. (2003) *Prostaglandins Other Lipid Mediat* **71**: 1–22. [PMID:12749590]
22. Brown KK et al. (1999) *Diabetes* **48**: 1415–1424. [PMID:10389847]
23. Brown PJ et al. (2001) *Bioorg Med Chem Lett* **11**: 1225–1227. [PMID:11354382]
24. Doebber TW et al. (2004) *Biochem Biophys Res Commun* **318**: 323–328. [PMID:15120604]
25. Graham TL et al. (2005) *Atherosclerosis* **181**: 29–37. [PMID:15939051]
26. Guo Q et al. (2004) *Endocrinology* **145**: 1640–1648. [PMID:14701675]
27. Henke BR et al. (1998) *J Med Chem* **41**: 5020–5036. [PMID:9836620]
28. Kane CD et al. (2009) *Mol Pharmacol* **75**: 296–306. [PMID:18971326]
29. Lee G et al. (2002) *J Biol Chem* **277**: 19649–19657. [PMID:11877444]
30. Leesnitzer LM et al. (2002) *Biochemistry* **41**: 6640–6650. [PMID:12022867]
31. Lehmann JM et al. (1997) *J Biol Chem* **272**: 3406–3410. [PMID:9013583]
32. McIntyre TM et al. (2003) *Proc Natl Acad Sci USA* **100**: 131–136. [PMID:12502787]
33. Michalik L et al. (2006) *Pharmacol Rev* **58**: 726–741. [PMID:17132851]
34. Nakamuta M et al. (2002) *Cell Biol Int* **26**: 235–241. [PMID:11991651]
35. Oliver WR et al. (2001) *Proc Natl Acad Sci USA* **98**: 5306–5311. [PMID:11309497]
36. Rocchi S et al. (2001) *Mol Cell* **8**: 737–747. [PMID:11684010]
37. Sakamoto J et al. (2000) *Biochem Biophys Res Commun* **278**: 704–711. [PMID:11095972]
38. Shearer BG et al. (2008) *Mol Endocrinol* **22**: 523–529. [PMID:17975020]
39. Sznajdman ML et al. (2003) *Bioorg Med Chem Lett* **13**: 1517–1521. [PMID:12699745]
40. Wang Y et al. (2000) *Mol Endocrinol* **14**: 1550–1556. [PMID:11043571]
41. Willson TM et al. (2000) *J Med Chem* **43**: 527–550. [PMID:10691680]
42. Xu HE et al. (2002) *Nature* **415**: 813–817. [PMID:11845213]
43. Xu Y et al. (2004) *J Med Chem* **47**: 2422–2425. [PMID:15115385]
44. Young PW et al. (1998) *J Pharmacol Exp Ther* **284**: 751–759. [PMID:9454824]
45. Benoit G et al. (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
46. Grant D et al. (2010) *ACS Chem Biol* **5**: 925–932. [PMID:20677822]
47. Kojetin D et al. (2010) *ACS Chem Biol* [Epub ahead of print]. [PMID:21043485]
48. Raghuram S et al. (2007) *Nat Struct Mol Biol* **14**: 1207–1213. [PMID:18037887]
49. Yin L et al. (2007) *Science* **318**: 1786–1789. [PMID:18006707]
50. Benoit G et al. (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
51. Bitsch F et al. (2003) *Anal Biochem* **323**: 139–149. [PMID:14622968]
52. Kallen J et al. (2004) *J Biol Chem* **279**: 14033–14038. [PMID:14722907]
53. Kallen JA et al. (2002) *Structure* **10**: 1697–1707. [PMID:12467577]
54. Paravicini G et al. (1996) *Biochem Biophys Res Commun* **227**: 82–87. [PMID:8858107]
55. Stehlin-Gaon C et al. (2003) *Nat Struct Biol* **10**: 820–825. [PMID:12958591]
56. Wiesenberg I et al. (1995) *Nucleic Acids Res* **23**: 327–333. [PMID:7885826]
57. Collins JL et al. (2002) *J Med Chem* **45**: 1963–1966. [PMID:11985463]
58. Downes M et al. (2003) *Mol Cell* **11**: 1079–1092. [PMID:12718892]
59. Lehmann JM et al. (1997) *J Biol Chem* **272**: 3137–3140. [PMID:9013544]
60. Makishima M et al. (1999) *Science* **284**: 1362–1365. [PMID:10334993]
61. Maloney PR et al. (2000) *J Med Chem* **43**: 2971–2974. [PMID:10956205]
62. Moore DD et al. (2006) *Pharmacol Rev* **58**: 742–759. [PMID:17132852]
63. Otte K et al. (2003) *Mol Cell Biol* **23**: 864–872. [PMID:12529392]
64. Parks DJ et al. (1999) *Science* **284**: 1365–1368. [PMID:10334993]
65. Pellicciari R et al. (2002) *J Med Chem* **45**: 3569–3572. [PMID:12166927]
66. Repa JJ et al. (2000) *Science* **289**: 1524–1529. [PMID:10968783]
67. Wu J et al. (2002) *Mol Endocrinol* **16**: 1590–1597. [PMID:12089353]
68. Bishop JE et al. (1994) *J Bone Miner Res* **9**: 1277–1288. [PMID:7976510]
69. Blumberg B et al. (1998) *Genes Dev* **12**: 3195–3205. [PMID:9784494]
70. Colston KW et al. (1992) *Biochem Pharmacol* **44**: 2273–2280. [PMID:1472092]
71. Erben RG et al. (2002) *Mol Endocrinol* **16**: 1524–1537. [PMID:12089348]
72. Fujishima T et al. (2003) *Bioorg Med Chem* **11**: 3621–3631. [PMID:12901907]
73. Herdick M et al. (2000) *Chem Biol* **7**: 885–894. [PMID:11094341]
74. Jones SA et al. (2000) *Mol Endocrinol* **14**: 27–39. [PMID:10628745]
75. Kanno Y et al. (2013) *J Toxicol Sci* **38**: 309–315. [PMID:23665929]
76. Lehmann JM et al. (1998) *J Clin Invest* **102**: 1016–1023. [PMID:9727070]
77. Maglich JM et al. (2003) *J Biol Chem* **278**: 17277–17283. [PMID:12611900]
78. Miura D et al. (1999) *J Biol Chem* **274**: 16392–16399. [PMID:10347199]
79. Moore DD et al. (2006) *Pharmacol Rev* **58**: 742–759. [PMID:17132852]
80. Moore LB et al. (2000) *Proc Natl Acad Sci USA* **97**: 7500–7502. [PMID:10852961]
81. Moore LB et al. (2000) *J Biol Chem* **275**: 15122–15127. [PMID:10748001]
82. Tzameli I et al. (2000) *Mol Cell Biol* **20**: 2951–2958. [PMID:10757780]
83. Wentworth JM et al. (2000) *J Endocrinol* **166**: R11–R16. [PMID:10974665]
84. Wiberg K et al. (1995) *Bone* **17**: 391–395. [PMID:8573413]
85. Benoit G et al. (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
86. Kiselyuk A et al. (2012) *Chem Biol* **19**: 806–818. [PMID:22840769]
87. Yuan X et al. (2009) *PLoS ONE* **4**: e5609. [ePMID:19440305]
88. Chambon P. (1996) *FASEB J* **10**: 940–954. [PMID:8801176]
89. Ebisawa M et al. (1999) *Chem Pharm Bull* **47**: 1778–1786. [PMID:10748721]
90. Germain P et al. (2006) *Pharmacol Rev* **58**: 760–772. [PMID:17132853]
91. Germain P et al. (2002) *Nature* **415**: 187–192. [PMID:11805839]
92. Mangelsdorf DJ, Evans RM. (1995) *Cell* **83**: 841–850. [PMID:8521508]
93. Nahoum V et al. (2007) *Proc Natl Acad Sci USA* **104**: 17323–17328. [PMID:17947383]
94. Benoit G et al. (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]



95. Shyr CR *et al.* (2002) *Mol Cell Biol* **22**: 4661–4666. [PMID:12052874]
96. Zhou XE *et al.* (2011) *J Biol Chem* **286**: 2877–2885. [PMID:21068381]
97. Benoit G *et al.* (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
98. Land PW, Monaghan AP. (2003) *Cereb Cortex* **13**: 921–931. [PMID:12902391]
99. Monaghan AP *et al.* (1997) *Nature* **390**: 515–517. [PMID:9394001]
100. Benoit G *et al.* (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
101. Pereira FA *et al.* (1999) *Genes Dev* **13**: 1037–1049. [PMID:10215630]
102. Qiu Y *et al.* (1997) *Genes Dev* **11**: 1925–1937. [PMID:9271116]
103. Warnecke M *et al.* (2005) *Genes Dev* **19**: 614–625. [PMID:15741322]
104. Benoit G *et al.* (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
105. Suetsugu M *et al.* (2003) *Mol Cancer Res* **1**: 981–991. [PMID:14638870]
106. Willy PJ *et al.* (2004) *Proc Natl Acad Sci USA* **101**: 8912–8917. [PMID:15184675]
107. Yu DD, Forman BM. (2005) *Bioorg Med Chem Lett* **15**: 1311–1313. [PMID:15713377]
108. Zuercher WJ *et al.* (2005) *J Med Chem* **48**: 3107–3109. [PMID:15857113]
109. Baker KD *et al.* (2003) *Cell* **113**: 731–742. [PMID:12809604]
110. Benoit G *et al.* (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
111. Flaig R *et al.* (2005) *J Biol Chem* **280**: 19250–19258. [PMID:15716272]
112. Wang Z *et al.* (2003) *Nature* **423**: 555–560. [PMID:12774125]
113. Zhan Y *et al.* (2008) *Nat Chem Biol* **4**: 548–556. [PMID:18690216]
114. Benoit G *et al.* (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
115. Del Tredici AL *et al.* (2008) *Mol Pharmacol* **73**: 900–908. [PMID:18055761]
116. Madoux F *et al.* (2008) *Mol Pharmacol* **73**: 1776–1784. [PMID:18334597]
117. Benoit G *et al.* (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
118. Benoit G *et al.* (2006) *Pharmacol Rev* **58**: 798–836. [PMID:17132856]
119. Compton DR *et al.* (2004) *J Med Chem* **47**: 5872–5893. [PMID:15537344]
120. Dahlman-Wright K *et al.* (2006) *Pharmacol Rev* **58**: 773–781. [PMID:17132854]
121. de Gasparo M *et al.* (1987) *J Pharmacol Exp Ther* **240**: 650–656. [PMID:2949071]
122. Gass EK *et al.* (1998) *Endocrinology* **139**: 1905–1919. [PMID:9528977]
123. Ge RS *et al.* (2005) *Mol Cell Endocrinol* **243**: 35–42. [PMID:16188378]
124. Giwercman A *et al.* (2000) *J Clin Endocrinol Metab* **85**: 2253–2259. [PMID:10852459]
125. Heikinheimo O *et al.* (1987) *J Steroid Biochem* **26**: 279–284. [PMID:3560943]
126. Hellal-Levy C *et al.* (1999) *FEBS Lett* **464**: 9–13. [PMID:10611474]
127. Hu X *et al.* (2005) *Mini Rev Med Chem* **5**: 709–718. [PMID:16101407]
128. Kraichely DM *et al.* (2000) *Endocrinology* **141**: 3534–3545. [PMID:11014206]
129. Leonhardt SA *et al.* (1998) *Mol Endocrinol* **12**: 1914–1930. [PMID:9849965]
130. Levin ER. (2008) *Am J Physiol Regul Integr Comp Physiol* **295**: R1425–R1430. [PMID:18784332]
131. Li JJ *et al.* (2008) *J Med Chem* **51**: 7010–7014. [PMID:18921992]
132. Lu NZ *et al.* (2006) *Pharmacol Rev* **58**: 782–797. [PMID:17132855]
133. Malamas MS *et al.* (2004) *J Med Chem* **47**: 5021–5040. [PMID:15456246]
134. Meyers MJ *et al.* (1999) *J Med Chem* **42**: 2456–2468. [PMID:10395487]
135. Meyers MJ *et al.* (2001) *J Med Chem* **44**: 4230–4251. [PMID:11708925]
136. Ning M *et al.* (2007) *Br J Pharmacol* **150**: 19–28. [PMID:17115070]
137. Prossnitz ER *et al.* (2008) *Annu Rev Physiol* **70**: 165–190. [PMID:18271749]
138. Prossnitz ER, Maggiolini M. (2009) *Mol Cell Endocrinol* **308**: 1–2. [PMID:19389460]
139. Rupprecht R *et al.* (1993) *Eur J Pharmacol* **247**: 145–154. [PMID:8282004]
140. Schindler AE *et al.* (2003) *Maturitas* **46**: S7–S16. [PMID:14670641]
141. Simard J *et al.* (1997) *Urology* **49**: 580–586; discussion 586–9. [PMID:9111629]
142. Smith JL *et al.* (2008) *Steroids* **73**: 1160–1173. [PMID:18603275]
143. Stauffer SR *et al.* (2000) *J Med Chem* **43**: 4934–4947. [PMID:11150164]
144. Stephenson G *et al.* (1984) *Am J Physiol* **246** (2 Pt 2): F227–F233. [PMID:6320679]
145. Sun J *et al.* (2002) *Endocrinology* **143**: 941–947. [PMID:11861516]
146. Sun J *et al.* (1999) *Endocrinology* **140**: 800–804. [PMID:9927308]
147. Tilley WD *et al.* (1989) *Proc Natl Acad Sci USA* **86**: 327–331. [PMID:2911578]
148. Vegeto E *et al.* (1993) *Mol Endocrinol* **7**: 1244–1255. [PMID:8264658]
149. Wang LG *et al.* (1998) *Biochem Pharmacol* **55**: 1427–1433. [PMID:10076535]
150. Yu DD, Forman BM. (2005) *Bioorg Med Chem Lett* **15**: 1311–1313. [PMID:15713377]
151. Zhu Y *et al.* (2003) *Proc Natl Acad Sci USA* **100**: 2237–2242. [PMID:12601167]
152. Zhu Y *et al.* (2003) *Proc Natl Acad Sci USA* **100**: 2231–2236. [PMID:12574519]

