



Palladium-catalyzed Exchange of Allylic Groups of Ethers and Esters with Carboxylic Anhydrides

Kuniyuki TAKAHASHI*, Go HATA, and Akihisa MIYAKE
Basic Research Laboratories, Toray Industries, Inc., Tebira, Kamakura

(Received October 12, 1972)

Previously we have reported^{1,2)} the palladium-catalyzed exchange reaction of allylic groups of ethers and esters with active-hydrogen compounds. In the course of the studies, it has been found that the reaction of the allylic compounds (I) with carboxylic anhydrides in the presence of palladium catalysts also gives the exchanged products (II and III) of allylic groups.



R = C₆H₅, CH₃CO, CH₃CH₂CO
R¹ = H, CH₂CH₂CH₂CH=CH₂
R² = H, CH₂CH₂CH₂CH=CH₂
R³ = CH₃, C₆H₅

An active and easily available catalyst is prepared by mixing palladium acetate and triphenylphosphine. Tetrakis(triphenylphosphine)palladium(0) is also an effective catalyst. However, the catalytic activity of the combination of dichlorobis(triphenylphosphine)-palladium (II) and sodium phenoxide is much smaller than that of the above two catalysts. Some platinum catalysts also showed the catalytic activity, but it was smaller than that of palladium catalysts.

The results on the reaction of allylic ethers and

TABLE 1. REACTIONS OF ALLYLIC ETHERS AND ESTERS WITH CARBOXYLIC ANHYDRIDES^{a)}

Anhydride (mol)	Allylic compound (mol)	Catalyst (mmol)	Time (hr)	Products ^{e)} and yields ^{f)} (%)
(CH ₃ CO) ₂ O	C ₆ H ₅ OC ₈ H ₁₃ ^{b)}	Pd(OCOCH ₃) ₂ -Ph ₃ P	40	CH ₃ CO ₂ X, 57 CH ₃ CO ₂ Y, 24
(CH ₃ CO) ₂ O	C ₆ H ₅ OC ₈ H ₁₃ ^{b)}	PdCl ₂ (Ph ₃ P) ₂ -PhONa	40	CH ₃ CO ₂ X, 18 CH ₃ CO ₂ Y, 8
(CH ₃ CO) ₂ O	C ₆ H ₅ OC ₈ H ₁₃ ^{b)}	Pd(Ph ₃ P) ₄	40	CH ₃ CO ₂ X, 36 CH ₃ CO ₂ Y, 15
(CH ₃ CO) ₂ O	C ₆ H ₅ OC ₈ H ₁₃ ^{b)}	PtCl ₂ (Ph ₃ P) ₂ -PhONa	40	CH ₃ CO ₂ X, 10 CH ₃ CO ₂ Y, 2
(CH ₃ CO) ₂ O	C ₆ H ₅ OC ₈ H ₁₃ ^{b)}	Pt(Ph ₃ P) ₄	40	CH ₃ CO ₂ X, 33 CH ₃ CO ₂ Y, 14
(C ₆ H ₅ CO) ₂ O	C ₆ H ₅ OC ₈ H ₁₃ ^{b)}	Pd(OCOCH ₃) ₂ -Ph ₃ P	40	C ₆ H ₅ CO ₂ X, 42 C ₆ H ₅ CO ₂ Y, 14
(C ₆ H ₅ CO) ₂ O	C ₆ H ₅ OCH ₂ CH=CH ₂ [*]	Pd(OCOCH ₃) ₂ -Ph ₃ P	40	C ₆ H ₅ CO ₂ CH ₂ CH=CH ₂ , 17
(CH ₃ CO) ₂ O	C ₂ H ₅ CO ₂ C ₈ H ₁₃ ^{e)}	Pd(OCOCH ₃) ₂ -Ph ₃ P	16	CH ₃ CO ₂ X, 55 CH ₃ CO ₂ Y, 22
(C ₆ H ₅ CO) ₂ O	CH ₃ CO ₂ C ₈ H ₁₃ ^{d)}	Pd(OCOCH ₃) ₂ -Ph ₃ P	16	C ₆ H ₅ CO ₂ X, 50 C ₆ H ₅ CO ₂ Y, 25
(C ₆ H ₅ CO) ₂ O	CH ₃ CO ₂ CH ₂ CH=CH ₂	Pd(OCOCH ₃) ₂ -Ph ₃ P	16	C ₆ H ₅ CO ₂ CH ₂ CH=CH ₂ , 33

a) Reaction conditions: anhydride 0.05 mol, allylic compound 0.025 mol (* 0.05 mol), palladium and platinum complex 0.05 mmol, Ph₃P 0.2 mmol, PhONa 0.5 mmol, reaction temperature 85°C, dimethylformamide 5 ml.

b) 1-Phenoxy-2,7-octadiene containing 1.3% of 3-phenoxy-1,7-octadiene.

c) A mixture of C₂H₅CO₂X(69%) and C₂H₅CO₂Y(31%).

d) A mixture of CH₃CO₂X(76%) and CH₃CO₂Y(24%).

e) X = -CH₂CH=CHCH₂CH₂CH₂CH=CH₂, Y = -CH(CH=CH₂)CH₂CH₂CH₂CH=CH₂.

f) Based on the allylic compound employed.

1) G. Hata, K. Takahashi, and A. Miyake, *Chem. Commun.*, **1970**, 1392.

2) K. Takahashi, A. Miyake, and G. Hata, *This Bulletin*, **45**, 230 (1972).

esters with carboxylic anhydrides are shown in Table 1.

Experimental

Reagents. Pd(OAc)₂,³⁾ PdCl₂(Ph₃P)₂,⁴⁾ Pd(Ph₃P)₄,⁴⁾ PtCl₂(Ph₃P)₂,⁵⁾ Pt(Ph₃P)₄,⁵⁾ 1-phenoxy-2,7-octadiene,⁶⁾

3) T. Matsuda, T. Mitsuyasu, and Y. Nakamura, *Kogyo Kagaku Zasshi*, **72**, 1751 (1969).

4) L. Malatesta and M. Angolleta, *J. Chem. Soc.*, **1957**, 1866.

5) L. Malatesta and C. Carielli, *ibid.*, **1958**, 2323.

6) E. J. Smutny, *J. Amer. Chem. Soc.*, **89**, 6793 (1968).

7) W. M. Iaver and W. F. Filbert, *ibid.*, **58**, 1388 (1936).

8) S. Hattori, H. Munakata, T. Suzuki, and N. Imaki, *Netherlands Patent*, 6816008 (1969).

1-phenoxy-2-propene,⁷⁾ octadienyl carboxylate⁸⁾ were prepared by previously reported methods. Carboxylic anhydrides and allyl acetate were purified by distillation. Dimethylformamide was dried on calcium hydride and purified by distillation.

Reaction Procedure. All the reactions were carried out under an argon atmosphere. In a 50 ml two necked flask, catalysts, reagents, and dimethylformamide were charged and the reaction mixture was stirred at 85°C. The reaction products were characterized by elemental analysis, molecular weight measurement, and IR and NMR spectral measurement or by comparison with authentic samples. The gas chromatography was used for the quantitative analysis of products.