

Formation of Ketone Diperoxides from Ozonation of *O*-Methyloximes


Yoshikatsu Ito,* Hiroaki YOKOYA, Yasutoshi UMEHARA, and Teruo MATSUURA

Department of Synthetic Chemistry, Faculty of Engineering, Kyoto University, Kyoto 606

(Received February 8, 1980)

Synopsis. The ozonation of the *O*-methyloximes has been investigated. Besides the corresponding ketones, ketone diperoxides, and *N*-methoxyamides were produced. The stereochemistry of the ketone diperoxides was studied by the NMR technique.

In our previous paper,¹⁾ it has been shown that the ozonation of valerophenone *O*-methyloxime (**1c**) in carbon tetrachloride at -20°C gives as the major products valerophenone diperoxide (**2c**) and *N*-methoxyamides **4c** and **5c**, in addition to the corresponding ketone **3c**. Erickson *et al.*, however, reported²⁾ that the ozonation of acetophenone *O*-methyloxime (**1a**) in dichloromethane at -78°C led to only the carbon-nitrogen double bond cleavage affording the corresponding ketone **3a**. This and our interest in a chemi-energized process occurring from **2c**³⁾ led us to study the ozonation of *O*-methyloximes in more detail.



Scheme 1.

Table 1 shows the results of the ozonation of *O*-methyloximes **1a—d** in carbon tetrachloride or dichloromethane at low temperatures. In addition to the corresponding ketones **3a—d**, the ketone diperoxides **2a—c** and *N*-methoxyamides **4a—c** and **5a—d** were isolated, as we found previously.¹⁾ In the case of **1d**, however, many by-products were also formed and only benzoic acid was characterizable.

The ketone diperoxides **2b** and **2c** were formed as a mixture of *cis* (**2b-c** and **2c-c**) and *trans* (**2b-t** and **2c-t**) isomers, while **2a** was obtained as a single isomer.⁴⁾ The yields of the *cis* and *trans* isomers were almost




Fig. 1. NMR spectra (100 MHz) of *cis*-propiophenone diperoxide (**2b-c**) and *cis*-valerophenone diperoxide (**2c-c**) at different temperatures.

equal, *i.e.* **2b-t/2b-c** = 1 and **2c-t/2c-c** = 1.2. From their NMR spectra at a number of temperatures the higher melting isomers were assigned to be *trans* and the lower melting isomers to be *cis*. Similarly, the stereochemistry of acetophenone diperoxide (**2a**) was assigned to be *trans*.

The chair-to-chair conformational isomerism of acetone diperoxide was studied by Murray and coworkers by the NMR technique.⁵⁾ The NMR spectra

TABLE 1. THE OZONATION OF *O*-METHYL OXIMES **1** AT LOW TEMPERATURE

<i>O</i> -Methyloxime 1 ($\text{M} \times 10^2$)	Solvent	O_3 passed (equiv)	Product (% yield)			
			2	3	4	5
1a (6.7)	$\text{CCl}_4^{\text{a)}$	6.3	6	39	16	9
1a (8.0)	$\text{CH}_2\text{Cl}_2^{\text{b)}$	57	10	43	11	8
1b (24.0)	$\text{CCl}_4^{\text{a)}$	6.5	13 ^{d)}	51	11	10
1c (5.5)	$\text{CCl}_4^{\text{a)}$	27	20 ^{d)}	34	13	16
1d (23.0)	$\text{CCl}_4^{\text{c)}$	5.6	0	6	0	10 ^{e)}

a) At -20°C . b) At -78°C . c) At 0°C . d) A combined yield of the *cis* and *trans* isomers. e) PhCO_2H (15%) as a by-product.

- 1) Y. Ito, M. Konishi, and T. Matsuura, *Photochem. Photobiol.*, **30**, 53 (1979).
- 2) R. E. Erickson, P. J. Andrusis, Jr., J. C. Collins, M. L. Lungle, and G. D. Mercer, *J. Org. Chem.*, **34**, 2961 (1969).
- 3) Y. Ito, T. Matsuura, and H. Yokoya, *J. Am. Chem. Soc.*, **101**, 4010 (1979).
- 4) The stereoselective formation of **2a** is probably due to the less bulkiness of the methyl group.
- 5) R. W. Murray, P. R. Story, and M. L. Kaplan, *J. Am. Chem. Soc.*, **88**, 526 (1966).
- 6) M. Schulz and K. Kirschke, "Organic Peroxides," ed by D. Swern, Wiley, New York (1972), Vol. 3, p. 73.
- 7) M. Schulz and K. Kirschke, *Adv. Heterocycl. Chem.*, **8**, 165 (1967).
- 8) For the reaction and generation of a carbonyl oxide, see P. S. Bailey, "Ozonation in Organic Chemistry," Academic Press, New York (1978).
- 9) A. H. Riebel, R. E. Erickson, C. J. Abshire, and P. S. Bailey, *J. Am. Chem. Soc.*, **82**, 1801 (1960).
- 10) G. J. Karabatsos and N. Hsi, *Tetrahedron*, **23**, 1079 (1967).
- 11) W. Dilthey, M. Inkel, and H. Stephan, *J. Prakt. Chem.*, **154**, 219 (1940).
- 12) P. M. Beart and A. D. Ward, *Aust. J. Chem.*, **27**, 1341 (1974).