

Synthesis of 2-(4-Azulenyl)ethanamine Derivatives as a Nonbenzenoid Analog of Biogenic Amine

Shinji KUROKAWA

Department of Chemistry, Faculty of Education, Saga University, Honjo-machi, Saga 840

(Received October 6, 1982)

2-(4-Azulenyl)ethanamine derivatives, nonbenzenoid analogs of the biologically active amine, were synthesized by the action of methyleneammonium salts on sodium 4-methylazulenide, sodium 4,6,8-trimethylazulenide, and sodium guaiazulenide. These compounds, as well as their hydrochlorides, were characterized by UV, IR, ^1H NMR, ^{13}C NMR, and MS data. The reaction of sodium guaiazulenide and *N*-ethylidenemethylamine also yielded the corresponding 2-(4-azulenyl)ethanamine derivatives, but the low yield of the product showed that the electrophilicity of azomethine carbon atom to 4-methylene carbanion was insufficient. The enzyme activity for some of those products was investigated. Negligible effect was found on prostaglandin 15-hydroxydehydrogenase, and considerable inhibition to cyclic AMP-phosphodiesterase, *in vitro*.

2-Phenylethanamine derivatives, together with their indole analogs, constitute an important class of biological active compounds related to biogenic amines.¹⁾ In studying azulene analogs of biological active amines, we previously reported the synthesis of some acyl and alkyl derivatives of 2-(1-azulenyl)ethanamine by aminoethylation of azulene or guaiazulene with *N*-acylaziridines or with aziridinium salts, as well as the effect of those compounds on enzyme activity.²⁾

Our study on azulene derivatives aminoethylated in the 5-membered ring of the azulene nucleus continued with those having aminoethyl side chains in the 7-membered ring. For this type of compounds we preliminarily reported the synthesis of 2-(4-azulenyl)ethanamine derivatives by the reaction of sodium guaiazulenide and methyleneammonium salts;³⁾ more details are given in this paper.

The present paper also deals with the synthesis of 2-(4-azulenyl)ethanamine derivatives by the action of *N*-methylenedimethylammonium iodide on sodium 4-methylazulenide or on sodium 4,6,8-trimethylazulenide.

Results and Discussion


Sodium *N*-methylanilide, used as the reagent, was conveniently prepared from sodium hydride and *N*-methylaniline in THF according to our modification.⁴⁾ This reagent reacted smoothly with 4-methylazulene (**1**)


to form a reddish-orange solution of sodium 4-methylazulenide (**2**).⁵⁾ Next **2** was mixed with *N*-methylenedimethylammonium iodide⁶⁾ at *ca.* -80°C , and left for 5 h to rise to room temperature.

From the hydrochloric acid extract of the reaction mixture, the intended *N,N*-dimethyl-2-(4-azulenyl)ethanamine, a major product, was isolated as its hydrochloride (**7**) (44%); this was easily converted to free amine (**3**) (90%) by the usual treatment with alkali. Both compounds (**3** and **7**) were characterized by spectral data.

In this reaction, three by-products: **4** (7%), **5** (2%), and **6** (2%), were also obtained. **5** and **6** are already known to occur in the aminomethylation of **1** with *N,N,N',N'*-tetramethylmethanethylenediamine.⁷⁾ The attempted aminomethylation of **1** with *N*-methylenedimethylammonium iodide was also found to yield **5** (7%) and **6** (13%). Consequently, the formation of **5** and **6** as by-products may be reasonably explained by aminomethylation of **1**, which coexists in the solution of **2** at equilibrium and increases in its ratio after the addition of an excess of methyleneammonium salt. Quaternary salt of *N*-methylaniline might act as a proton source.

The assigned position of the dimethylaminomethyl group of the other product, **4**, was indicated by low-field shift of 8-proton (0.15 ppm) of **4** relative to **3** in ^1H NMR. This conclusion is consistent with the observation that a dimethylaminomethyl group gives





rise a considerable low-field shift at *peri*-position, as is shown in 8-proton of **5** relative to **1** (0.31 ppm) and in 2'-methylene protons of **22** relative to **21** (0.38 ppm).

Moreover, the structure of **4** was established by the action of sodium *N*-methylanilide on **5** leading to sodium 1-dimethylaminomethyl-4-methylazulenide, followed by the reaction of it with *N*-methylenedimethylammonium iodide to yield **4** (6%). This process, even if possible, could not be a major course of formation of **4** in the reaction of **2** and methyleneammonium salt, judging from the low yields of **5** from **2** and of **4** from **5**. Thus most of **4** may be formed from the main product **3** by successive aminomethylation after aminoethylation of **2**.

Reaction of 4,6,8-trimethylazulene and sodium *N*-methylanilide at -15°C is known to give rise to **9** selectively, though proton abstraction from both 4(8)- and 6-methyl group is possible for this compound at higher temperatures.⁵⁾ Sodium 4,6,8-trimethylazulenide was thus prepared at -15°C and reacted with *N*-methylenedimethylammonium iodide, at first at *ca.* -80°C , followed by gradual completion at room temperature.

Hydrochloric acid treatment of the reaction products afforded **13** (13%), **14** (15%), and **16** (7%), all of which regenerated the corresponding 2-aminoethyl (**11** and **12**) and aminomethyl derivatives (**15**) almost quantitatively when treated with alkali.

Formation of **12** in this reaction is inconsistent with the observation in Ref. 5, but the same type of product from **10** is also noted in sodium 4,6,8-trimethylazulenide, prepared in THF at -15°C , when it was successively treated with carbon dioxide at -10°C .⁸⁾ The present reaction, in which the sodium 4,6,8-trimethylazulenide formed is used continuously for the next step, is thought to involve equilibrated species **8**, **9**, and **10** in the reaction mixtures, since *N*-methylaniline coexists also. Thus, **9** formed at first at -15°C may change into **10** via **8** in the following reaction with *N*-methylenedimethylammonium iodide to yield **12**. Another pathway

of direct conversion from **9** to **10** would require an ionic-attack of the nucleophile on anion **9** and seems improbable.⁸⁾

The yield ratio of **11** and **12** (46 : 54) is very close to that (45 : 55) of carbonation products at 4- and 6-methylene positions in sodium 4,6,8-trimethylazulenide prepared at ordinary temperature,⁸⁾ indicating that the equilibrium between **9** and **10** is faster than the reaction of those with methyleneammonium salt.

On the other hand, product **15** cannot be formed from **8**, because the electrophilic substitution of azulene is well known to occur at 1(3)-position,⁹⁾ and in fact, aminomethylation of **8** was found to yield 1-substituted and 1,3-disubstituted products.¹⁰⁾ Since the substitution at 2-position, along with that at 1(3)-position, is known in carbonation of azulenide anion of Meisenheimer type,⁸⁾ this product is thought to be formed from **9** and/or **10** by aminomethylation in a similar manner. So far as we know, it is the first case in methyleneazulenide-type anion that the electrophilic substitution was observed in the 5-membered ring of azulenide anion. Moreover, this fact also suggests that **5** and **6** may arise, at least in part, from **2** in the former reaction of **2** and methyleneammonium salt.

Similarly to the cases of **1** and **8**, **18** was prepared from **17** and allowed to react with some methyleneammonium salts.³⁾ The reaction of **18** and *N*-methylenedimethylammonium iodide was carried out as described above for **2** to give **21** (50%), **22** (7%), **19** (1%), and **20** (14%). By-products **20** and **22** may be produced by aminomethylation of **17** and/or **18**, or of **21**, as already discussed in cases of **1** and **8**. **19** is thought to be formed by nucleophilic attack of anilide anion on the 3-methylene position of conjugate acid of **20** ($-\text{CH}_2-\text{NH}^+(\text{CH}_3)_2$), as described before.³⁾

By virtually the same procedure as described above, the reaction of **18** with *N*-isopropylidenedimethylammonium perchlorate, *N*-isopropylidenepyrrolidinium perchlorate, or with *N*-(α -methylbenzylidene)pyrrolidinium perchlorate¹¹⁾ at -20°C or at room temperature yielded




TABLE 1. INHIBITION OF CYCLIC AMP-PHOSPHODIESTERASE (a) AND PROSTAGLANDIN 15-HYDROXYDEHYDROGENASE (b) BY 2-(4-AZULENYL)ETHANAMINE DERIVATIVES (ADDITION OF 20 $\mu\text{g cm}^{-3}$ IN EACH)

Compound	Inhibition/%	
	a	b
21	31.7	0
22	39.7	0
24	49.2	4.1
25	45.3	9.2
26	26.4	10.3

the corresponding 2-(4-azulenyl)ethanamine derivatives **24** (64%), **25** (62%), and **26** (54%). No appreciable amount of by-product was found in these cases. Of these 2-aminoethyl derivatives, **21** and **24**—**26** were converted to crystalline hydrochlorides **27** (50%), **28** (65%), **29** (51%), and **30** (28%), respectively. Both free amines and hydrochlorides were characterized by spectral data.

To clarify the reactivity of **18** further, the reaction of **18** and *N*-ethylidenemethylamine¹² was also tried and found to yield the corresponding amine, **23** (8%). However, the low yield of **23** is thought to show that the electrophilicity of azomethine carbon atom is rather insufficient to allow a smooth reaction of it with the methylene carbanion of **18**.

An additional investigation was undertaken on the effect of 2-(4-azulenyl)ethanamine derivatives (**21**—**26**) for enzyme activity, *in vitro*.¹³ As is observed in Table 1, considerable activities are found in inhibitions of cyclic AMP-phosphodiesterase in all compounds and especially in **24** and **25**, though less active than *N*-butyl-2-(5-isopropyl-3,8-dimethyl-1-azulenyl)ethanamine (69.9%).²

Different from the amide or primary amine derivatives of 2-(1-azulenyl)ethanamine previously reported,² these tertiary amine derivatives (**21**—**26**) had almost a negligible effect on prostaglandin 15-hydroxydehydrogenase. This observation well accords with that about tertiary amine derivatives of 2-(1-azulenyl)ethanamine, in which no appreciable effect was found on the inhibi-

tion by prostaglandin I₂ of ADP-induced aggregation of blood platelets.¹⁴

Experimental

All the melting points are uncorrected. Spectral data were recorded on the following instruments: UV, Hitachi 624 digital spectrometer with a Hitachi 056 recorder; IR, Hitachi 260-10 spectrometer; ¹H NMR, JEOL MH-100; ¹³C NMR, JNM-FX 100; MS, JEOL-D300 at 70 eV.

Reaction of Sodium 4-Methylazulene (2) and N-Methylenedimethylammonium Iodide.

Under dry nitrogen, a stirred solution of sodium *N*-methylanilide, prepared from 200 mg (5.1 mmol) of 61% sodium hydride and 1.81 g (16.9 mmol) of *N*-methylaniline in 20 cm³ of THF according to our modified procedure,⁴ was mixed with another solution of 710 mg (4.99 mmol) of 4-methylazulene (**1**)¹⁵ in 5 cm³ of THF over a period of 15 min, and stirring was continued for 30 min more (violet color changed to reddish orange). To the stirred solution of **3** thus obtained was added, under Dry Ice-acetone cooling, 2.00 g (10.8 mmol) of *N*-methylenedimethylammonium iodide over a period of 20 min (washed in with 5 cm³ of THF); stirring was continued for 1 h with Dry Ice-acetone cooling and then for 5 h more, during which period the reaction mixture was gradually brought to room temperature. A facile color change from reddish orange to violet was observed at ca. -20°C. The reaction mixture was taken up into 200 cm³ of benzene, washed with 50% potassium carbonate solution (100 cm³ × 2), dried on sodium sulfate, and concentrated (reduced pressure) to give a violet oil. Successive alumina chromatography (elution with 9 : 1 benzene-ethyl acetate) of the oil gave 2.15 mg (0.3%) of unchanged 4-methylazulene and 637 mg of violet oil, which was again dissolved in 200 cm³ of benzene, extracted with 1 mol dm⁻³ hydrochloric acid (100 cm³ × 3), and the hydrochloric acid solution extracted with chloroform (100 cm³ × 10). The chloroform solution was dried (sodium sulfate), concentrated (reduced pressure), and chromatographed on a silica-gel column (elution with 8 : 2 chloroform-methanol) to give 521 mg (44%) of **7** as violet crystals, mp 228—230°C: UV (C₂H₅OH) 240 (log ϵ 4.41), 270 (sh, 4.55), 275 (sh, 4.66), 278 (4.68), 283 (4.67), 298 (3.78), 318 (sh, 3.48), 329 (3.62), 342 (3.72), 354 (3.24), 531 (sh, 2.51), 553 (sh, 2.59), 571 (2.65), 588 (sh, 2.61), 617 (2.57), 644 (sh, 2.33), and 677 nm (2.14); IR (CHCl₃) 2545, 2510,

2400, 2310 (NH⁺), and 1192 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ=2.44 (6H, s, N(CH₃)₂), 2.90 (2H, m, 1'-H₂), 3.48 (2H, m, 2'-H₂), 6.96–7.58 (3H, m, 5-, 6-, and 7-H), 7.36 (1H, d, *J*=4 Hz, 1-H), 7.50 (1H, d, *J*=4 Hz, 3-H), 7.82 (1H, t, *J*=4 Hz, 2-H), and 8.28 (1H, d, *J*=9 Hz, 8-H).

The hydrochloric acid solution, after completion of chloroform extraction, was made alkaline by addition of ice-cooled 50% potassium carbonate solution, and extracted with benzene to give 449 mg of violet oil after removal of the solvent. Alumina chromatography of this oil with THF gave 19.6 mg (2%) of **5**, 29.7 mg (2%) of **6**, and 92.0 mg (7%) of **4** as violet oils in each case. **5** and **6** were identical with those in Ref. 7, as well as with authentic specimen obtained by the reaction of **1** and *N*-methylenedimethylammonium iodide, in UV, IR, and ¹H NMR. These compounds were also characterized with MS data. **5**: MS *m/e* (rel intensity) 199 (M⁺; 18) and 155 (M⁺ - N(CH₃)₂; 100). **6**: MS *m/e* (rel intensity) 256 (M⁺; 17), 212 (M⁺ - N(CH₃)₂; 23), and 167 (M⁺ - 2N(CH₃)₂ - H; 100).

Product **4** was characterized as follows: UV (C₂H₅OH) 242 (log ε 4.28), 278 (sh, 4.49), 282 (4.54), 286 (4.53), 290 (sh, 4.49), 300 (sh, 3.98), 336 (3.55), 345 (3.64), 359 (3.38), 543 (sh, 2.48), 575 (2.76), 617 (2.51), and 677 nm (sh, 2.09); IR (CCl₄) 2870, 2775, 2720 (N(CH₃)₂), 2980, 2950, 2860, 1468, 1456, 1440, 1430, 1374 (CH₃ and CH₂), and 1040 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ=2.25 (6H, s, N(CH₃)₂), 2.35 (6H, s, N(CH₃)₂), 2.77 (2H, m, 1'-H₂), 3.39 (2H, m, 2'-H₂), 3.86 (2H, s, 1-CH₂), 7.04 (1H, t, *J*=10 Hz, 7-H), 7.18 (1H, d, *J*=10 Hz, 5-H), 7.43 (1H, d, *J*=4 Hz, 3-H), 7.47 (1H, m, 6-H), 7.78 (1H, d, *J*=4 Hz, 2-H), and 8.47 (1H, d, *J*=9.5 Hz, 8-H); MS *m/e* (rel intensity) 256 (M⁺; 37), 212 (M⁺ - N(CH₃)₂; 7), 198 (M⁺ - CH₂N(CH₃)₂; 3), 154 (M⁺ - CH₂N(CH₃)₂ - N(CH₃)₂; 20), and 58 ((CH₃)₂NCH₂⁺; 100).

Found: C, 79.80; H, 9.35; N, 10.46%. Calcd for C₁₇H₂₄N₂: C, 79.64; H, 9.44; N, 10.93%.

In order to obtain **3** from **7**, **7** was dissolved in a mixed solvent of benzene–water (100 cm³ each) and slowly treated with 200 cm³ of 50% potassium carbonate solution under ice-cooling and stirring. The benzene layer was separated, the aqueous layer was extracted with benzene (30 cm³ × 2), and the combined benzene extract, after being dried (sodium sulfate) and concentrated (reduced pressure), was chromatographed over alumina (elution with 98 : 2 benzene–methanol) to give 126 mg (90%) of **3** as a violet oil: UV (C₂H₅OH) 240 (log ε 4.36), 265 (sh, 4.35), 270 (sh, 4.49), 275 (sh, 4.60), 278 (4.62), 283 (4.61), 298 (3.70), 317 (sh, 3.34), 328 (3.51), 342 (3.64), 354 (3.05), 530 (sh, 2.48), 550 (sh, 2.56), 568 (2.61), 587 (sh, 2.58), 615 (2.55), 642 (sh, 2.29), and 675 nm (2.12); IR (CCl₄) 2820, 2770, 2720 (NCH₃), 2980, 2945, 2860, 1456, 1440, 1362 (CH₃ and CH₂), 1045, and 1037 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ=2.35 (6H, s, N(CH₃)₂), 2.77 (2H, m, 1'-H₂), 3.40 (2H, m, 2'-H₂), 7.06 (1H, t, *J*=10.5 Hz, 7-H), 7.15 (1H, d, *J*=10.5 Hz, 5-H), 7.43 (1H, d, *J*=4 Hz, 1-H), 7.52 (1H, m, 6-H), 7.58 (1H, d, *J*=4 Hz, 3-H), 7.91 (1H, t, *J*=4 Hz, 2-H), and 8.33 (1H, d, *J*=9.5 Hz, 8-H); ¹³C NMR (CDCl₃) δ=36.5 (t, 2'-C), 45.1 (q, N(CH₃)₂), 60.6 (t, 1'-C), 115.1 (d, 3-C), 118.8 (d, 1-C), 121.5 (d, 7-C), 125.8 (d, 5-C), 135.3 (d, 2-C), 136.3 (d, 6-C), 136.7 (d, 8-C), 137.4 (s, 10-C), 139.9 (s, 9-C), and 148.8 (s, 4-C) [cf. 4-methylazulene in Ref. 16]¹⁶; MS *m/e* (rel intensity) 199 (M⁺; 20), 141 (M⁺ - (CH₃)₂NCH₂⁺; 3), and 58 ((CH₃)₂NCH₂⁺; 100).

Found: C, 84.34; H, 8.51; N, 7.07%. Calcd for C₁₄H₁₇N: C, 84.37; H, 8.60; N, 7.03%.

Reaction of 4-Methylazulene (1) and N-Methylenedimethylammonium Iodide. To a stirred solution of 300 mg (2.11 mmol) of 4-methylazulene in 5 cm³ of dry dichloromethane was added 600 mg (3.24 mmol) of *N*-methylenedimethyl-

ammonium iodide, and the suspension was stirred for 43 h. The reaction mixture was diluted with 100 cm³ of water, made alkaline by addition of 50% potassium carbonate solution, and extracted with chloroform (60 cm³ × 3). The chloroform extract was dried (sodium sulfate), concentrated (reduced pressure), and taken up into 100 cm³ of benzene. This benzene solution was extracted with 1 mol dm⁻³ hydrochloric acid (35 cm³ × 3), and the hydrochloric acid solution was extracted with chloroform (50 cm³ × 4). The chloroform extract, after being dried (sodium sulfate) and concentrated (reduced pressure), was chromatographed over silica gel (elution with 8 : 2 chloroform–methanol) to give 37.0 mg (7%) of hydrochloride of **5** as dark violet crystals, mp 177–179 °C: UV (C₂H₅OH) 241 (log ε 4.42), 271 (sh, 4.47), 277 (sh, 4.62), 281 (4.67), 286 (4.66), 290 (sh, 4.57), 301 (3.98), 318 (sh, 3.45), 332 (3.60), 341 (3.68), 353 (3.40), 550 (2.70), 584 (sh, 2.66), 617 (sh, 2.45), 643 (2.28), and 680 nm (1.72); IR (CHCl₃) 2650, 2550, 2500, 2400, 2325 (NH⁺), and 1195 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ=2.74 (6H, s, N(CH₃)₂), 3.02 (3H, s, 4-CH₃), 4.70 (2H, s, CH₂), 6.54 (1H, broad s, NH), 7.48 (1H, t, *J*=10 Hz, 7-H), 7.53 (1H, d, *J*=10 Hz, 5-H), 7.68 (1H, d, *J*=4 Hz, 3-H), 7.92 (1H, t, *J*=10 Hz, 6-H), 8.34 (1H, d, *J*=4 Hz, 2-H), and 8.90 (1H, d, *J*=10 Hz, 8-H).

The hydrochloric acid layer, after completion of chloroform extraction, was made alkaline by addition of 200 cm³ of 50% potassium carbonate solution, and extracted with benzene (50 cm³ × 3). Benzene extract, after removal of the solvent (reduced pressure), was submitted to alumina chromatography (elution with 3 : 1 benzene–THF) to afford 68.0 mg (13%) of **6**, identical with that of Ref. 7 and the authentic specimen (described above) in its UV, IR, and ¹H NMR spectra.

In order to obtain **5**, its hydrochloride was suspended in 80 cm³ of benzene and slowly mixed with 200 cm³ of ice-cooled 50% potassium carbonate solution. This mixture was worked up as described for **3** from **7** to give, without chromatographic separation, 17.9 mg (90%) of **5**, identical with that of Ref. 7 and the authentic specimen (described above) in its UV, IR, and ¹H NMR spectra.

Reaction of Sodium 1-Dimethylaminomethyl-4-methylazulene and N-Methylenedimethylammonium Iodide. To a stirred solution of 0.114 mmol of sodium *N*-methylaniline in 4 cm³ of THF

was added, under dry nitrogen, another solution of 21.8 mg (0.110 mmol) of **5** in 1 cm³ of THF during a period of 15 min, and then, under Dry Ice–acetone cooling, 30 mg (0.162 mmol) of *N*-methylenedimethylammonium iodide (washed in with 1 cm³ of THF). Stirring was continued at ca. -80 °C for 1 h and then 5 h; after the first hour the reaction mixture was gradually brought to room temperature. The reaction mixture was taken up into 20 cm³ of benzene and extracted with 1 mol dm⁻³ hydrochloric acid (20 cm³ × 3). The hydrochloric acid extract was washed with benzene (20 cm³ × 3), made alkaline by addition of 100 cm³ of ice-cooled 50% potassium carbonate solution, and extracted with benzene (20 cm³ × 3) to give 19.6 mg of brown oil after removal of the solvent (reduced pressure). Preparative TLC on an alumina sheet (3 : 1 benzene–THF) of the oil afforded 0.56 mg (3%) of unchanged **5** and 1.76 mg (6%) of **4**, identical with authentic specimen in TLC, UV, and IR results.

Reaction of Sodium 4,6,8-Trimethylazulene and N-Methylenedimethylammonium Iodide. Under dry nitrogen, a stirred solution of 5.1 mmol of sodium *N*-methylaniline in 20 cm³ of THF was mixed at -15 °C with another solution of 850 mg (4.99 mmol) of **8** in 5 cm³ of THF (violet color changed to orange), and then, under Dry Ice–acetone cooling, with 2.00 g (10.8 mmol) of *N*-methylenedimethylammonium iodide (washed in with 5 cm³ of THF). The suspension was stirred for 1 h at ca. -80 °C and for 5 h more with a gradual rising

of the temperature. The reaction mixture was worked up as described in **2** to give a violet oil, which was taken up into 200 cm³ of benzene, extracted with 1 mol dm⁻³ hydrochloric acid (100 cm³ × 3); the hydrochloric acid layer was extracted with chloroform (100 cm³ × 6). The benzene solution, after being dried (sodium sulfate) and concentrated (reduced pressure), was chromatographed on a silica-gel column with hexane to afford 220 mg (26%) of unchanged **8**. The chloroform extract was also dried (sodium sulfate), concentrated (reduced pressure), and submitted to silica gel chromatography (elution with 8 : 2 chloroform-methanol) to allow separation into three fractions.

From the first eluate, 174 mg (13%) of **13** was obtained as a violet oil; UV (C₂H₅OH) 244 (log ϵ 4.48), 270 (sh, 4.32), 275 (sh, 4.50), 281 (sh, 4.67), 285 (4.73), 294 (sh, 6.67), 305 (sh, 4.00), 328 (sh, 3.62), 334 (3.70), 343 (3.74), 348 (3.80), 359 (sh, 2.73), 547 (2.78), 565 (sh, 2.76), 580 (sh, 2.73), and 635 nm (sh, 2.31); IR (CHCl₃) 2500, 2400, 2360, 2325 (NH⁺), and 1195 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ =2.40 (6H, s, N(CH₃)₂), 2.63 (3H, s, 6-CH₃), 2.84 (2H, m, 1'-H₂), 2.86 (3H, s, 8-CH₃), 3.42 (2H, m, 2'-H₂), 7.13 (1H, s, 7-H), 7.16 (1H, s, 5-H), 7.48 (1H, q, J =4 and 1.8 Hz, 1-H), 7.58 (1H, q, J =4 and 1.8 Hz, 3-H), and 7.80 (1H, t, J =4 Hz, 2-H).

From the second eluate, 201 mg (15%) of **14** was obtained as violet crystals; these showed no certain melting point but decomposed above 200 °C: UV (C₂H₅OH) 245 (log ϵ 4.43), 270 (sh, 4.26), 275 (sh, 4.46), 281 (sh, 4.63), 285 (4.69), 294 (sh, 4.61), 306 (sh, 3.90), 327 (sh, 3.69), 333 (3.75), 342 (sh, 3.78), 348 (3.84), 520 (sh, 2.53), 555 (2.64), 575 (sh, 2.61), 592 (sh, 2.56), and 650 nm (sh, 2.09); IR (CHCl₃) 2550, 2510, 2450, 2390, 2355, 2330, 2280 (NH⁺), and 1195 cm⁻¹ (C-N); ¹H NMR (CD₃OD) δ =2.88 (6H, s, N(CH₃)₂), 2.90 (6H, s, 4- and 8-CH₃), 3.33 (4H, m, 1'- and 2'-H₂), 7.13 (2H, s, 5- and 7-H), 7.36 (2H, d, J =4 Hz, 1- and 3-H), and 7.66 (1H, t, J =4 Hz, 2-H).

From the last fraction, 85.4 mg (7%) of **16** was obtained as violet crystals, mp 41–45 °C: UV (C₂H₅OH) 246 (log ϵ 4.53), 287 (4.79), 293 (4.79), 335 (4.00), 351 (4.04), 536 (2.72), 560 (sh, 2.70), 615 (sh, 2.33), and 624 nm (sh, 2.27); IR (CHCl₃) 2500, 2400, 2350, 2330 (NH⁺), and 1190 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ =2.17 (6H, s, N(CH₃)₂), 2.61 (3H, s, 6-CH₃), 2.86 (6H, s, 4- and 8-CH₃), 3.83 (2H, s, CH₂), 4.48 (1H, broad s, NH⁺), 7.08 (2H, s, 5- and 7-H), and 7.33 (2H, s, 1- and 3-H).

In order to obtain **11** from **13**, a solution of 51.6 mg (0.196 mmol) of **13** in 70 cm³ of benzene-methanol (98 : 2) was mixed dropwise with 200 cm³ of ice-cooled 50% potassium carbonate solution. After the mixture was worked up in the usual manner as described before, 44.5 mg (100%) of **11** was obtained as a violet oil: UV (C₂H₅OH) 244 (log ϵ 4.43), 271 (sh, 4.31), 275 (sh, 4.45), 281 (sh, 4.61), 285 (4.67), 290 (4.67), 294 (sh, 4.61), 305 (sh, 3.93), 322 (sh, 3.45), 328 (sh, 3.51), 334 (3.59), 344 (sh, 3.64), 349 (3.71), 359 (sh, 3.06), 547 (2.71), 564 (sh, 2.69), 581 (sh, 2.65), and 637 nm (sh, 2.20); IR (CCl₄) 2825, 2775, 2725 (N(CH₃)₂), 2980, 2950, 2860, 1467, 1458, 1438, 1372 (CH₂ and CH₃), 1045, and 1040 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ =2.37 (6H, s, N(CH₃)₂), 2.60 (3H, s, 6-CH₃), 2.78 (2H, m, 1'-H₂), 2.84 (3H, s, 8-CH₃), 3.38 (2H, m, 2'-H₂), 7.06 (1H, s, 7-H), 7.09 (1H, s, 5-H), 7.40 (1H, q, J =4 and 1.5 Hz, 1-H), 7.50 (1H, q, J =4 and 1.5 Hz, 3-H), and 7.73 (1H, t, J =4 Hz, 2-H); ¹³C NMR (CDCl₃) δ =25.0 (q, 8-CH₃), 28.7 (q, 6-CH₃), 37.3 (t, 2'-C), 45.2 (q, N(CH₃)₂), 61.1 (t, 1'-C), 115.5 (d, 3-C), 116.2 (d, 1-C), 126.7 (d, 5-C), 127.0 (d, 7-C), 132.6 (d, 2-C), 136.0 (s, 9- and 10-C), 145.4 (s, 8-C), 146.3 (s, 6-C), and 147.8 (s, 4-C) [cf. 4,6,8-trimethylazulene in Ref. 16]¹⁰; MS m/e (rel intensity) 227 (M⁺; 12), 169 (M⁺-(CH₃)₂NCH₂; 3), and 58 ((CH₃)₂NCH₂⁺; 100).

Found: C, 84.23; H, 9.21; N, 6.04%. Calcd for C₁₆H₂₁N: C, 84.53; H, 9.31; N, 6.16%.

Similar treatment of 53.4 mg (0.202 mmol) of **14** gave 43.9 mg (0.202 mmol) of **12** as a violet oil: UV (C₂H₅OH) 244 (log ϵ 4.51), 275 (sh, 4.53), 281 (4.68), 284 (4.74), 289 (4.73), 294 (sh, 4.66), 304 (sh, 4.05), 322 (sh, 3.50), 328 (sh, 3.57), 333 (3.64), 343 (sh, 3.69), 348 (3.75), 548 (2.69), 562 (sh, 2.67), 584 (sh, 2.62), and 638 nm (sh, 2.18); IR (CCl₄) 2825, 2780, 2725 (N(CH₃)₂), 2985, 2955, 1470, 1460, 1435, 1375 (CH₃ and CH₂), 1050, and 1042 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ =2.32 (6H, s, N(CH₃)₂), 2.62 (2H, m, 1'-H₂), 2.88 (6H, s, 4- and 8-CH₃), 2.94 (2H, m, 2'-H₂), 7.08 (2H, s, 5- and 7-H), 7.42 (2H, d, J =4 Hz, 1- and 3-H), and 7.74 (1H, t, J =4 Hz, 2-H); ¹³C NMR (CDCl₃) δ =25.0 (q, 4- and 8-CH₃), 41.2 (t, 2'-C), 45.4 (q, N(CH₃)₂), 62.4 (t, 1'-C), 116.0 (d, 1- and 3-C), 126.8 (d, 5- and 7-C), 132.7 (d, 2-C), 136.2 (s, 9- and 10-C), 145.7 (s, 4- and 8-C), and 148.2 (s, 6-C); MS m/e (rel intensity) 227 (M⁺; 5), 169 (M⁺-(CH₃)₂NCH₂; 1), and 58 ((CH₃)₂NCH₂⁺; 100).

Found: C, 84.27; H, 9.03; N, 6.11%. Calcd for C₁₆H₂₁N: C, 84.53; H, 9.31; N, 6.16%.

Similar treatment of 37.9 mg (0.144 mmol) of **16** afforded 32.8 mg (100%) of **15** as a violet oil: UV (C₂H₅OH) 246 (log ϵ 5.40), 287 (5.72), 293 (5.72), 335 (4.55), 351 (4.68), 537 (2.65), 560 (sh, 2.62), and 616 nm (sh, 2.20); IR (CCl₄) 2820, 2775, 2730 (N(CH₃)₂), 2980, 2945, 2930, 2860, 1455, 1445, 1376, 1357 (CH₃ and CH₂), 1040, and 1030 cm⁻¹ (C-N); ¹H NMR (CDCl₃) δ =2.35 (6H, s, N(CH₃)₂), 2.61 (3H, s, 6-CH₃), 2.87 (6H, s, 4- and 8-CH₃), 3.82 (2H, s, CH₂), 7.12 (2H, s, 5- and 7-H), and 7.37 (2H, s, 1- and 3-H); MS m/e (rel intensity) 227 (M⁺; 12), 184 (M⁺-CH₃NCH₂; 100), 183 (M⁺-(CH₃)₂N; 28), 169 (M⁺-(CH₃)₂NCH₂; 41), and 58 ((CH₃)₂NCH₂⁺; 13).

Found: C, 84.59; H, 9.30; N, 5.94%. Calcd for C₁₆H₂₁N: C, 84.53; H, 9.31; N, 6.16%.

Reaction of Sodium Guaiazulene (18) with Methylenammonium Salts or with N-Ethylidenemethylamine. A) *With N-Methylenedimethylammonium Iodide:*

To a stirred solution of 10 mmol of sodium guaiazulene (**18**) in 30 cm³ of THF [from 440 mg (10.2 mmol) of sodium hydride, 3.62 g (33.8 mmol) of *N*-methylaniline, and 1.99 g (10.0 mmol) of guaiazulene, as described in the preparation of **2**], under dry nitrogen and with Dry Ice-acetone cooling, was added 3.70 g (20 mmol) of *N*-methylenedimethylammonium iodide over a period of 20 min (washed with 5 cm³ of THF), and the suspension was stirred for 1 h at ca. -80 °C and for an additional 5 h with a gradual elevation of the temperature. The reaction mixture, after being worked up as described above for **2**, was chromatographed over alumina (elution with 9 : 1 hexane-ethyl acetate) and separated into four fractions. Further elution with ethanol afforded one more fraction.

Rechromatography of the first fraction on a silica-gel column with hexane gave 30.2 mg (2%) of unchanged guaiazulene. Successively the solvent was changed into benzene to give 32.2 mg (1%) of **19** as a blue oil: UV (C₂H₅OH) 248 (log ϵ 4.44), 283 (sh, 4.54), 290 (4.60), 307 (4.22), 341 (sh, 3.58), 354 (3.73), 371 (3.63), 584 (sh, 2.49), 625 (2.58), 676 (sh, 2.49), and 748 (sh, 2.05); IR (CCl₄) 3440 (NH), 1375 and 1360 cm⁻¹ (CH₃); ¹H NMR (CDCl₃)¹⁸ δ =1.32 (6H, d, J =7 Hz, CH(CH₃)₂), 2.60 (3H, s, 1-CH₃), 2.70 (3H, s, NCH₃), 2.84 (3H, s, 4-CH₃), 2.98 (1H, m, J =7 Hz, CH(CH₃)₂), 3.34 (1H, broad s, NH), 4.48 (2H, s, CH₂), 6.48 (2H, d, J =8 Hz, 3'- and 5'-H), 6.76 (1H, d, J =10 Hz, 5-H), 6.88 (2H, d, J =8 Hz, 2''- and 6''-H), 7.22 (1H, q, J =10 and 2 Hz, 6-H), 7.40 (1H, s, 2-H), and 8.08 (1H, d, J =2 Hz, 8-H); MS m/e (rel intensity) 317 (M⁺; 100), 302 (M⁺-CH₃; 57), and 120 (CH₃NHC₆H₄CH₂⁺; 29).

Found: C, 86.73; H, 8.53; N, 4.30%. Calcd for C₂₃H₂₇N:

C, 87.02; H, 8.57; N, 4.41%.

Rechromatography of the second fraction on an alumina column (elution with 98 : 2 hexane–ethyl acetate) gave 346 mg (14%) of **20** as blue crystals, mp 37–38 °C [Ref. 10; blue oil].¹⁰⁾ UV (cyclohexane) 215 (log ϵ 4.02), 247 (4.28), 282 (sh, 4.49), 288 (4.56), 292 (4.58), 305 (4.14), 339 (sh, 3.54), 347 (sh, 3.60), 353 (3.73), 371 (3.73), 568 (sh, 2.50), 595 (sh, 2.58), 613 (2.61), 640 (sh, 2.59), 667 (2.53), 708 (sh, 2.22), and 741 nm (2.08) [Ref. 10 (hexane); 613 (2.62), 662 (2.52), 714 (2.18), and 736 nm (2.04)]; IR (CCl₄) 2820, 2775, 2720 (N(CH₃)₂), 1460, 1435, and 1420 cm⁻¹ (CH₂); ¹H NMR (CDCl₃) δ =1.30 (6H, d, J =7 Hz, CH(CH₃)₂), 2.22 (6H, s, N(CH₃)₂), 2.60 (3H, s, 1-CH₃), 2.99 (1H, m, J =7 Hz, CH(CH₃)₂), 3.12 (3H, s, 4-CH₃), 3.80 (2H, s, 3-CH₂), 6.90 (1H, d, J =11 Hz, 5-H), 7.26 (1H, q, J =11 and 2 Hz, 6-H), 7.48 (1H, s, 2-H), and 8.10 (1H, d, J =2 Hz, 8-H); MS m/e (rel intensity) 255 (M⁺; 49) and 211 (M⁺–N(CH₃)₂; 100).

Rechromatography of the third fraction on an alumina column (elution with 9 : 1 hexane–ethyl acetate) yielded 1.28 g (50%) of **21** as a blue oil: UV (C₂H₅OH) 217 (log ϵ 4.01), 245 (4.35), 280 (sh, 4.52), 284 (4.57), 288 (4.56), 302 (3.98), 327 (sh, 3.23), 338 (sh, 3.43), 350 (3.60), 367 (3.45), 563 (sh, 2.53), 585 (sh, 2.60), 607 (2.65), 632 (sh, 2.61), 654 (2.58), and 725 nm (2.16); IR (CCl₄) 2820, 2772 (NCH₃), 2964, 2864, 1468, 1445, 1424, 1391, and 1373 cm⁻¹ (CH₃ and CH₂); ¹H NMR (CDCl₃) δ =1.31 (6H, d, J =7 Hz, CH(CH₃)₂), 2.28 (6H, s, N(CH₃)₂), 2.66 (3H, s, 1-CH₃), 2.76 (2H, m, 1'-H₂), 2.98 (1H, m, CH(CH₃)₂), 3.30 (2H, m, 2'-H₂), 6.98 (1H, d, J =10 Hz, 5-H), 7.36 (1H, q, J =10 and 1.6 Hz, 6-H), 7.40 (1H, d, J =4 Hz, 3-H), 7.68 (1H, d, J =4 Hz, 2-H), and 8.26 (1H, d, J =1.6 Hz, 8-H); ¹³C NMR (CDCl₃) δ =12.9 (q, 1-CH₃), 24.7 (q, CH(CH₃)₂), 36.3 (t, 2'-C), 38.1 (d, CH(CH₃)₂), 45.2 (q, N(CH₃)₂), 60.7 (t, 1'-C), 112.2 (d, 3-C), 124.7 (d, 5-C), 125.0 (s, 1-C), 133.0 (d, 8-C), 135.0 (d, 6-C), 136.0 (s, 10-C), 136.4 (d, 2-C), 137.3 (s, 9-C), 139.6 (s, 7-C), and 146.4 (s, 4-C) [cf. 5-guaiazulene in Ref. 17]¹⁷⁾; MS m/e (rel intensity) 255 (M⁺; 100), 211 (M⁺–(CH₃)₂N; 6), and 197 (M⁺–(CH₃)₂NCH₂; 6).

Found: C, 84.47; H, 9.61; N, 5.61%. Calcd for C₁₈H₂₅N: C, 84.65; H, 9.87; N, 5.48%.

The combined fourth and ethanol fractions were chromatographed on an alumina column (elution with 8 : 2 benzene–ethyl acetate) to afford 218 mg (7%) of **22** as dark blue prisms, mp 66–68 °C; UV (C₂H₅OH) 217 (sh, ϵ 4.09), 247 (4.34), 289 (4.56), 304 (sh, 4.12), 342 (sh, 3.61), 353 (3.72), 372 (3.69), 407 (sh, 2.09), 573 (sh, 2.55), 609 (2.64), 653 (sh, 2.57), and 727 nm (sh, 2.13); IR (CCl₄) 2820, 2770 (NCH₃), 2968, 2865, 1460, 1440, 1430, 1420, 1385, and 1370 cm⁻¹ (CH₃ and CH₂); ¹H NMR (CDCl₃) δ =1.34 (6H, d, J =7 Hz, CH(CH₃)₂), 2.23 (6H, s, N(CH₃)₂), 2.36 (6H, s, N(CH₃)₂), 2.61 (3H, s, 1-CH₃), 2.72 (2H, m, 1'-H₂), 3.02 (1H, m, J =7 Hz, CH(CH₃)₂), 3.68 (2H, m, 2'-H₂), 3.84 (2H, s, 3-CH₂), 7.03 (1H, d, J =11 Hz, 5-H), 7.38 (1H, q, J =11 and 2 Hz, 6-H), 7.56 (1H, s, 2-H), and 8.16 (1H, d, J =2 Hz, 8-H); MS m/e (rel intensity) 312 (M⁺; 3), 254 (M⁺–(CH₃)₂NCH₂; 9), and 58 ((CH₃)₂NCH₂⁺; 100).

Found: C, 80.35; H, 10.10; N, 8.81%. Calcd for C₂₁H₃₂N₂: C, 80.71; H, 10.32; N, 8.96%.

B) With N-Isopropylidenedimethylammonium Perchlorate: Reaction of 10 mmol of **18** and 3.71 g (20.0 mmol) of *N*-isopropylidenedimethylammonium perchlorate¹¹⁾ was carried out in a manner similar to *A* except that the reagents were reacted at room temperature for 2 h. Chromatography of the reaction mixture on an alumina column (elution with 8 : 2 benzene–chloroform) gave 746 mg of blue oil, which on rechromatography over silica gel with hexane afforded 597 mg (30%) of unchanged **17**. Successive blue eluate gave 2.11 g of blue

oil, which was also rechromatographed over alumina (elution with 9 : 1 hexane–ethyl acetate) to give 1.81 g (64%) of **24** as a blue oil: UV (C₂H₅OH) 247 (log ϵ 4.29), 282 (sh, 4.51), 285 (4.54), 289 (4.53), 305 (sh, 3.94), 340 (sh, 3.55), 351 (3.66), 368 (3.52), 566 (sh, 2.53), 588 (sh, 2.60), 609 (2.66), 634 (sh, 2.61), 659 (2.58), and 728 nm (2.14); IR (CCl₄) 2820, 2780, 2730 (NCH₃), 2960, 2925, 2870, 1460, 1385, 1365 (CH₃ and CH₂), and 1145 cm⁻¹ (C–N); ¹H NMR δ =1.01 (6H, s, 1'-C(CH₃)₂), 1.29 (6H, d, J =7 Hz, CH(CH₃)₂), 2.44 (6H, s, N(CH₃)₂), 2.64 (3H, s, 1-CH₃), 3.05 (1H, m, J =7 Hz, CH(CH₃)₂), 3.30 (2H, s, 2'-H₂), 7.08 (1H, d, J =11 Hz, 5-H), 7.40 (1H, d, J =4 Hz, 3-H), 7.46 (1H, q, J =11 and 2 Hz, 6-H), 7.70 (1H, d, J =4 Hz, 2-H), and 8.24 (1H, d, J =2.0 Hz, 8-H); MS m/e (rel intensity) 283 (M⁺; 0.7), 198 (M⁺–C₅H₁₁N; 15), and 86 (C₅H₁₂N⁺; 100).

Found: C, 84.64; H, 10.24; N, 4.41%. Calcd for C₂₀H₂₉N: C, 84.75; H, 10.31; N, 4.94%.

C) With N-Isopropylidenepyrrolidinium Perchlorate: Reaction of 10 mmol of **18** and 4.25 g (20.1 mmol) of *N*-isopropylidenepyrrolidinium perchlorate¹¹⁾ was carried out in a manner similar to *A*, except that the reaction mixture was stirred for 3 h at –20 °C (ice salt mixture) and an additional 1 h at room temperature. Elution of the reaction mixture on an alumina column with hexane gave 156 mg (8%) of unchanged **17**. Then the solvent was changed into hexane–ethyl acetate (9 : 1) to give 1.93 g (62%) of **25** as a blue oil: UV (C₂H₅OH) 218 (sh, log ϵ 4.00), 246 (4.31), 275 (sh, 4.37), 281 (sh, 4.47), 285 (4.50), 289 (4.48), 304 (sh, 3.92), 568 (sh, 2.49), 592 (2.57), 611 (2.61), 634 (sh, 2.58), 661 (2.55), and 729 nm (2.11); IR (CCl₄) 2968, 2940, 2916, 2875, 2814, 1468, 1453, 1440, 1423, 1390, and 1370 cm⁻¹ (CH₃ and CH₂); ¹H NMR (CDCl₃)¹⁸⁾ δ =1.06 (6H, s, 1'-C(CH₃)₂), 1.31 (6H, d, J =7 Hz, CH(CH₃)₂), 1.75 (4H, m, J =6 Hz, β - and γ -H₂), 2.64 (3H, s, 1-CH₃), 2.79 (4H, m, J =6 Hz, α - and δ -H₂), 3.00 (1H, m, J =7 Hz, CH(CH₃)₂), 3.32 (2H, s, 2'-H₂), 7.01 (1H, d, J =11 Hz, 5-H), 7.32 (1H, q, J =11 and 2 Hz, 6-H), 7.36 (1H, d, J =4 Hz, 3-H), 7.60 (1H, d, J =4 Hz, 2-H), and 8.16 (1H, d, J =2 Hz, 8-H); MS m/e (rel intensity) 309 (M⁺; 3), 197 (M⁺–(CH₃)₂CNC₄H₈; 4), 112 (C₄H₈NC(CH₃)₂⁺; 100), and 70 (C₄H₈N⁺; 37).

Found: C, 85.56; H, 10.09; N, 4.24%. Calcd for C₂₂H₃₁N: C, 85.38; H, 10.10; N, 4.53%.

D) With N-(α -Methylbenzylidene)pyrrolidinium Perchlorate: Reaction of 10 mmol of **18** and 5.00 g (18.3 mmol) of *N*-(α -methylbenzylidene)pyrrolidinium perchlorate was carried out in a manner similar to *C*. Elution of the reaction mixture on a silica-gel column with hexane afforded 399 mg (20%) of unchanged **17**. Then the solvent was changed into benzene–methanol (98 : 2) to give 2.02 g (54%) of **26** as a blue oil: UV (C₂H₅OH) 247 (log ϵ 4.32), 286 (4.51), 290 (sh, 4.49), 304 (sh, 3.93), 327 (sh, 3.40), 341 (sh, 3.51), 351 (3.61), 368 (3.50), 415 (2.27), 435 (2.26), 566 (sh, 2.49), 529 (sh, 2.62), 633 (sh, 2.58), 658 (2.55), and 730 nm (2.13); IR (CCl₄) 3110, 3090, 3070, 3035 (aromatic CH), 2975, 2880, 1465, 1450, 1440, 1423, 1390, 1380, 1370, and 1368 cm⁻¹ (CH₃ and CH₂); ¹H NMR (CDCl₃)¹⁸⁾ δ =1.23 (6H, d, J =7 Hz, CH(CH₃)₂), 1.24 (3H, s, 1'-C(CH₃)₂), 1.74 (4H, m, β - and γ -H₂), 2.60 (3H, s, 1-CH₃), 2.70 (4H, t, J =6 Hz, α - and δ -H₂), 2.90 (1H, m, J =7 Hz, CH(CH₃)₂), 3.31 (1H, d, J =12.5 Hz, 2'-H), 4.01 (1H, d, J =12.5 Hz, 2'-H), 6.03 (1H, d, J =11 Hz, 5-H), 6.85 (1H, q, J =11 and 2 Hz, 6-H), 7.10–7.38 (6H, m, 3-H and Ph), 7.52 (1H, d, J =4 Hz, 2-H), and 8.04 (1H, d, J =2 Hz, 8-H); ¹³C NMR (CDCl₃)¹⁸⁾ δ =13.0 (1-CH₃), 14.3 (1'-C(CH₃)₂), 24.1 (β - and γ -C), 24.6 (CH(CH₃)₂), 37.9 (CH(CH₃)₂), 46.1 (α - and δ -C), 49.7 (2'-C), 63.3 (1'-C), 113.1 (3-C), 124.4 (5-C), 126.2 (1-C), 126.7 (4'-C), 127.5 (2'', 3'', 5'', and 6''-C), 132.3 (8-C), 133.5 (6-C), 135.6 (2-C), 136.1 (10-C), 139.0 (9-C), 139.7 (7-C), 144.9 (1'-C), and 146.2 (4-C); MS m/e (rel intensi-

ty) 371 (M^+ ; 2), 174 ($C_6H_5C(CH_3)NC_4H_9^+$; 100), and 70 ($C_4H_9N^+$; 4).

Found: C, 87.17; H, 8.67; N, 3.35%. Calcd for $C_{27}H_{33}N$: C, 87.28; H, 8.95; N, 3.77%.

E) With N-Ethylidenemethylamine: Reaction of 10 mmol of **18** and 3.0 cm³ (40 mmol) of N-ethylidenemethylamine was carried out as described in B. Chromatography of the reaction mixture on an alumina column (elution with 8 : 2 benzene-methanol) yielded 4.08 g of blue oil, which on rechromatography (hexane-silica gel) afforded 1.74 g (88%) of unchanged **17**. The second blue eluate was rechromatographed on an alumina column (elution with 8 : 2 benzene-methanol) to afford 210 mg (8%) of **23** as a blue oil: UV (C_2H_5OH) 219 (log ϵ 4.02), 246 (4.36), 285 (4.58), 289 (4.58), 303 (3.99), 349 (3.62), 367 (3.49), 565 (sh, 2.55), 607 (2.66), 656 (2.59), and 725 nm (sh, 2.15); IR (CCl_4) 3330 (NH), 2790 (NCH_3), 2960, 2925, 2870, 1470, 1460, 1445, 1438, 1420, 1390, and 1370 cm⁻¹ (CH_3 and CH_2); ¹H NMR ($CDCl_3$) δ =1.12 (3H, d, J =6.3 Hz, 1'-C(CH_3)), 1.35 (6H, d, J =6.9 Hz, $CH(CH_3)_2$), 1.71 (1H, broad s, NH), 2.39 (3H, s, NCH_3), 2.64 (3H, s, 1- CH_3), 2.88-3.40 (4H, m; 1'-H, 2'-H₂, and $CH(CH_3)_2$), 6.96 (1H, d, J =10.3 Hz, 5-H), 7.28 (1H, d, J =3.4 Hz, 3-H), 7.36 (1H, q, J =10.3 and 1.7 Hz, 6-H), 7.59 (1H, d, J =3.4 Hz, 2-H), and 8.14 (1H, d, J =1.7 Hz, 8-H); MS m/e (rel intensity) 255 (M^+ ; 7), 198 ($M^+ - C_3H_5N$; 32), and 58 ($C_3H_5N^+$; 100).

Found: C, 84.70; H, 9.56; N, 5.62%. Calcd for $C_{18}H_{25}N$: C, 84.65; H, 9.87; N, 5.48%.

Preparation of Hydrochlorides from 2-(7-Isopropyl-1-methyl-4-azulenyl)ethanamine Derivatives.

A) From 21: A solution of 209 mg (0.818 mmol) of **21** in 100 cm³ of benzene was extracted with 1 mol dm⁻³ hydrochloric acid (50 cm³ × 3); the hydrochloric acid layer was then extracted with chloroform (50 cm³ × 3). The chloroform solution was dried (sodium sulfate) and concentrated (reduced pressure) to leave a blue crystalline mass, which was washed with hexane (20 cm³ × 5) and chromatographed on a silica-gel column (elution with 9 : 1 chloroform-methanol) to give 120 mg (50%) of **27** as blue crystals, mp 200-204 °C: UV (C_2H_5OH) 218 (sh, log ϵ 4.05), 245 (4.29), 280 (sh, 4.46), 285 (4.52), 289 (4.51), 303 (3.95), 350 (3.58), 367 (3.41), 565 (sh, 2.51), 588 (sh, 2.57), 605 (2.61), 633 (sh, 2.56), 658 (2.53), and 730 nm (2.09); IR ($CHCl_3$) 2500, 2400, 2350, 2328 (NH^+), and 1097 cm⁻¹ (C-N); ¹H NMR ($CDCl_3$) δ =1.28 (6H, d, J =7 Hz, $CH(CH_3)_2$), 2.32 (6H, s, $N(CH_3)_2$), 2.61 (3H, s, 1- CH_3), 2.74 (2H, m, 1'-H₂), 3.02 (1H, m, J =7 Hz, $CH(CH_3)_2$), 3.31 (2H, m, 2'-H₂), 7.00 (1H, d, J =11 Hz, 5-H), 7.32 (1H, d, J =4 Hz, 3-H), 7.38 (1H, q, J =11 and 2 Hz, 6-H), 7.64 (1H, d, J =4 Hz, 2-H), and 8.18 (1H, d, J =2 Hz, 8-H).

B) From 24: By treating 210 mg (0.741 mmol) of **24** as described above in A, 153 mg (65%) of **28** was obtained as blue crystals, mp 200-204 °C: UV (C_2H_5OH) 215 (log ϵ 4.64), 246 (5.00), 285 (5.28), 290 (5.27), 303 (sh, 4.65), 330 (sh, 2.52), 351 (3.19), 368 (2.90), 440 (sh, 2.43), 565 (sh, 2.35), 614 (2.45), 666 (2.36), and 740 nm (1.89); IR ($CHCl_3$) 2615, 2540, 2510, 2440, 2350, 2340 (NH^+), and 1144 cm⁻¹ (C-N); ¹H NMR ($CDCl_3$) δ =1.34 (6H, s, 1'-C(CH_3)), 1.40 (6H, d, J =7 Hz, $CH(CH_3)_2$), 2.72 (3H, s, 1- CH_3), 2.82 (6H, s, $N(CH_3)_2^+$), 3.16 (1H, m, J =7.0 Hz, $CH(CH_3)_2$), 3.68 (2H, s, 2'-H₂), 7.12 (1H, d, J =11 Hz, 5-H), 7.54 (1H, q, J =11 and 4 Hz, 6-H), 7.60 (1H, d, J =4 Hz, 3-H), 7.84 (1H, d, J =4 Hz, 2-H), and 8.38 (1H, d, J =2 Hz, 8-H).

C) From 25: By treating 321 mg (1.04 mmol) of **25** as described in A, except that 6 mol dm⁻³ hydrochloric acid and a mixed solvent of chloroform-methanol (7 : 3) were used for extraction and for chromatographic separation respectively, 183 mg (51%) of **29** was obtained as blue crystals, mp 82-84

°C: UV (C_2H_5OH) 216 (log ϵ 3.21), 245 (3.69), 284 (4.03), 290 (4.02), 304 (sh, 2.54), 338 (sh, 3.38), 352 (3.46), 368 (3.38), 608 (2.59), 660 (2.49), and 730 nm (2.02); IR ($CHCl_3$) 2690, 2645, 2560, 2500-2400, 2360, 2340 (NH^+), and 1194 cm⁻¹ (C-N); ¹H NMR ($CDCl_3$) δ =1.32 (6H, s, C(CH_3)₂), 1.43 (6H, d, J =7 Hz, $CH(CH_3)_2$), 2.08 (4H, m, β - and γ -H₂), 2.24 (1H, s, NH^+), 2.76 (3H, s, 1- CH_3), 3.18 (1H, m, J =7 Hz, $CH(CH_3)_2$), 3.32 (4H, t, J =6 Hz, α - and δ -H₂), 3.68 (2H, s, 2'-H₂), 7.19 (1H, d, J =11 Hz, 5-H), 7.58 (1H, q, J =11 and 2 Hz, 6-H), 7.64 (1H, d, J =4 Hz, 3-H), 7.84 (1H, d, J =4 Hz, 2-H), and 8.36 (1H, d, J =2 Hz, 8-H).

D) From 26: By treating 415 mg (1.12 mmol) of **26** as described in C, 128 mg (28%) of **30** was obtained as blue crystals, mp 80-85 °C: UV (C_2H_5OH) 246 (log ϵ 4.03), 286 (4.34), 291 (4.31), 303 (3.52), 329 (sh, 3.46), 335 (sh, 3.54), 351 (3.66), 368 (3.54), 416 (2.90), 436 (2.89), 560 (sh, 2.50), 602 (sh, 2.59), 611 (2.60), 660 (2.52), and 730 nm (2.08); IR ($CHCl_3$) 2670, 2560, 2490-2430, 2400, 2360, 2340 (NH^+), and 1192 cm⁻¹ (C-N); ¹H NMR ($CDCl_3$) δ =1.23 (3H, s, 1'-C(CH_3)), 1.32 (6H, d, J =7 Hz, $CH(CH_3)_2$), 1.88 (4H, m, β - and γ -H₂), 2.64 (3H, s, 1- CH_3), 3.02 (4H, m, α - and δ -H₂), 3.70 (1H, d, J =12.5 Hz, 2'-H), 4.15 (1H, d, J =12.5 Hz, 2'-H), 6.10 (1H, d, J =11 Hz, 5-H), 6.96 (1H, q, J =11 and 2 Hz, 6-H), 7.30 (3H, m, 3'', 4'', and 5''-H), 7.34 (1H, d, J =4 Hz, 3-H), 7.64 (2H, m, 2''- and 6''-H), 7.62 (1H, d, J =4 Hz, 2-H), and 8.16 (1H, d, J =2 Hz, 8-H).

We wish to thank Professor Hitoshi Takeshita of Kyushu University, for his kind measurements of ¹³C NMR, and Konan-kako Co. Ltd., for generously giving us the sample of 4,6,8-trimethylazulene.

References

- 1) See, for example, W. E. Coyne, "Medicinal Chemistry," 3rd ed, ed by A. Burger, Wiley Interscience, New York (1970), Part II, pp. 953-968; W. T. Comer and A. W. Gomoll, *ibid.*, pp. 1019-1056; D. J. Triggle, *ibid.*, pp. 1235-1282.
- 2) S. Kurokawa and A. G. Anderson, Jr., *Bull. Chem. Soc. Jpn.*, **56**, 2059 (1983).
- 3) S. Kurokawa, *Chem. Lett.*, **1981**, 1569.
- 4) S. Kurokawa, *Bull. Chem. Soc. Jpn.*, **52**, 1748 (1979).
- 5) K. Hafner, H. Pelster, and H. Patzelt, *Justus Liebigs Ann. Chem.*, **650**, 80 (1961).
- 6) J. Schreiber, H. Maag, N. Hashimoto, and A. Eschenmoser, *Angew. Chem., Int. Ed. Engl.*, **10**, 330 (1971).
- 7) R. N. McDonald, J. M. Richmond, J. R. Curtis, H. E. Petty, and R. J. Mobley, *J. Am. Chem. Soc.*, **99**, 5739 (1977); Identifications by spectral data were performed according to N. L. Wolfe, Ph. D. Thesis, Kansas State University, Kansas, U. S. A., (1974), which was kindly sent us by Professor R. N. McDonald.
- 8) R. N. McDonald, H. E. Petty, N. L. Wolfe, and J. V. Paukstelis, *J. Org. Chem.*, **39**, 1877 (1974).
- 9) E. Heilbronner, "Non-Benzenoid Aromatic Compounds," ed by D. Ginsburg, Interscience, New York (1959), pp. 171-276.
- 10) K. Hafner and W. Senf, *Justus Liebigs Ann. Chem.*, **656**, 34 (1962).
- 11) N. J. Leonard and J. V. Paukstelis, *J. Org. Chem.*, **28**, 3021 (1963).
- 12) R. Tiollais, *Bull. Soc. Chim. Fr.*, **14**, 708 (1947); K. N. Campbell, A. H. Sommers, and B. K. Campbell, *J. Am. Chem. Soc.*, **66**, 82 (1944).
- 13) The tests for enzyme activity were performed by Masao Kidokoro at the Central Research Laboratory of Sankyo Co. Ltd., using established procedures.

14) Unpublished data available in our laboratory: These data were obtained by Cynthia H. Wilson and John B. Plastino at the University of Washington using established procedures.

15) K. Hafner and H. Weldes, *Justus Liebigs Ann. Chem.*, **606**, 90 (1957); K. Hafner, C. Bernhard, and R. Müller, *ibid.*, **650**, 35 (1961).

16) S. Braun and J. Kinkeldei, *Tetrahedron*, **33**, 1827 (1977).

17) K. Sato, M. Yamaguchi, and I. Ogura, *Bunseki Kagaku*, **31**, 185 (1982).

18) This numbering was used for pyrrolidine ring and for benzene ring:

