

Cycloaddition of Diphenylcyclopropenone with Carboximide, Carboximidamide, and Carboximidothioate

Hiroshi YOSHIDA,* Shingo SOGAME, Shoichi BANDO, Shosuke NAKAJIMA,
Tsuyoshi OGATA, and Kiyoshi MATSUMOTO†


Department of Applied Chemistry, Faculty of Engineering, Shizuoka University, Hamamatsu 432

†College of Liberal Arts and Sciences, Kyoto University, Sakyo-ku, Kyoto 606

(Received March 23, 1983)

Synopsis. The reaction of diphenylcyclopropenone with $R^1N=C(R^2)X$ (R^1, R^2 =alkyl, aryl, X =MeO, EtO, MeS, or Me₂N) gave 2-pyrrolin-4-one (**3**) in good yield. The less reactive carboximidothioate (**2**, R^1 =4-MeC₆H₄, R^2 =Ph, X =MeS) yielded isomeric 3-pyrrolin-2-one together with **3**.

In a previous paper¹⁾ we have reported the reaction of *N*-imidoyl sulfoximide with diphenylcyclopropenone to yield the sulfoximide substituted with pyrrolinone ring together with pyrimidinone *via* [2+3] and [3+3]cycloaddition reactions:



Eicher and his coworkers have reported that ketimine reacts with diphenylcyclopropenone **1** to yield 2-pyrrolin-4-one derivative by [2+3]cycloaddition reaction.²⁾ However, no work has been undertaken on the *N,N,N'*-trisubstituted carboximidamide, carbiximide, and carboximidothioate (**2**). In this paper we report the reactivities of these compounds with diphenylcyclopropenone.

An equimolar mixture of **1** and **2** in benzene (reaction at 80 °C) or xylene (over 100 °C) was heated at suitable temperature. The reaction was followed by TLC. The products isolated by column chromatography were 2-pyrrolin-4-one (**3**), but in the case of **2j**, **3j**, and 3-pyrrolin-2-one (**4j**) were obtained. The yields of **3** and **4**

and their physical properties are summarized in Table 1.

The structures of the products were elucidated on the basis of their elemental analyses, ¹H-NMR and mass spectroscopic studies as well as chemical transformations. Mass spectra of **3c**, **3g**, and **3j** showed peaks corresponding to the groups PhC=C(Ph)-NR¹ together with the parent peaks. Although **4j** showed a peak corresponding to PhC=C(Ph)-CPh, no such peak was observed for **3j**.


Some products were subjected to drastic hydrolysis. On treating with hot mixture of aqueous acetic and sulfuric acids, **3b**, **3e**, **3g**, **3i**, and **3j** gave the known triphenylfuranone (**5**)^{2c)} in moderate yields, while the isomer **4j** yielded the acid **6**.³⁾

It has been reported that *N*-methyl and *N*-phenylketimines react completely with **1** in boiling 1,2-dimethoxyethane (at about 83 °C) within 20 min and 2–5 h respectively.^{2c)} Table 1 indicates that the reactivity of **2** decreased in orders on R¹: alkyl>aryl, X: NMe₂>OMe, and OEt>SMe. Although the reactivities of **2** were lower than those of ketimines, the electron-releasing effects of substituents on R¹ and X would account for the reactivities of **2**. Nucleophilic attack of the imino nitrogen on the cyclopropenone **1** gave **3** or **4** depending on the attacking site. The major products were 2-pyrrolin-4-ones (**3**) whereas the less reactive **2j** showed lower selectivity under more drastic conditions giving a mixture of **3j** and **4j**. In conclusion, the reaction of **1** with

TABLE 1. YIELDS AND PHYSICAL PROPERTIES OF **3** AND **4**

Reactant	Reaction conditions		Product	Yield %	Mp θ _m /°C	(Recryst solvent) ^{a)}	¹ H-NMR (δ in CDCl ₃) ^{b)}	M ⁺	Found (Calcd) (%)		
	Temp/°C	Time/h							C	H	N
2a	110	2	3a	90	105–106	(E-P)	1.65 (3H, s, MeC-N), 2.00 (3H, s, MeS) 3.05 (3H, s, MeN)	309	73.79 (73.75)	6.27 (6.19)	4.44 (4.53)
2b	80	3	3b	73	157–158	(E-P)	1.36 (3H, t, <i>J</i> =7 Hz, MeCH ₂), 2.76 (3H, s, MeN), 3.65 (2H, q, CH ₂)	369	81.51 (81.27)	6.18 (6.27)	3.83 (3.79)
2c	110	4	3c	88	96	(E-P)	2.09 (3H, s, MeS), 2.97 (3H, s, MeN)	371	77.24 (77.57)	5.81 (5.70)	3.91 (3.77)
2d	110	2	3d	85	142	(C-P)	0.55 (3H, t, <i>J</i> =7 Hz, MeCH ₂), 2.56 (6H, s, Me ₂ N), 3.30 (2H, dq, <i>J</i> =5 and 7 Hz, CH ₂)	382	81.69 (81.64)	6.77 (6.85)	7.41 (7.32)
2e	110	16	3e	71	177–179	(C-P)	2.64 (6H, s, Me ₂ N)	430	83.47 (83.69)	6.11 (6.09)	6.49 (6.51)
2f	130	8	3f	61	165–168	(E-P)	3.65 (3H, s, MeO)	417	83.42 (83.43)	5.39 (5.55)	3.04 (3.35)
2g	130	8	3g	87	190–191	(E-P)	1.28 (3H, t, <i>J</i> =7 Hz, Me), 3.86 (2H, q, CH ₂)	431	83.61 (83.50)	5.92 (5.84)	3.16 (3.25)
2h	110	10	3h	82	208–210	(C-P)	2.12 (3H, s, MeC ₆ H ₄), 2.52 (6H, s, Me ₂ N)	444	83.83 (83.75)	6.35 (6.35)	6.43 (6.30)
2i	130	8	3i	83	178–180	(E-P)	1.27 (3H, t, <i>J</i> =7 Hz, MeCH ₂), 2.16 (3H, s, MeC ₆ H ₄), 3.86 (2H, q, CH ₂)	445	83.21 (83.57)	6.28 (6.11)	3.04 (3.15)
2j	140	18	3j	48	97–99	(P)	2.27 (3H, s, MeC ₆ H ₄), 2.34 (3H, s, MeS)	400 ^{c)}	80.77 (80.50)	5.64 (5.63)	2.99 (3.13)
			4j	33	104–106	(P)	1.98 (3H, s, MeS), 2.28 (3H, s, MeC ₆ H ₄)	400 ^{c)}	80.39 (80.50)	5.45 (5.63)	3.61 (3.13)
2k	110	10	3k	73	163	(C-P)	2.58 (6H, s, Me ₂ N), 3.68 (3H, s, MeO)	460	80.62 (80.81)	6.24 (6.12)	6.11 (6.08)
2l	110	10	3l	80	98–100	(C-P)	2.14 (3H, s, MeC ₆ H ₄), 2.46 (6H, s, Me ₂ N)	444	83.72 (83.75)	6.29 (6.35)	6.54 (6.30)

a) C: CHCl₃, E: EtOH, P: petroleum ether. b) Aryl ring protons were observed at δ 6.4–8.1 as multiplets. c) M⁺ –SMe.



2 has proven to be a useful method for the preparation of 2-pyrrolin-4-ones (**3**).

Experimental

General. Melting points are uncorrected. ¹H-NMR spectra were recorded on a Hitachi Perkin Elmer R-24 (60 MHz) spectrometer using TMS as an internal standard, and mass spectra on a Hitachi RMU-7M mass spectrometer.

Preparation of Carboximide, Carboximidate, and Carboximidothioate. The compounds **2** were prepared by the usual methods.⁴⁾ Carboximidamides **2d**, **e**, **h**, **k**, and **2l** were obtained from the corresponding imidoyl chlorides and excess dimethylamine in benzene. Carboximidates **2b**, **f**, **g**, and **2i** were prepared from the imidoylchlorides and sodium alkoxide in alcohol. Carboximidothioates **2a**, **c**, and **2j** were obtained by the reaction of the corresponding thioamides with methyl iodide followed by treatment with aqueous sodium hydroxide. The purity of the products was confirmed by ¹H-NMR. New

compounds were as follows: **2h** (85%): oil; ¹H-NMR (CDCl₃) δ=2.04 (3H, s, Me), 2.83 (6H, s, Me₂N), and 6.2–7.6 (9H, m, Arom); *m/e* 238 (M⁺). **2l** (93%): oil; ¹H-NMR (CDCl₃) δ=2.17 (3H, s, Me), 2.89 (6H, s, Me₂N), and 6.4–7.4 (9H, m, Arom); *m/e* 238 (M⁺).

The Reaction of 1 with 2. A solution of **1** (1.5 mmol) and **2** (2 mmol) was refluxed in a minimum amount of benzene or xylene (3–5 cm³). The reaction was checked by TLC (aluminum oxide, petroleum ether : ethyl acetate, 1 : 1, v/v) until **1** reacted completely. After cooling, the mixture was chromatographed over silica gel (petroleum ether : ethyl acetate, 4 : 1, v/v). The products were crystallized from appropriate solvents.

Hydrolyses of 3 and 4. In a boiling mixture of acetic acid and sulfuric acid in water (50 : 1 : 25, v/v, 5 cm³) pyrrolinone (50 mg) was hydrolyzed for 2 d. The resulting mixture was poured into water and extracted with chloroform. The extract was dried over sodium sulfate and evaporated under reduced pressure. **3b**, **e**, **g**, **i**, and **3j** gave triphenyl-3-(2H)-furanone **5**: mp 191–192 °C (lit, mp 192–193 °C).^{2c)} In contrast **4j** yielded the acid **6**: mp 172–174 °C (lit, mp 172 °C).³⁾

References

- 1) H. Yoshida, S. Sogame, Y. Takishita, and T. Ogata *Bull. Chem. Soc. Jpn.*, **56**, 2438 (1983).
- 2) a) T. Eicher, F. Abdesaken, G. Franke, and J. L. Weber, *Tetrahedron Lett.*, **1975**, 3915; b) T. Eicher, G. Franke, and F. Abdesaken, *ibid.*, **1977**, 4067; c) T. Eicher, J. L. Weber, and G. Chatila, *Justus Liebigs Ann. Chem.*, **1978**, 1203; d) T. Eicher and G. Franke, *ibid.*, **1981**, 1337.
- 3) J. Hoch and R. Teixier, *C. R. Acad. Sci.*, **261**, 4132 (1965).
- 4) S. R. Sandler and W. Karo, "Organic Functional Group Preparations," Academic Press, New York (1972), Vol. III.