

Synthesis of 4,4-Disubstituted 2,3-Dicyano-5-imino-2-pyrrolines. Reaction of Alkylidenemalononitriles with Trimethylsilyl Cyanide


Naoto CHATANI*,† and Terukiyo HANAFUSA

The Institute of Scientific and Industrial Research, Osaka University, Ibaraki, Osaka 567


(Received March 17, 1990)

Synopsis. The reaction of alkylidenemalononitriles with trimethylsilyl cyanide in the presence of KCN/18-crown-6 gave 4,4-disubstituted 2,3-dicyano-5-imino-2-pyrrolines in high yields.


In a series of transition metal-catalyzed reaction of trimethylsilyl cyanide (**1**),¹⁾ we have already reported that **1** reacted with aryl iodides in the presence of Pd(PPh₃)₄ to give aryl cyanides in good yields (Eq. 1).²⁾



During the examination of scope and limitations of the cyanation reaction, we found that 1,1-dibromo-2,2-diphenylethylene (**2**) reacted with **1** in refluxing toluene in the presence of Ni(PPh₃)₂(CO)₂ gave (diphenylmethylene)malononitrile (**3a**) in 92% yield (Eq. 2). However, 2,3-dicyano-4,4-diphenyl-5-imino-2-






pyrroline (**4a**) was obtained along with **3a** when the reaction was carried out without solvent. It was expected that **4a** would be produced by a further reaction of the primary product **3a** with **1**. Actually, **4a** was obtained in a high yield from the Ni-catalyzed reaction of **3a** with **1**. As a result of optimization of the reaction conditions, we have found that the nickel catalyst was not necessary to proceed the construction of pyrroline ring from **3**. We now wish to report that the reaction of alkylidenemalononitriles **3** with **1** gave 4,4-disubstituted 2,3-dicyano-5-imino-2-pyrrolines (**4**) in high yields (Eq. 3).³⁾




The results are summarized in Table 1. Cyclohexylidenemalononitrile (**3b**) was used as substrate for optimization of the reaction conditions. Higher temperatures gave higher yields (Entries 1–3). Formation of **4b** proceeded even the reaction temperature was 25 °C and even no catalyst was present (Entry 3). Although the presence of KCN did not affect the yield of the **4b** (Entry 4), we have found that a small

Table 1. Reaction of Alkylidenemalononitriles with Me₃SiCN (**1**)^{a)}

Entry	Substrate	Temp/°C	Catalyst	Product	Yield/% ^{b)}
1		Reflux	—		92
2	3b	80	—	4b	82
3	3b	25	—	4b	52
4	3b	25	KCN	4b	60
5	3b	25	KCN/18-crown-6	4b	98
6		25	KCN/18-crown-6		50
7		25	KCN/18-crown-6		26
8	3a	50	KCN/18-crown-6	4a	86

a) Reaction conditions: alkylidenemalononitrile (2.5 mmol), Me₃SiCN (15 mmol, 2.0 ml), KCN (0.1 mmol), 18-crown-6 (0.1 mmol) for 20 h. b) Isolated yields based on the substrate.

† Present address: Department of Applied Chemistry, Faculty of Engineering, Osaka University, Suita, Osaka 565.




Scheme 1.

amount of KCN/18-crown-6 are exceptionally mild and efficient catalyst for the formation of **4b** (Entry 5).⁴ Phenyl-substituted alkylidenemalononitriles **3a** and **3c** gave the corresponding 2,3-dicyano-5-imino-2-pyrrolines **4a** and **4c**, respectively (Entries 6–8). The reaction of benzylidenemalononitrile (**3d**), which has a hydrogen at the vinylic position, with **1** gave a complex mixture.

A plausible reaction path is shown in Scheme 1. Michael addition of **1** to alkylidenemalononitrile **3** catalyzed by CN^- ⁵ gave a ketenimine **5** to which **1** added to give intermediate **6**. Ring closure of **6** by an intramolecular nucleophilic attack of the silylamino group at the cyano group led silyl-protected 2,3-dicyano-5-imino-2-pyrroline derivative **7**. Desilylation of **7** by EtOH gave a pyrroline **4**.

The reaction of cyclohexylidenecyanoacetate (**8**) did not give an expected product but hydrocyanation product **9** was obtained as a sole product (Eq. 4).



In summary, we found that alkylidenemalononitriles are the precursor for the preparation of 4,4-disubstituted 2,3-dicyano-5-imino-2-pyrrolines,⁶ which would be useful intermediates for organic synthesis because of their multi-functionalities.⁷

Experimental

General. Melting points (mp) were determined on a Yanagimoto micro melting point apparatus and uncorrected. Infrared spectra were obtained on a Hitachi 260-10 spectrometer. Peaks are reported in units of cm^{-1} . ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker WM 360 spectrometer and are reported in ppm from tetramethylsilane or chloroform as an internal standard on the δ scale. Data are reported as follows: chemical shift, multiplicity (s=singlet, d=doublet, t=triplet, m=multiplet, br=broad), coupling constant, integration, and interpretation. Mass spectra were obtained on a JMS-DX 300 with ionization voltages of 70 eV. Elemental analyses were performed on a Perkin-Elmer 240C by the ISIR Material Analysis Center in Osaka University.

Materials. 1,1-Dibromo-2,2-diphenylethylene (**2**) was prepared by the literature method.⁸ Alkylidenemalononitriles (**3**) were prepared from the Knoevenagel reaction⁹ of corresponding ketones with malononitrile according to the method reported previously by our group.¹⁰ Trimethylsilyl

cyanide was obtained from Tokyo Kasei Co., and distilled from CaH_2 . 18-Crown-6 was purchased from Aldrich Co., and recrystallized from CH_3CN .

General Procedure for the Reaction of Alkylidenemalononitriles with Trimethylsilyl Cyanide. To a solution of alkylidenemalononitrile (2.5 mmol) and Me_3SiCN (15 mmol, 2.0 ml) was added KCN/18-crown-6 (0.1 mmol/0.1 mmol). The solution was stirred at 25°C for 20 h and excess Me_3SiCN was distilled in vacuo. To the residue was added EtOH (20 ml) and the mixture was stirred at 25°C for 1 h. Recrystallization of the residue from acetone/hexane gave 4,4-disubstituted 2,3-dicyano-5-imino-2-pyrrolines. Spectral samples were obtained by recrystallization from EtOH.

2,3-Dicyano-5-imino-4,4-pentamethylene-2-pyrroline (4a). White solid, mp 220°C (decomp); IR (Nujol) 3360, 3330, 2230, 2205, 1675, 1650, 1550 cm^{-1} ; ^1H NMR (acetone- d_6) δ =1.74–1.96 (c, 10 H, CH_2), 7.63 (br, 1 H, NH), 8.04 (br, 1 H, NH); ^{13}C NMR (acetone- d_6) δ =22.58, 25.24, 34.12, 58.54, 112.2, 114.3, 116.7, 142.0, 185.6; MS m/z 200 (M^+). Found: C, 66.01; H, 5.85; N, 27.77%. Calcd for $\text{C}_{11}\text{H}_{12}\text{N}_4$: C, 65.98; H, 6.04; N, 27.98%.

2,3-Dicyano-5-imino-4-methyl-4-phenyl-2-pyrroline (4b). White solid; mp 218°C (decomp); IR (Nujol) 2420, 2230, 2200, 1655, 1570; ^1H NMR (acetone- d_6) δ =1.84 (s, 3 H, CH_3), 7.43 (d, J =7.0 Hz, 2 H, Ph), 7.44–7.46 (m, 3, Ph), 7.65 (br, 1 H, NH), 8.14 (br, 1 H, NH); ^{13}C NMR (acetone- d_6) δ =20.52, 61.58, 113.7, 114.1, 116.2, 127.0, 129.5, 130.2, 136.7, 140.2, 185.5. MS m/z 222 (M^+); Found: C, 69.96; H, 4.16; N, 24.55%. Calcd for $\text{C}_{13}\text{H}_{10}\text{N}_4$: C, 70.26; H, 4.54; N, 25.21%.

2,3-Dicyano-4,4-diphenyl-5-imino-2-pyrroline (4c). White solid; mp 280°C (decomp); IR (Nujol) 3400, 2210, 1670, 1560; ^1H NMR (acetone- d_6) δ =7.29–7.51 (m, 10 H, Ph), 8.69 (br, 2 H, NH); MS m/z 284 (M^+). Found: C, 76.25; H, 4.13; N, 19.29%. Calcd for $\text{C}_{18}\text{H}_{12}\text{N}_4$: C, 76.04; H, 4.25; N, 19.71%.

Ethyl Cyano(1-cyanocyclohexyl)acetate (11). Colorless oil; IR (neat) 2250, 2240, 1745; ^1H NMR (CDCl_3) δ =4.33 (q, J =7.2 Hz, 2 H, CH_2O), 3.65 (s, 1 H, CH), 2.24 (m, 1 H, CH_2), 2.08 (m, 1 H, CH), 1.53–1.88 (m, 8 H, CH_2), 1.37 (t, J =7.2 Hz, 3 H, CH_3); MS, m/z 220 (M^+).

References

- For previous papers of this series, see: N. Chatani, N. Horiuchi, and T. Hanafusa, *J. Org. Chem.*, **55**, 3393 (1990). N. Chatani, T. Takeyasu, and T. Hanafusa, *Tetrahedron Lett.*, **29**, 3979 (1988). N. Chatani, T. Takeyasu, N. Horiuchi, and T. Hanafusa, *J. Org. Chem.*, **53**, 3539 (1988).
- N. Chatani and T. Hanafusa, *J. Org. Chem.*, **51**, 4714 (1986).
- For a paper on the construction of 2-imino-4-pyrroline ring, see: P. Merot, C. Gadreau, and A. Foucaud, *Tetrahedron*, **37**, 2595 (1981).
- KCN/18-crown-6 was used as a catalyst for the reaction of aldehydes with **1**. D. A. Evans and L. K. Truesdale, *Tetrahedron Lett.*, **1973**, 4929.
- It is known that alkylidenemalononitriles react with KCN to give tricyanoethanes. A recent paper on the reaction, see: J. A. Ciller, N. Martin, M. Quinteiro, C. Seoane, and J. L. Soto, *J. Chem. Res. Synop.*, **1987**, 166.
- For a paper on the preparation of a variety of heterocyclic compounds by the use of alkylidenemalononitriles, see: Ref. 5 and references cited therein.
- "Advances in Heterocyclic Chemistry," ed by A. R. Katritzky, Academic Press, Oxford (1986).
- R. E. Buckless, E. A. Hausman, and N. G. Wheeler, *J. Am. Chem. Soc.*, **72**, 2494 (1950).
- G. Jones, *Org. React.*, **15**, 204 (1967).
- J. Yamawaki, T. Kawate, T. Ando, and T. Hanafusa, *Bull. Chem. Soc. Jpn.*, **56**, 1885 (1983).