

Conjugate Reduction of 2-Butene-1,4-diones with LiAlH₄-SbCl₃

Shinsei SAYAMA* and Yutaka INAMURA

Department of Chemistry, Fukushima Medical College, Hikarigaoka, Fukushima 960-12

(Received June 15, 1990)

Synopsis. The reagent LiAlH₄-SbCl₃ was found to be more effective for a conjugate reduction of 2-butene-1,4-diones in comparison with the reagent LiAlH₄-other metal halides.

The effect of the addition of AlCl₃ to LiAlH₄ was to produce more mild and specific reagents for reducing epoxides and halo ketones.¹⁾ Therefore, a chemoselective reduction of polyfunctional compounds with LiAlH₄ in the presence of various other metal halides has been the subject of much study.²⁾ It was reported that the reduction of α,β -unsaturated ketones with LiAlH₄-Cu₂I₂ or LiAlH₄-metal halides (e.g., Cu₂Br₂, Cu₂Cl₂, TiCl₃, FeCl₃) gave the corresponding ketones arising from a conjugate reduction.^{3,4)} Since γ -keto- α,β -unsaturated ketones, such as 2-butene-1,4-diones, afforded 1,2-reduction products predominantly with LiAlH₄ or NaBH₄, there has been much interest in mild, convenient methods for a conjugate reduction of γ -keto- α,β -unsaturated ketones. The reduction of γ -keto- α,β -unsaturated ketones with LiAlH₄-metal halides, however, has not been carried out.^{1–7)} Moreover, SbCl₃ has not been used as a metal halide in a conjugate reduction of α,β -unsaturated ketones and γ -keto- α,β -unsaturated ketones. Since antimony is one of the representative elements which possess both metallic and nonmetallic properties and Sb³⁺ is a borderline acid in the classification of hard and soft acids,^{2a,6,7)} it can be expected that the reduction of carbonyl compounds with LiAlH₄-SbCl₃ has a new reactivity in comparison with LiAlH₄-transition metal halides. We would like to report on the results of studies concerning the selective conjugate reduction of γ -keto- α,β -unsaturated ketones, such as 2-butene-1,4-diones, with LiAlH₄-SbCl₃.

The reduction of *trans*-1-phenyl-2-pentene-1,4-dione (**1**), chosen as a representative 2-butene-1,4-dione for

this study, with LiAlH₄ (LAH)-SbCl₃ in various stoichiometric ratios was carried out. The results are summarized in Table 1. At the ratio of enedione **1**, LAH, and SbCl₃ (1:1:1, 1:1:3, or 1:3:1), a mixture of 4-hydroxy-1-phenyl-1-pentanone, 1-phenyl-2-pentene-1,4-diol and 1-phenylpentane-1,4-diol and **1** was mainly obtained, accompanied by a small amount of 1-phenylpentane-1,4-dione (**1a**). At a stoichiometric ratio of 1:3:3 or 1:3:4, enedione **1** was reduced to give a conjugate reduction product **1a** in nearly quantitative yield. The optimum conditions for the conjugate reduction of **1** with LAH-SbCl₃ found in the present experiments are as follows: i) SbCl₃ is essential for effecting a regioselective conjugate reduction. ii) There is need to use either an equal or excess molar equivalent of SbCl₃ over LAH in order to suppress the reaction of 1,2-reduction with LAH. iii) There is need to use more than two molar equivalents of LAH and SbCl₃ over **1** for obtaining conjugate reduction products in high yield.

The results of a LAH-SbCl₃ reduction of other 2-butene-1,4-diones are shown in Table 2. The reductions of *trans*-3-decene-2,5-dione (**2**), ethyl 3-benzoylacrylate (**3**), *trans*- and *cis*-1,4-diphenyl-2-butene-1,4-diones (**5** and **6**), even diethyl azodicarboxylate (**4**) and 3-benzoyl-1-phenyl-2-pentene-1,4-dione (**7**) with LAH-SbCl₃ took place to give the corresponding conjugate reduction products in good yields.











The results of a LAH-SbCl₃ reduction of α,β -unsaturated carbonyl compounds are shown in Table 3. The reduction of α,β -unsaturated ketones (**9** and **10**) was not selective, as shown. Only the formyl group of **8** was reduced to a hydroxyl group in 92% yield. The chemoselective reduction of aldehydes in the presence of ketones with this reagent was also examined in the following competition experiments. Equimolar amounts of 3,7-dimethyl-6-octenal and ethyl levulinate were

Table 1. Reduction of *trans*-1-Phenyl-2-pentene-1,4-dione **1** with LiAlH₄-SbCl₃ in THF at 0°C for 4 h^{a)}

Entry	Molar ratio			Products, yield/% ^{b)}			
	Enedione 1	LiAlH ₄	SbCl ₃	1-Phenylpentane-1,4-dione 1a	Hydroxy compound		Recovered enedione 1
					Hydroxy ketone 1b	Diol 1c	
1	1	2	0	—	56 ^{c)}	38 ^{d)}	—
2	1	1	1	32	30 ^{e)}	—	33
3	1	1	3	23	—	—	76
4	1	2	2	60	10 ^{e)}	—	21
5	1	3	3	84	6 ^{e)}	—	—
6	1	3	1	14	42 ^{e)}	43 ^{f)}	—
7	1	3	9	87	—	—	—
8	1	3	4	93	—	—	—
9	1	0	3	—	—	—	99

a) LAH; 1.0–3.0 mmol; Solvent; 10–15 ml. b) Yield is based on enedione **1** used. c) 4-Hydroxy-1-phenyl-1-pentanone. d) 1-Phenyl-2-pentene-1,4-diol. e) Mixture of **1b** and **1c**. f) Mixture of enediol and saturated diol.

Table 2. Reduction of 2-Butene-1,4-dione with $\text{LiAlH}_4\text{-SbCl}_3$ in THF at 0°C for 4 h^{a)}

Substrate	(S)	Molar ratio			Products	Yield/% ^{b)}
		S	LAH	SbCl_3		
	1	1	3	3		1a ^{a)} 84
	2	1	3	3		2a ^{a,9)} 94
$\text{PhCOCH=CHCO}_2\text{Et}$	3	1	3	3	$\text{PhCOCH}_2\text{CH}_2\text{CO}_2\text{Et}$	3a ^{a)} 86
$\text{EtOOCN=NCOC}_2\text{Et}$	4	1	3	3	$\text{EtOOCNHNHCO}_2\text{Et}$	4a 81
	5	1	3	3		5a ^{12,13)} 78
	6	1	3	3		5a ^{12,13)} 81
	7	1	2.5	2.5		7a 64

a) LAH; 1.0–3.0 mmol; Solvent; 10–15 ml. b) Yield is based on 2-butene-1,4-dione used.

Table 3. Reduction of Carbonyl Compounds with $\text{LiAlH}_4\text{-SbCl}_3$ in THF at 0°C for 4 h^{a)}

Substrate	(S)	Molar ratio			Products	Yield ^{b)} /%
		S	LAH	SbCl_3		
$\text{C}_6\text{H}_5\text{CH=CHCHO}$	8	1	2	2	$\text{C}_6\text{H}_5\text{CH=CHCH}_2\text{OH}$ $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	8a 92 8b 7
$\text{C}_6\text{H}_5\text{CH=CHCOCH}_3$	9	1	3	3	$\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{COCH}_3$ $\text{C}_6\text{H}_5\text{CH=CHCH(OH)CH}_3$ Recovered	9a 12 9b 44 9 42
$\text{C}_6\text{H}_5\text{CH=CHCOC}_6\text{H}_5$	10	1	3	3	$\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{COC}_6\text{H}_5$ Recovered	10a 16 10 50

a) LAH; 1.0–3.0 mmol; Solvent; 10–15 ml. b) Yield is based on the carbonyl compounds used.

allowed to compete for the reduction with three molar equivalents of LAH-SbCl_3 . 3,7-Dimethyl-6-octenal was selectively reduced to 3,7-dimethyl-6-octene-1-ol in 70% yield and ethyl levulinate was recovered unchanged. In addition, carbonyl groups of 1-phenylbutane-1,3-dione and 1-phenylpentane-1,4-dione and ester, nitrile, and acetal groups in ethyl levulinate and 4-phenyl-4,4-ethylenedioxybutanenitrile were not affected under the above-mentioned reduction conditions.

On the other hand, the reduction of enedione **1** with LAH —other metal halides such as Cu_2I_2 , CuCl_2 , AlCl_3 , FeCl_3 , and TiCl_4 gave no 1,4-reduction products, and complex mixture of hydroxy derivatives were obtained under various ratios of LAH and metal halides.

Thus, the reagent LAH-SbCl_3 provides a new convenient method for a mild, selective conjugate

reduction of 2-butene-1,4-diones. Further application of reduction with LAH-SbCl_3 to aldehyde and other functional groups is now in progress.

Experimental

IR spectra were recorded on a JASCO A-100 spectrometer. ^1H NMR spectra were taken on a Hitachi R-24B spectrometer with TMS as an internal standard. The products were identified by spectroscopic data.

General Procedure. To a suspension of LAH (113 mg, 3 mmol) in THF (8 ml) at 0°C was added SbCl_3 (684 mg, 3 mmol) dissolved in THF (2 ml). The resulting mixture was stirred for 5 min at 0°C , and then 2-butene-1,4-dione (1 mmol) in THF (2 ml) was added. After stirring for 4 h at 0 – 18°C , the reaction mixture was treated with 1 M aq Na_2CO_3 and extracted with ethyl acetate. The organic layer was washed by 0.5 M aq HCl and successively saturated aq

NaCl and dried by MgSO_4 . After removal of the solvent in vacuo, the residue was purified by column chromatography on silica gel (Wakogel C-300) with CCl_4 and CHCl_3 (3:1). Conjugate reduction products **1a**—**7a** were obtained in 64—94% yield.

2-Butene-1,4-diones were prepared according to a procedure from the literature.

trans-1-Phenyl-2-pentene-1,4-dione (**1**)¹¹ and *trans*-3-decene-2,5-dione (**2**) were prepared by pyridinium chlorochromate (PCC) oxidation of the corresponding 2,5-dialkylfuran derivatives.¹⁰

trans-1,4-Diphenyl-2-butene-1,4-dione (**5**)^{11,12} was prepared by Friedel-Crafts acylation of benzene with fumaryl chloride.¹⁴ *cis*-1,4-Diphenyl-2-butene-1,4-dione (**6**)^{11,12} and 3-benzoyl-1-phenyl-2-pentene-1,4-dione (**7**) were prepared by oxidation of corresponding furan derivatives with HNO_3 in acetic acid.¹⁵

3-Benzoyl-1-phenylpentane-1,4-dione (7a): Mp 83—84°C (from CCl_4); IR (KBr) 1720, 1690, and 1675 cm^{-1} ; ^1H NMR (CDCl_3) δ =2.22 (3H, s), 3.70 (2H, m), 5.30 (1H, t, J =6.0), 7.26—8.26 (10H, m).

***trans*-3-Decene-2,5-dione (2):** Mp 46—47°C (from hexane); IR (KBr) 1670 cm^{-1} ; ^1H NMR (CDCl_3) δ =0.85 (3H, t, J =7.0), 1.13—2.00 (6H, m), 2.33 (3H, s), 2.47—2.83 (2H, m), 6.70 (2H, s).

3-Benzoyl-1-phenyl-2-pentene-1,4-dione (7): Mp 112—113°C (from ethanol); IR (KBr) 1685 and 1660 cm^{-1} ; ^1H NMR (CDCl_3) δ =2.35 (3H, s), 7.20—8.10 (11H, m).

References

- 1) a) H. O. House, "Modern Synthetic Reactions," W. A. Benjamin, Inc., Menlo Park, California (1972), p. 45; b) E. C. Ashby and B. Cooke, *J. Am. Chem. Soc.*, **90**, 1625 (1968).
- 2) a) T. Imamoto, T. Takeyama, and T. Kusumoto, *Chem. Lett.*, **1985**, 1491; b) E. C. Ashby and J. J. Lin, *Tetrahedron Lett.*, **1977**, 4481.
- 3) E. C. Ashby, J. J. Lin, and R. Kover, *J. Org. Chem.*, **41**, 1939 (1976), and references cited therein.
- 4) T. Tsuda, T. Fujii, K. Kawasaki, and T. Saegusa, *J. Chem. Soc., Chem. Commun.*, **1980**, 1013.
- 5) a) M. Fieser and L. F. Fieser, "Reagents for Organic Synthesis," John Wiley & Sons, Inc., New York, Vol. 1—13; b) S. S. Pizey, "Synthetic Reagents," John Wiley & Sons, Inc., New York (1974), Vol. 1, p. 101.
- 6) T.-L. Ho, "Hard and Soft Acids and Bases Principle in Organic Chemistry," Academic Press, New York (1977), p. 93.
- 7) J. Bottin, O. Eisenstein, C. Minot, and N. T. Anh, *Tetrahedron Lett.*, **1972**, 3015.
- 8) H. Stetter, *Angew. Chem., Int. Ed. Engl.*, **15**, 639 (1976).
- 9) H. Stetter and H. Kuhlmann, *Tetrahedron Lett.*, **1974**, 4505.
- 10) G. Piancatelli, A. Scettri, and M. D'Auria, *Tetrahedron*, **36**, 661 (1980).
- 11) C.-S. Chien, T. Kawasaki, M. Sakamoto, Y. Tamura, and Y. Kita, *Chem. Pharm. Bull.*, **33**, 2743 (1985).
- 12) Y. Inamura and Y. Inamura, *Nippon Kagaku Kaishi*, **1985**, 134.
- 13) Y. Kobayashi, T. Taguchi, T. Morikawa, E. Tokuno, and S. Sekiguchi, *Chem. Pharm. Bull.*, **28**, 262 (1980).
- 14) R. E. Lutz, "Organic Syntheses," John Wiley & Sons, Inc., New York (1955), Coll. Vol. III, p. 248.
- 15) a) R. E. Lutz and R. J. Rowlett, Jr., *J. Am. Chem. Soc.*, **70**, 1359 (1948); b) R. E. Lutz and F. N. Wilder, *ibid.*, **56**, 978 (1934).