

One-Electron Reductive Cleavage of the C–Cl Bond of ArCF_2Cl : Convenient Route for the Synthesis of ArCF_2 Derivatives

Masato Yoshida,* Ayako Morishima, Daiki Suzuki, Masahiko Iyoda, Kozo Aoki,[†] and Shigeru Ikuta[†]

Department of Chemistry, Faculty of Science, Tokyo Metropolitan University, Hachioji, Tokyo 192-03

[†]Computer Center, Tokyo Metropolitan University, Hachioji, Tokyo 192-03

(Received January 23, 1996)

When ArCF_2Cl was reacted with PhSeNa in DMF in the presence of light, it was found that a $\text{S}_{\text{RN}}1$ type reaction occurred to give ArCF_2SePh . As the reaction did not proceed without light, the substitution reaction should be initiated by photo-induced electron transfer from PhSe^- to ArCF_2Cl . The cleavage of the C–Cl bond of PhCF_2Cl by one-electron reduction with samarium(II) iodide to give a PhCF_2 radical and chloride ion was also observed. The PhCF_2 radical, thus formed, was trapped by styrene and phenylacetylene before it was reduced to PhCF_2 anion. UHF/3-21G calculation demonstrated that one-electron reduction in PhCF_2Cl led the spontaneous dissociation of the C–Cl bond to the PhCF_2 radical and Cl^- , but the PhCF_3 radical anion existed as a stable form.

Recently, the molecules bearing a *gem*-difluoromethylene group ($-\text{CF}_2-$) have attracted much interest, because the replacement of $-\text{CH}_2-$ to $-\text{CF}_2-$ brings about dramatic changes in the physical, chemical, and biological properties of the parent molecules.¹⁾ However, convenient methods for the preparation of these molecules are limited,²⁾ and the development of such methodology is still an important and challenging target for organic chemists. Since there are many ArCH_2 branched natural and biological active compounds, a development of a convenient route for the synthesis of ArCF_2 derivatives is especially interesting. As a precursor for the synthesis of ArCF_2 derivatives, ArCF_3 is inconvenient due to the strong C–F bond, and little is known about the routes from ArCF_3 ; the C–F bond is too stable to replace the fluorine with other functional groups.³⁾ The carbon–chlorine bond is predicted to be more reactive than the C–F bond, and ArCF_2Cl is, accordingly, expected to be a useful precursor for the synthesis of ArCF_2 derivatives. We have been exploring the chlorodifluoromethylation of aromatic rings with bis(chlorodifluoroacetyl) peroxide and successive conversion of the chlorine into other functional groups.⁴⁾ In our previous paper, we reported that the chlorine atom of ArCF_2Cl was readily replaced by a hydrogen or allyl group under radical conditions.⁴⁾ We have now explored the reductive cleavage of the C–Cl bond of ArCF_2Cl by a one-electron reducing agent such as PhSe^- and samarium(II) iodide (SmI_2), and the results are described in this paper with theoretical considerations using *ab initio* molecular orbital calculation.


Results and Discussion

It is known that the C–Cl bond in chlorofluorocarbons is usually inert to normal nucleophilic attack due to the strong electron-withdrawing power of fluorine. The atomic

charge of Cl in PhCF_2Cl was calculated by UHF/3-21G to be -0.0137 ; this means that the C–Cl bond in PhCF_2Cl is little polarized and therefore nonreactive to nucleophilic attack. In fact, PhCF_2Cl was found to be resistant to the usual nucleophiles, such as alkyllithium and alkylmagnesium halide. However, the replacement of the chlorine of ArCF_2Cl with PhSe^- was observed in the reaction of PhCF_2Cl with PhSeNa . As the reaction did not proceed without light, the substitution reaction is probably initiated by a photo-induced electron transfer from PhSe^- to ArCF_2Cl as shown in Scheme 1. In this reaction, PhSe^- did not act as a nucleophile but as a one-electron reducing agent. PhS^- and PhSe^- have been reported to act as a one-electron reducing agent in the $\text{S}_{\text{RN}}1$ type reactions of halofluoro compounds.^{5,6)}




Thus, the C–Cl bond of ArCF_2Cl was found to be readily dissociated by one-electron reduction to give a ArCF_2 radical and chloride ion.⁷⁾ Uneyama et al. have reported perfluoroalkyl-chalcogenations (sulfenylation, selenenylation, and tellurenylation) of olefins and acetylenes by the reaction of PhX^- ($\text{X} = \text{S}, \text{Se}, \text{and Te}$) with perfluoroalkyl halide. The reaction is initiated by a one-electron transfer from PhX^- to perfluoroalkyl halide to give a perfluoroalkyl radical followed by a radical chain reaction via addition to olefin or acetylene.⁸⁾ We therefore investigated the reaction of PhCF_2Cl with PhSeNa in the presence of styrene or phenylacetylene to introduce the PhCF_2 unit into an unsaturated bond. However, the desired adduct of PhCF_2 and PhSe to styrene was not obtained but only PhSeCF_2Ph was produced; this means that the PhCF_2 radical reacted with PhSe^- (path b in Scheme 2) prior to styrene and phenylacetylene (path a in Scheme 2).

Samarium(II) iodide (SmI_2) was investigated as an alternative agent for one-electron reduction of PhCF_2Cl . SmI_2 has recently attracted much attention as a very useful and facile one-electron reducing reagent⁹⁾ and already been used




Ar = Phenyl (89%)
1-Naphthyl (84%)
2-Naphthyl (93%)

(Mechanism)



Scheme 1.




Scheme 2.


as a catalyst to promote the radical addition of perfluoroalkyl iodide to olefins and acetylenes.¹⁰ PhCF_2Cl underwent one-electron reduction with SmI_2 in the presence of HMPA to give the PhCF_2 radical and chloride ion, and the resulting PhCF_2 radical was trapped by styrene and phenylacetylene (Scheme 3).¹¹ 2-Dimethylaminoethanol (DMAE) was added as a proton source; in the absence of a proton source the yield of the adduct was very low and considerable amounts of polymeric compounds were produced. When a radical trapping reagent did not exist in the reaction system, PhCF_2H was obtained as a sole product. In the presence of CH_3OD , PhCF_2D was obtained almost selectively. This means that the reduction proceeds on the ionic path; the PhCF_2 radical is further

reduced with SmI_2 to a PhCF_2 anionic species (Scheme 3).

UHF/3-21G calculation demonstrated that reduction in PhCF_2Cl led the spontaneous and selective dissociation of the C–Cl bond to the PhCF_2 radical and Cl^- (Fig. 1). The selective reduction of the C–Cl bond in PhCF_2Cl has also been indicated by an electrochemical study using cyclic voltammetry; the difference in reduction potential between that of the C–Cl bond and that of the C–F bond was reported to be about 0.7 V, sufficiently for a selective reduction.⁷ On the contrary, benzyldiene trifluoride (PhCF_3) did not undergo one-electron reductive cleavage of the C–F bond with either PhSe^- or SmI_2 , and UHF/3-21G calculation also indicated that the PhCF_3 radical anion existed as a stable form without



(Mechanism)



Scheme 3.





Fig. 1. Energy diagram for dissociation of PhCF_2Cl radical anion calculated by UHF/3-21G; C-Cl bond distance is fixed and the other internal coordinates (bond distance, angle, and dihedral angle) are optimized for each of a series of specified C-Cl bond distances.



	neutral molecule	radical anion
C-F ¹ (Å)	1.352	1.377
C-F ² (Å)	1.352	1.377
C-F ³ (Å)	1.355	1.406
$\angle \text{CC}_\alpha\text{F}^3$ (deg)	112.9	118.4

$$E_{\text{HF}}[(\text{PhCF}_3)^{\bullet-}] = -563.1695 \text{ (Hartrees)}$$


$$E_{\text{HF}}(\text{PhCF}_2\bullet) + E_{\text{HF}}(\text{F}^-) = -563.0567 \text{ (Hartrees)}$$

Fig. 2. UHF/3-21G optimized structures of PhCF_3 and its radical anion.

the C-F bond dissociation (Fig. 2).

It is interesting to note that the optimized structure of the PhCF_2 radical is nonplanar. The optimized structures of the PhCF_2 radical, cation, and anion, and those of PhCH_2 calculated by UHF/3-21G are shown in Fig. 3.

The PhCH_2 radical, cation, and anion have planar structures and are stabilized by delocalization of non-bonding orbitals to the benzene ring. However, in the series of PhCF_2 , calculated structures of the radical and anion were nonplanar. The PhCF_2 cation is stabilized by resonance effect with a benzene ring as shown in Scheme 4 (A); meanwhile, in the PhCF_2 anion, repulsion between the anionic charge on the benzylic carbon and the lone-pair of fluorine (Scheme 4 (B)) leads the structure to a nonplanar form. The long carbon-fluorine bond (1.45 Å) in the PhCF_2 anion reflects the contri-



Scheme 4.

bution of the repulsion. In the PhCF_2 radical, the repulsion of the unpaired electron for the lone pair of fluorines also predominates, thus leading to a nonplanar structure, though the effect is less than in PhCF_2 anion. As the unpaired electron is localized on the benzylic carbon, the reactivity of the PhCF_2 radical is predicted to be higher than that of the PhCH_2 radical where the unpaired electron is delocalized to the benzene ring. In fact, the ArCF_2 radical was experimentally shown in our previous work to be moderately reactive and useful for the synthesis of ArCF_2 derivatives.⁵⁾

Thus, in this study, the importance of one-electron reduction of the C-Cl bond of ArCF_2Cl as a convenient route for the synthesis of ArCF_2 derivatives is shown experimentally and theoretically.

Experimental

Melting points were measured with Yanaco MP-500D melting point apparatus, and are uncorrected. ^1H , ^{13}C , and ^{19}F NMR spectra were taken with a JEOL JNM EX400 (400 MHz ^1H , 100 MHz ^{13}C , and 376 MHz ^{19}F NMR) spectrometer. Fluorine chemical shifts are given in ppm from external $\text{CF}_3\text{CO}_2\text{H}$. Mass spectra were obtained with a JEOL JMS AX-505W spectrometer with a JEOL JMA 5000 mass data system using an electron-impact (EI) ionization technique at 70 eV. Gel-permeation chromatography (GPC) was done with a JAI model LC-908 liquid chromatograph with two JAIGEL-1H columns (20 mm $\times$ 600 mm) with chloroform as eluent. Standard ab initio molecular orbital calculations were done with the GAUSSIAN 90 program¹²⁾ using IBM 3090/30J Computer.

Materials. PhCF_2Cl was prepared by the reaction of benzene with bis(chlorodifluoroacetyl) peroxide as described in our previous paper.⁴⁾ 1-(Chlorodifluoromethyl)- and 2-(chlorodifluoromethyl)naphthalenes were obtained by the reaction of naphthalene with bis(chlorodifluoroacetyl) peroxide, and the respective isomers were separated by GPC.⁴⁾ Diphenyl diselenide was available from Nacalai Tesque, Inc. and recrystallized from hexane before use. Samarium(II) iodide (SmI_2) was used as a THF solution (0.1 mol dm^{-3}); it was synthesized from samarium powder by the reaction with I_2 in THF under nitrogen according to the literature,¹³⁾ or obtained from Aldrich Co., Ltd., as a THF solution (0.1 mol dm^{-3}). The concentration of SmI_2 was measured by iodometry before use as described in the literature.¹³⁾

Reaction of ArCF_2Cl with PhSe^- . Diphenyl diselenide (0.4 mmol) was treated with NaBH_4 (1.2 mmol) in ethanol (0.2 cm^3) at room temperature under nitrogen to give PhSeNa , and then PhCF_2Cl (0.4 mmol) in DMF (3 cm^3) was added. The resulting solution was allowed to react at 100 $^\circ\text{C}$ under visible light. The reaction mixture was worked up with water (25 cm^3), and organic products were extracted with ether (20 $\text{cm}^3 \times 2$). The combined organic layer was washed with water, and then dried over MgSO_4 . After removal of the ether, the residue was purified by GPC to give PhCF_2SePh .

PhCF_2SePh : Colorless crystals (from hexane); mp 75.4—

76.1 °C; $^1\text{H NMR}$ (CDCl_3) δ = 7.22–7.66 (m, 10H); $^{13}\text{C NMR}$ (CDCl_3) δ = 122.40 (t, J_{CF} = 289 Hz), 125.00, 125.75, 128.27, 129.11, 129.48, 130.38, 136.94 (t, J_{CCF} = 20.1 Hz), 137.16; $^{19}\text{F NMR}$ (CDCl_3) δ = 5.4; MS m/z 284 (M^+ for ^{80}Se), 157, 128. Found: C, 55.30; H, 3.40%. Calcd for $\text{C}_{13}\text{H}_{10}\text{F}_5\text{Se}$: C, 55.14; H, 3.56%.

Similarly, the reactions of 1-(chlorodifluoromethyl)naphthalene and 2-(chlorodifluoromethyl)naphthalene with PhSeNa in DMF were done.

PhSeCF₂Np-1: Colorless crystals (from hexane); mp 75.0–76.0 °C; $^1\text{H NMR}$ (CDCl_3) δ = 7.29 (m, 2H), 7.38 (m, 2H), 7.52–7.65 (m, 5H), 7.89 (m, 2H), 8.56 (d, J = 8.3 Hz, 1H); $^{13}\text{C NMR}$ (CDCl_3) δ = 123.92 (t, J_{CCF} = 8.3 Hz), 124.19 (2C), 125.55, 125.77 (t, J_{CF} = 286.7 Hz), 126.08, 126.17, 128.64, 128.91, 129.00, 129.41, 131.56, 131.84 (t, J_{CCF} = 20.1 Hz), 134.01, 137.12; $^{19}\text{F NMR}$ (CDCl_3) δ = 8.8; MS m/z 334 (M^+ for ^{80}Se), 178, 128. Found: C, 61.25; H, 3.56%. Calcd for $\text{C}_{17}\text{H}_{12}\text{F}_2\text{Se}$: C, 61.27; H, 3.62%.

PhSeCF₂Np-2: Colorless crystals (from hexane); mp 127.6–130.0 °C; $^1\text{H NMR}$ (CDCl_3) δ = 7.28–7.49 (m, 2H), 7.38 (m, 1H), 7.49–7.67 (m, 5H), 7.79–7.91 (m, 4H); $^{13}\text{C NMR}$ δ = 122.00 (t, J_{CCF} = 3.7 Hz), 124.47 (t, J_{CF} = 294.4 Hz), 124.80 (t, J_{CCF} = 5.5 Hz), 125.79, 126.79, 127.45, 127.74, 128.33, 128.75, 129.08, 129.39, 132.22, 133.89, 134.05 (t, J_{CCF} = 23.8 Hz), 137.12; $^{19}\text{F NMR}$ (CDCl_3) δ = 5.5; MS m/z 334 (M^+), 178, 157, 127. Found: C, 61.09; H, 3.39%. Calcd for $\text{C}_{17}\text{H}_{12}\text{F}_2\text{Se}$: C, 61.27; H, 3.62%.

Reaction of PhCF_2Cl with SmI_2 in the Presence of Styrene and Phenylacetylene. To a solution of PhCF_2Cl in benzene (0.15 mol dm^{-3}) in the presence of HMPA (3 mol amt. to SmI_2), 2-PrOH, and styrene (25 mol amt. to PhCF_2Cl), a THF solution of SmI_2 (3–4 mol amt. to PhCF_2Cl) was added at room temperature under nitrogen. The reaction completed at once. PhCF_3 was added to the reaction mixture as an internal standard, and the yields of the products were measured by $^{19}\text{F NMR}$; PhCF_2Cl was completely consumed under these conditions, and the production of $\text{PhCF}_2\text{CH}_2\text{CH}_2\text{Ph}$ and PhCF_2H was recognized. However, PhCF_2Cl remained when a theoretical amount of SmI_2 (2 mol amt. to PhCF_2Cl) was used, and excess SmI_2 was required to consume PhCF_2Cl completely. After the workup with aq-HCl (1 mol dm^{-3}), organic products were extracted with ether ($3 \times 10 \text{ cm}^3$), and the combined organic layer was washed with water. The solution was dried over MgSO_4 , and then the solvent was removed. The purification of the residue by GPC gave pure $\text{PhCF}_2\text{CH}_2\text{CH}_2\text{Ph}$.

$\text{PhCF}_2\text{CH}_2\text{CH}_2\text{Ph}$: Colorless oil; $^1\text{H NMR}$ (CDCl_3) δ = 2.37–2.45 (m, 2H), 2.78 (t, J = 8.3 Hz, 2H), 7.15–7.20 (m, 2H), 7.25–7.29 (m, 2H), 7.42–7.45 (m, 3H), 7.49–7.52 (m, 2H); $^{13}\text{C NMR}$ (CDCl_3) δ = 28.77, 40.95 (t, J_{CCF} = 27.6 Hz), 124.93, 126.19, 128.23, 128.46, 128.97, 129.74, 137.16, 140.43; $^{19}\text{F NMR}$

(CDCl_3) δ = –20.8; MS m/z 232 (M^+), 127, 105. Exact MS: Found: m/z 232.1054. Calcd for $\text{C}_{15}\text{H}_{14}\text{F}_2$: 232.1063.

Similarly, the reaction in the presence of phenylacetylene was done. *E* and *Z* isomers of the adducts were separated by GPC.

***E*-PhCH=CHCF₂Ph:** Colorless oil; $^1\text{H NMR}$ (CDCl_3) δ = 6.44 (d, t, J_{HH} = 16.1 Hz, J_{HF} = 9.8 Hz, 1H), 6.82 (d, J = 16.1 Hz, 1H), 7.28–7.60 (m, 10H); $^{13}\text{C NMR}$ (CDCl_3) δ = 119.96 (t, J_{CF} = 239 Hz), 124.29 (t, J_{CCF} = 29.4 Hz), 125.52, 127.20, 128.44, 128.72, 129.05, 129.91, 134.28, 134.74, 136.66; $^{19}\text{F NMR}$ (CDCl_3) δ = –15.6; MS m/z 127 (M^+), 103. Exact MS: Found: m/z 230.0872. Calcd for $\text{C}_{15}\text{H}_{12}\text{F}_2$: M, 230.0907.

***Z*-PhCH=CHCF₂Ph:** Colorless oil; $^1\text{H NMR}$ (CDCl_3) δ = 6.01 (q, $J_{\text{HH}} = J_{\text{HF}}$ = 12.7 Hz, 1H), 6.83 (d, J_{HH} = 12.7 Hz, 1H), 7.25–7.55 (m, 10H); $^{13}\text{C NMR}$ (CDCl_3) δ = 119.36 (t, J_{CF} = 238 Hz), 125.44, 125.96 (t, J_{CCF} = 29.4 Hz), 127.80, 128.33, 129.10, 129.87, 134.92, 136.07, 137.17; $^{19}\text{F NMR}$ (CDCl_3) δ = –7.3.

References

- 1) “Biological Aspects of Fluorine Chemistry,” ed by R. Filler and Y. Kobayashi, Kodansha Ltd., Tokyo (1982); J. T. Welch, *Tetrahedron*, **43**, 3123 (1987).
- 2) K. Uneyama, *Yuki Gosei Kagaku Kyokai Shi (J. Synth. Org. Chem. Jpn.)*, **51**, 232 (1993); I. L. Knunyants and G. G. Yakobson, “Syntheses of Fluoroorganic Compounds,” Springer-Verlag, Berlin (1985).
- 3) Y. Kobayashi and I. Kumadaki, *Acc. Chem. Res.*, **11**, 197 (1978); C. Saboureaux, M. Troupel, S. Sibille, and Porichon, *J. Chem. Soc., Chem. Commun.*, **1989**, 1138; M. Kako, T. Morita, T. Torihara, and Y. Nakadaira, *J. Chem. Soc., Chem. Commun.*, **1993**, 678.
- 4) M. Yoshida, M. Ueda, Y. Morinaga, N. Kamigata, and M. Iyoda, *Chem. Lett.*, **1992**, 227; M. Yoshida, Y. Morinaga, and M. Iyoda, *J. Fluorine Chem.*, **68**, 33 (1994).
- 5) For PhS^- : A. E. Feiring, *J. Fluorine Chem.*, **24**, 191 (1984); V. N. Voiko, G. M. Shupak, and L. M. Yagupolskii, *J. Org. Chem., USSR (Engl. Transl.)*, **13**, 972 (1977); V. I. Popov, V. N. Boiko, N. V. Kondratenko, V. P. Sambur, and L. M. Yagupolskii, *J. Org. Chem., USSR (Engl. Transl.)*, **13**, 1985 (1977); V. N. Boiko, T. A. Dashevskaya, G. M. Shshupak, and L. M. Yagupolskii, *J. Org. Chem., USSR (Engl. Transl.)*, **15**, 347 (1979); V. N. Boiko, G. N. Shchapak, N. V. Ignat’ev, and L. M. Yagupolskii, *J. Org. Chem., USSR (Engl. Transl.)*, **15**, 1111 (1975); V. I. Popov, V. N. Boiko, and L. M. Yagupolskii, *J. Fluorine Chem.*, **21**, 365 (1982); I. Rico and C. Wakselman, *Tetrahedron*, **37**, 4209 (1981); I. Rico and C. Wakselman, *Tetrahedron Lett.*, **22**, 323 (1981); I. Rico, D.




Fig. 3. Optimized structures for cation, radical, and anion of PhCX_2 ($\text{X} = \text{F}$ and H) calculated by UHF/3-21G.

- Cantacuzene, and C. Wakselman, *J. Org. Chem.*, **48**, 1979 (1983); C. Wakselman and M. Tordeux, *J. Org. Chem.*, **50**, 4047 (1985); C. Wakselman and M. Tordeux, *J. Chem. Soc., Chem. Commun.*, **1984**, 793.
- 6) For PhSe^- : V. G. Volashchuk, V. N. Boiko, and L. M. Yagupolskii, *J. Org. Chem., USSR (Engl. Transl.)*, **13**, 1866 (1977).
- 7) Electrochemical reduction following C–Cl bond dissociation of PhCF_2Cl has been reported: H. Lund and N. J. Jensen, *Acta Chem. Scand., Ser. B*, **B28**, 263 (1974).
- 8) Y. Ueda, M. Kanai, and K. Uneyama, *Bull. Chem. Soc. Jpn.*, **67**, 2273 (1994); K. Uneyama and K. Kitagawa, *Tetrahedron Lett.*, **32**, 375 (1991); K. Uneyama and K. Kitagawa, *Tetrahedron Lett.*, **32**, 3385 (1991); K. Uneyama and M. Kanai, *Tetrahedron Lett.*, **32**, 7425 (1991).
- 9) J. Inanaga, *Yuki Gosei Kagaku Kyokai Shi (J. Synth. Org. Chem. Jpn.)*, **47**, 200 (1989); G. A. Molander, *Chem. Rev.*, **92**, 29 (1992); D. P. Curran, T. L. Fevig, C. P. Jasperse, and M. J. Tottleben, *Synlett*, **1992**, 943; J. A. Soderquist, *Aldrichimica*, **24**, 5 (1991).
- 10) X. Lu, S. Ma, and J. Zhu, *Tetrahedron Lett.*, **29**, 5219 (1988); S. Ma and X. Lu, *Tetrahedron*, **46**, 357 (1990).
- 11) The reactions of PhCF_2Cl with SmI_2 in the presence of various olefins and acetylenes have been investigated and the results were reported as preliminary communication: M. Yoshida, D. Suzuki, and M. Iyoda, *Chem. Lett.*, **1994**, 2357.
- 12) M. J. Frisch, M. Head-Gordon, G. W. Trucks, J. B. Foresman, K. Raghavachari, H. B. Schlegel, M. Robb, J. S. Binkley, C. Gonzalez, D. J. Defrees, D. J. Fox, R. A. Whiteside, R. Seeger, C. F. Melius, J. Baker, L. R. Kahn, J. J. P. Stewart, E. M. Fluder, S. Topiol, and J. A. Pople, "Gaussian 90," Gaussian, Inc., Pittsburgh, PA (1990).
- 13) T. Imamoto and M. Ono, *Chem. Lett.*, **1987**, 501.
-