

Characteristics of Phosphorous Components in Drainage Water

J. Lu

College of Resources and Environment, Northwest Sci-Tech University of Agriculture and Forestry, Yangling, Shaanxi 712100 People's Republic of China

Received: 31 August 2002/Accepted: 28 November 2003

Phosphorous (P) is an element for plant growth and its input has long been recognized as necessary to maintain profitable crop production. Phosphorous inputs can also increase the biological productivity of surface waters. Although nitrogen (N) and carbon (C) are essential to the growth of aquatic biota, most attention has focused on P inputs, because of the difficulty in controlling the exchange of N and C between the atmosphere and water and fixation of atmospheric N by some blue-green algae. Thus, P is often the limiting element and its control is of prime importance in reducing the accelerated eutrophication of fresh waters (Sharpley AN et al 1997). Since the late 1960s, point sources of water pollution have been reduced, due to their relative ease of identification and control. Even so, water-quality problems remain, and, as further point-source control becomes less cost-effective, attention is now being directed towards the contribution of agricultural P non-point sources to P in surface waters. Thus, a greater understanding of where P is coming from, how much P in soil and water is too much and how and where we can reduce these inputs and losses must be gained through research and extension programmes, in order to develop agricultural resource systems that sustain production and environmental quality, as well as farming communities. Phosphorus may be lost from agricultural land to water by several processes. These include erosion, surface runoff and subsurface flow (leaching) (Brookes PC et al. 1997). As most soils have a high absorption capacity for P, usually far exceeding the quantities of P added as manures or fertilizers (van Riemsdijk et al. 1987), it has long been considered that leaching losses of P from soil to water are negligible in most cases (Cooke 1976; Ryden et al. 1973; Lu Jialong et al. 1999).

Here I present data from drainage water of the Broadbalk Continuous Wheat Experiment at Rothamsted in UK (Johnston 1969), which question these findings.

MATERIALS AND METHODS

A detailed description of the Broadbalk experiment is given by Avery and

Bullock(1969). The silt loam and silty clay loam of the Batcombe series (Chromic Luvisol) together make up >90% of topsoil, which overlies clay-with-flints at < 60 cm, in turn resting on chalk. The experiment consists of 20 plots, 18 of which run the full length of the field and are 0.24 ha each. They receive different fertilizer treatments without replication. The experiment is divided, at right angles to the plots, into 10 sections, which test different agronomic treatments. Five are in continuous wheat, and five are in a rotation of fallow, potato and three successive wheat crops. Tile drains were installed along the middle of each plot in 1849 (1884 on the FYM+N2 strip) at a depth of about 65 cm, which opened into a ditch at one end of the experiment. In 1993 the old drains were replaced by perforated plastic tube drains on one section of continuous wheat nearest the ditch (section 9). Unlike the old drains, where the overlying clay had settled to the same bulk density as the soil mass, the new drains had a permeable gravel backfill up to 30 cm soil depth. The old drains were intercepted by a cross drain to avoid water from the rest of the experiment interfering with the new drains. Fertilizer treatments are listed in table 1. Chalk is applied every 5 year to maintain the pH at around 7.5 (Table 1). Water flowing from drains, 65 cm under soil surface, between winter 2000 and early summer 2001 was collected and analyzed for total and dissolved P. My aim was to see if significant quantities of P were lost through drainage and what are the percents of different phosphorus forms.

Drainage water samples were collected in Winchester bottles at the drain outlets on five occasions between December 2000 and April 2001. In each case, water was collected after the drains had been running freely for a few hours. The entire time of collection usually took about 1 hr. Drainage water was filtered through

Table 1. Details of the treatment and the soil from the Broadbalk experiment.

Plot no.	Treatment	Annotation
2.2	FYM	FYM: 35 t ha ⁻¹ farmyard manure N0-N5: supplying 0, 48, 96, 144, 192 and 240 kg N ha ⁻¹ both autumn and spring application. Supplying N to plots 17 and 18 in spring (April), to other plots in autumn (before ploughing) P: supplying 35 kg P ha ⁻¹ K: supplying 90 kg K ha ⁻¹ Na: supplying 35 kg Na ha ⁻¹ Mg: supplying 20 kg Mg ha ⁻¹ to plot 14, 35 kg Mg ha ⁻¹ every 3 rd year to other plots since 1974
3	NIL	
5	P K Mg	
6	N1 P K Mg	
7	N2 P K Mg	
8	N3 P K Mg	
9	N4 P K Mg	
10	N2	
11	N2 P	
12	N2 P Na	
13	N2 P K	
14	N2 P K Mg	
15	N5 P K Mg	
17	N1+N3, 1/2 [P K Mg]	
18	N0+N3, 1/2 [P K Mg]	

Table 2. Average concentrations of phosphorus and particulate matter in drainage water.

Plot	TP (mg/L)	TDP (mg/L)	MRP (mg/L)	DOP (mg/L)	PP (mg/L)	PM (g/L)
2.2	1.202	0.796±0.10	0.696±0.08	0.100±0.01	0.502	0.223
3	0.099±0.02	0.040±0.01	0.014±0.01	0.026±0.01	0.064±0.02	0.025±0.01
5	0.677±0.15	0.349±0.15	0.305±0.06	0.044±0.01	0.330±0.11	0.106±0.04
6	1.470±0.42	0.615±0.40	0.550±0.09	0.065±0.04	0.854±0.35	0.268±0.11
7	2.355±0.61	0.621±0.59	0.548±0.06	0.074±0.03	1.732±0.56	0.601±0.21
8	1.713±0.48	0.452±0.47	0.398±0.06	0.053±0.02	1.268±0.43	0.465±0.16
9	0.985±0.27	0.287±0.29	0.245±0.07	0.042±0.02	0.700±0.25	0.278±0.11
10	0.454±0.24	0.082±0.24	0.029±0.01	0.053±0.02	0.374±0.23	0.177±0.10
11	1.294±0.36	0.559±0.34	0.509±0.09	0.050±0.02	0.733±0.32	0.301±0.14
12	1.993	0.894	0.884	0.010	1.100	0.316
13	1.538±0.41	0.602±0.46	0.563±0.16	0.040±0.02	1.007±0.32	0.282±0.10
14	1.681±0.47	0.673±0.51	0.626±0.22	0.047±0.02	1.024±0.32	0.281±0.10
15	0.831±0.26	0.272±0.25	0.235±0.07	0.038±0.02	0.569±0.20	0.176±0.07
17	0.648±0.25	0.204±0.24	0.157±0.04	0.047±0.01	0.449±0.26	0.188±0.11
18	0.997±0.24	0.127±0.24	0.079±0.01	0.048±0.003	0.870±0.24	0.294±0.08

0.45- μ m Millipore membrane filters (mixed cellulose ester) with subsequent measurement of molybdate reactive phosphorus (MRP), i. e., mainly inorganic P, (Murphy and Riley, 1962) on the same day. The particulate matters under 0.45- μ m Millipore membrane filters were collected and dried in oven at 105-110 °C, then, weighted them on balance. The contents of particulate matter in drainage water were calculated. For further analysis, samples were stored in Winchester bottles at 5 °C in the dark. Total P in drainage water (TP), i. e., before filtration, and total dissolved P (TDP), i. e., after filtration, were determined after a modified perchloric acid digestion. An aliquot containing between 5 and 20 μ g P was digested with 3 ml of perchloric acid (60%) and 0.5 ml saturated $MgCl_2$ nearly to dryness, redissolved in 0.6 M HCl, neutralized with 5 M NaOH (Brookes and Powlson, 1981) and analyzed for P by the method of Murphy and Riley (1962). Total particulate P (PP) was calculated from the difference between TP and TDP and dissolved organic P (DOP) from the difference between TDP and MRP. All measurements are the means of three replicate determinations.

RESULTS AND DISCUSSION

Arithmetic means of total P (TP), molybdate reactive P (MRP), total dissolved P (TDP), dissolved organic P (DOP), total particulate P (PP) and particulate matter (PM) and the mean proportions of MRP, DOP, TDP and PP in TDP or TP of drainage waters from Broadbalk plots of 5 drainage events monitored between

Table 3. Average proportions MRP, DOP, TDP and PP in TDP or TP of drainage waters from Broadbalk.

Plot	TDP	PP	MRP	DOP	MRP	DOP
	Percentage of TP				Percentage of TDP	
2.2	58.24	41.76	48.50	9.73	86.99	13.01
3	45.04	59.53	13.22	31.82	52.34	47.66
5	58.89	41.42	47.92	10.96	83.79	16.21
6	51.42	48.42	48.05	3.37	90.91	9.09
7	32.74	67.15	29.42	3.32	88.60	11.40
8	34.49	65.66	31.40	3.09	89.11	10.89
9	39.63	60.56	32.86	6.77	80.28	19.72
10	41.09	60.70	11.31	29.79	49.62	50.38
11	56.98	42.89	52.07	4.91	90.65	9.35
12	44.86	55.19	44.36	0.50	98.88	1.12
13	48.91	53.91	43.28	5.63	89.22	10.78
14	48.74	51.77	43.37	5.37	85.20	14.80
15	46.05	57.29	38.93	7.12	82.42	17.58
17	57.94	44.67	42.67	15.28	72.50	27.50
18	18.87	81.13	12.34	6.54	61.56	38.44

December 2000 and April 2001 were tabulated in table 2 and table 3 respectively. There was substantial variation both in total P concentration and the relative contribution of the P fractions between the five events and between the different plots. The Broadbalk experiment does not have facilities for monitoring of the drainflows. Therefore results given have mainly represent snapshots. The Table 2 and Figure 2 had showed, On Broadbalk, the lowest total P concentration, 0.099 mg L⁻¹, for each event was generally measured on the plot which had never received fertilizer (the Nil plot) and ranged from 0.07 to 0.152 mg L⁻¹. The exception was the 6 April event in 2001 when lowest total P concentration in drainage water was measured on Plot 10. Despite receiving no P fertilizer inputs, drainage waters from Plot 10 (N2) had frequently higher total P concentrations, reaching 0.454 mg L⁻¹. This was mainly attributable to larger concentration. Thus, drainage water MRP from Plot 10 only varied between 0 to 0.043 mg L⁻¹, with an average concentration, 0.454 mg L⁻¹, slightly lower than Plot 17 (0.468 mg L⁻¹). The lowest MRP concentrations were always measured on the NIL plot.

The highest TP concentration in drainage water of 4.834 mg L⁻¹ was measured on the N2PKMg plot. Dominated by PP, frequently, TP concentrations ranged 0.354 to 4.076 mg L⁻¹, especially on Plots 6,7,8,12,13 & 14 (Table 2 and Figure 1). Similarly, the highest MRP concentrations were usually found on plots 2.2 (exception), 7, 12, 13, & 14, ranging from 0.548 to 0.884 mg L⁻¹ (Table 2 and Figure 1). Dissolved organic P (DOP) concentration, as determined by the




Figure 1. Phosphorus concentrations in drainage waters.

difference between total and molybdate reactive P in the $0.45\ \mu\text{m}$ filtrate of drainage waters, was ranged from 0.01 to $0.1\ \text{mg DOP L}^{-1}$. Most of them were low-remaining frequently below $0.05\ \text{mg L}^{-1}$ (Table 2). This is line with findings from other drainage experiment on similar soil types receiving inorganic P fertilizer (e.g. Culley 1983, Grant 1996, Ulén 1995) although they usually measured lower TP in general. A high mobility of DOP in soils is associated with large organic matter inputs (Table 2). Although DOP in drainage waters from the FYM plot tended to be higher than from plots receiving inorganic fertilizer (Table 2), the magnitude of DOP concentrations did not imply a significant contribution to leaching losses. The Broadbalk results are contrasted by findings on P fractions in the soil solution from an iron humus podzol. Ron Vaz (1993) found dissolved organic P to be the most significant fraction in soil solution below 10 cm soil depth on plots which had received up to $80\ \text{kg P ha}^{-1}\text{yr}^{-1}$ as superphosphate.

Table 2 and Table 3 illustrated the importance of P associated with Particulate matter (PP) in drainage waters and its treatment variation with the proportion of PP varying from 41.4% to 81.1% of the TP on plots. Particulate matter (PM) concentrations in drainage waters reached about $0.6\ \text{g L}^{-1}$ which is comparatively high for subsurface drainage systems (Pilgrim and Huff 1983, Skaggs 1994). The variability of PP concentrations in drainage waters was matched by the broad distribution of particulate matter concentration (PM) from different plots and drainage events. Accordingly, highly significant linear relationships ($P < 0.001$) were observed between PP and PM in drainage waters. Linear relationships between PP and PM in drainage waters were also reported from four agricultural drainage catchments on sandy loam in Denmark (Grant 1996). The proportion of




Figure 2. Mean proportions of MRP, DOP and PP in drains.

DOP, 1-50%, was lower than that of MRP, 50-99%, in TDP, including FYM plot. The proportions MRP, DOP and PP in total P (TP) and MRP and DOP in TDP were shown on Table 3 and Figure 2. Particulate P comprised by far the highest P concentrations in drainage waters on individual plots reaching occasionally 81% of the TP. The average proportions of PP, 54%, was higher than that of MRP, 36%, in drainage waters—opposite result was obtained by Heckrath GJ and Brookes PC (1998). Their conclusion was the average proportions of PP and MRP of the TP in drainage waters across all plots and drainage events were relatively similar. With 55 and 40% for MRP and PP, respectively, the dissolved P fraction was on the whole more important. The findings from Broadbalk are broadly in accordance with those of Culley (1983) who found on average 62% to 68% molybdate reactive P, 17% to 25% particulate P and 11% to 17% soluble organic P in tile drainage of a moderately fertilized clay soil in arable cropping. Phosphorus concentrations tended to be larger in the drainage water after the installation of the new drains in September 1993 (Heckrath G et al. 1995). The inevitable disturbance of soil during the installation might have caused enhanced P losses and so these must be considered with caution, especially as particulate P was generally greater than in earlier samplings. This would be consistent with the movement of topsoil into drains following disturbance. Due to the gravel backfill used, it is also possible that some water passing through a smaller cross section of soil reached the new drains more quickly, thus decreasing the potential for P retention by the soil.

Acknowledgment. I thank professor P C Brookes (IACR-Rothamsted, UK) for helpful direction and Dr. S Fortune (IACR-Rothamsted, UK) for helpful discussions. This work has been funded by IACR-Rothamsted International, UK.

The financial contribution of IACR-RI is gratefully acknowledgment. Presented at "First International Conference on Pollution Eco-chemistry and Processes", Shenyang, China, Aug 25-31, 2002.

REFERENCES

- Addiscott TM, Brockie D, Catt JA Christian DG, Harris GL, Howse KR, Mirza NA and Pepper TJ (2000) Phosphate losses through field drains in a heavy cultivated soil. *J Environ Qual* 29: 522-532
- Avery BW and Bullock P (1969) Morphology and classification of Broadbalk soils. P. 63-81. In Rothamsted Report for 1968. Part 2. The soils of Broadbalk. Rothamsted Exp. Stn., Harpenden, UK
- Brookes PC and Powlson DS (1981) Preventing phosphorus losses during perchloric acid digestion of sodium bicarbonate soil extracts. *J Sci Food Agric* 32:671-674
- Brookes PC, Heckrath G, De Smet J, Hofman G and Vanderdeelen J (1997) Losses of phosphorus in drainage water. In: Tunney H, Carton OT, Brookes PC and Johnston AE (eds.) *Phosphorus Loss from Soil to Water*. CAB International, UK, p253-271.
- Bridgham SD, Johnston CA, Schubauer JP and Weishampel P (2001) Phosphorus sorption dynamics and coupling with surface and pore water in riverine wetlands. *Soil Sci Soc America J* 65: 577-588.
- Cooke GW (1976) A review of the effects of agriculture on the chemical composition and quality of surface and underground waters. In *Agriculture and Water Quality*. MAFF Tech Bull 32 HMSO, London, UK, p5-57.
- Culley JLB, Bolton EF and Bernyk V (1983) Suspended soils and phosphorus loads from a clay soil: I. Plot studies *J Environ Qual* 12:493-498
- Edwards AC and Withers PJA (1998) Soil phosphorus management and water quality: a UK perspective. *Soil Use Manage* 14: 124-130
- Gburek WJ, Sharpley AN, Heathwaite L and Folmar GJ (2000) Phosphorus management at the watershed scale: A modification of the phosphorus index. *J Environ Qual* 29: 130-14
- Gillingham AG and Thorrold BS (2000) A review of New Zealand research measuring phosphorus in runoff from pasture. *J Environ Qual* 29: 88-96
- Grant R, Laubel A, Kronvang B Anderson HE, Svendsen LM and Fuglsang (1996) Loss of dissolved and particulate phosphorus from arable catchments by subsurface drainage. *Wat Res Res* 30: 2633-2642
- Haygarth PM and Sharpley AN (2000) Terminology for phosphorus transfer. *J Environ Qual* 29: 10-15
- Hesketh N and Brookes PC (2000) Development of an Indicator for risk of phosphorus leaching. *J Environ Qual* 29: 105-110

- Heckrath G, Brookes PC, Poulton PR and Goulding WT (1995) Phosphorus leaching from soils containing different phosphorus concentrations in the Broadbalk experiment. *J Environ Qual* 24: 904-910
- Hooda PS, Rendell AR, Edwards AC, Withers PJA, Aitken MN and Truesdale VW (2000) Relating soil phosphorus indices to potential phosphorus release to water. *J Environ Qual* 29: 1166-1171
- Hughes S, Reynolds B, Bell SA and Gardner C (2000) Simple phosphorus saturation index to estimate risk of dissolved P in runoff from arable soils. *Soil Use Manage* 16: 206-210
- Jorden C, McGuckin SO and Smith RV (2000) Increased predicted losses of phosphorus to surface waters from soils with high Olsen-P concentrations. *Soil Use Manage* 16: 27-35
- Laubel A, Jacobsen OH, Kronvang B, Grant R and Anderson HE (1999) Subsurface drainage loss of particles and phosphorus from field plot experiments and a tile-drained catchment. *J Environ Qual* 28: 576-584
- Lu JL, Zhang YP, Zhang JC and Su SP (1999) Studies on phosphorus transport in soils. *Acta Pedologica Sinica* 36:82-89
- McDowell RW and Sharpley AN. Approximating phosphorus release from soils to surface runoff and subsurface drainage. *J Environ Qual* 30: 508-520
- Morel C, Tunney H, Plénet D and Pellerin S (2000) Transfer of phosphate ions between soil and solution: Perspectives in soil testing. *J Environ Qual* 29: 50-59
- Murphy J and Riley JR (1962) A modified single solution method for the determination of phosphate in natural waters. *Anal Chim Acta* 27:31-36
- Pautler MC and Sins JT (2000) Relationships between soil test phosphorus, soluble phosphorus, and phosphorus saturation in Delaware soils. *Soil Sci Soc America J* 64:765-773
- Pilgrim DH and Huff DD (1983) Suspended sediment in rapid subsurface stormflow on a large field plot. *Earth Surf Proc Landforms* 8:451-463
- Riemsdijk WH van, Lexmond TM, Enfield CG and Zee SEATM van der (1987) Phosphorus and heavy metals: accumulation and consequences. In: HG van der Meer, RJ Unwin, TA van Dijk and GC Ennik (eds) *Animal Manure on Grassland and Fodder Crops. Fertilizer or Waste?* P.213-227. Martinus Nijhoff, Publishers, Dordrecht
- Ron Vaz ML, Edwards AC, Shand C and Cresser M (1993) Quantification of phosphorus fractions in soil solution. *Sci Tot Environ* 135:67-71
- Ryden JC, Syers JK and Harris RF (1973) Phosphorus in runoff and streams. *Adv Agron* 25:1-45
- Schoumans OF and Groenendijk P (2000) Modeling soil phosphorus levels and phosphorus leaching from agricultural land in the Netherlands. *J Environ Qual* 29: 111-116

- Sharpley AN, Chapra SC, Wedepohl R, Sims JT, Daniel TC and Reddy KR (1994) Managing agricultural phosphorus for protection of surface waters: issues and options. *J Environ Qual* 23: 437-451
- Sharpley AN, Foy B and Withers PJA (2000) Practical and innovative measures for the control of agricultural phosphorus losses to water: an overview *J Environ Qual* 29: 1-9
- Sharpley AN and Moyer B (2000) Phosphorus forms in manure and compost and their release during simulated rainfall. *J Environ Qual* 29: 1462-1467
- Sharpley AN and Tunney H (2000) Phosphorus research strategies to meet agricultural and environmental challenges of 21st century. *J Environ Qual* 29: 176-181
- Skaggs RW, Brevé MA and Gilliam JW (1994) Hydrologic and water quality impacts of agricultural drainage. *Crit Rev Environ Sci Technol* 24:1-32
- Siddique MT, Robinson JS and Alloway BJ (2000) Phosphorus reactions and leaching potential in soils amended with sewage sludge. *J Environ Qual* 29: 1931-1938
- Sui Y and Thompson ML (2000) Phosphorus sorption, desorption, and buffering capacity in a biosolids-amended mollisol. *Soil Sci Soc America J* 64:164-169
- Turner BL and Hayarth PM (2000) Phosphorus forms and concentrations in leachate under four grassland soil types. *Soil Sci Soc America J* 64:1090-1099
- Ulen B (1995) Episodic precipitation and discharge events and their influence on losses of phosphorus and nitrogen from tiledrained arable fields. *Swedish J Agr Res* 25:25-31
- Withers PJA, Davidson IA and Foy RH (2000) Prospects for controlling nonpoint phosphorus loss to water: A UK perspective. *J Environ Qual* 29: 167-175
- Withers PJA, Edwards AC and Foy RH (2001) Phosphorus cycling in UK agriculture and implications for phosphorus loss from soil. *Soil Use Manage* 17: 139-149