

ERRATUM

Iwao Iwasaki · Hisahiko Sugiyama
Shinsaku Kanazawa · Hiromichi Hemmi

Establishment of cisplatin-resistant variants of human neuroblastoma cell lines, TGW and GOTO, and their drug cross-resistance profiles

Published online: 29 August 2002
© Springer-Verlag 2002

Cancer Chemother Pharmacol (2002) 49:438–444

Table 1 (referred to on p 440) in the above article was inadvertently omitted. Table 1 is as follows:

Table 1. Cytotoxicity of anticancer agents against TGW and GOTO cell lines and their resistant variants. Cytotoxicity is expressed as the 50% inhibitory concentration (IC_{50} , μM) calculated from the growth inhibitory curves from at least three independent

experiments and are expressed as means $\pm$ SD. The cytotoxicity indexes were calculated using the expression: mean IC_{50} of the parental cell line/mean IC_{50} of the resistant cell line (*ND* not done)

Cell line	CDDP		ADR		MNNG		Vincristine		Ara-C		Chlorambucil	
	IC_{50}	Index	IC_{50}	Index	IC_{50}	Index	IC_{50}	Index	IC_{50}	Index	IC_{50}	Index
TGW	21.7 $\pm$ 3.6	1.0	14.2 $\pm$ 3.4	1.0	17.8 $\pm$ 2.3	1.0	1.53 $\pm$ 0.71	1.0	81.1 $\pm$ 10.2	1.0	64.1 $\pm$ 8.6	1.0
TR1	45.0 $\pm$ 5.2	2.1*	16.9 $\pm$ 6.1	1.2	26.7 $\pm$ 9.9	1.4	1.73 $\pm$ 0.5	1.1	74.0 $\pm$ 15.3	0.91	66.1 $\pm$ 5.6	1.03
TR2	34.3 $\pm$ 2.9	1.6*	19.9 $\pm$ 2.2	1.4*	37.5 $\pm$ 0.7	2.2*	2.33 $\pm$ 0.59	2.1	64.0 $\pm$ 11.5	0.79	75.0 $\pm$ 9.6	1.17
TR3	40.8 $\pm$ 3.7	1.9*	17.2 $\pm$ 4.6	1.2	41.6 $\pm$ 5.3	2.5*	2.9 $\pm$ 1.19	1.9	80.5 $\pm$ 12.8	0.98	68.6 $\pm$ 5.2	1.07
GOTO	25.7 $\pm$ 5.8	1.0	16.4 $\pm$ 7.9	1.0	15.1 $\pm$ 3.4	1.0	1.2 $\pm$ 0.87	1.0	81.3 $\pm$ 15.2	1.0	79.0 $\pm$ 5.4	1.0
GR1	39.5 $\pm$ 2.0	1.5*	19.3 $\pm$ 2.1	1.2	16.5 $\pm$ 0.7	1.1	1.0 $\pm$ 0.14	0.8	73.6 $\pm$ 20.6	0.91	70.4 $\pm$ 6.2	0.89
GR2	43.3 $\pm$ 5.5	1.7*	17.8 $\pm$ 2.9	1.1	21.6 $\pm$ 3.4	1.5	0.8 $\pm$ 0.46	0.7	ND		ND	
GR3	28.2 $\pm$ 2.1	1.1	ND		17.1 $\pm$ 4.0	1.1	ND		ND		ND	

* $P < 0.01$ vs respective parental cell line, Student's *t*-test

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00280-002-0452-4>

I. Iwasaki · H. Sugiyama · S. Kanazawa · H. Hemmi (✉)
Department of Molecular Biology,
Toho University School of Medicine,
5-21-16 Ohmori-Nishi, Ohta-Ku, Tokyo 143-8540, Japan
E-mail: hhemmi@med.toho-u.ac.jp
Tel.: +81-3-37624151
Fax: +1-3-37646923