

Imatinib mesylate enhances the malignant behavior of human breast carcinoma cells

Germana Rappa · Fabio Anzanello · Aurelio Lorico

Received: 10 March 2010 / Accepted: 20 June 2010 / Published online: 2 July 2010
© Springer-Verlag 2010

Abstract

Purpose Imatinib mesylate (Imatinib), clinically employed for chronic myeloid leukemia and gastrointestinal stromal tumors, is a selective inhibitor of the tyrosine kinases, c-abl, c-kit and PDGFRs. Due to the frequent expression of these genes in breast cancer cells, the clinical efficacy of Imatinib has recently been investigated in patients with advanced and metastatic breast cancer. Here, we have studied the effects of Imatinib on human MA-11 breast carcinoma cells, expressing both c-abl and PDGFRbeta, in vitro and in mouse xenografts.

Methods The effects of Imatinib mesylate on the human MA-11 breast carcinoma cell line were studied in vitro and in xenografts.

Results Daily intraperitoneal treatment with 60 mg/kg Imatinib for 9 days of athymic nude mice pre-implanted subcutaneously with MA-11 cells did not result in an anti-tumor effect, but rather increased the take rate of 3×10^4 cells from 30.8 to 84.6% and caused the appearance of large abdominal masses in 30% of mice. To investigate the mechanism(s) of the observed effects of Imatinib on MA-11 tumors, we exposed the cells in vitro to Imatinib for 9 days. The surviving population, expanded in culture, showed increased motility and over-expressed a set of genes associated with aggressive behavior. Also, several genes belonging to the Wnt and the MAPK pathway were differentially

expressed. In promoter activation assays, Imatinib increased the promoter activity driven by both Wnt and MAPK/ERK-1/2.

Conclusions Our data suggest caution in the clinical use of Imatinib in breast cancer patients; the comparison of Imatinib-surviving breast cancer cells with parental cells may help define the regulatory pathways involved in the increased malignancy of residual tumor cells that survive therapy, ultimately providing important therapeutic targets.

Keywords Breast cancer · Imatinib · Cancer stem cells · CD24 · Microarray

Introduction

Every year, more than 1 million women are newly diagnosed with breast cancer worldwide [1]. One of the most exciting areas of improvement in the treatment of breast cancer is the development of novel targeted and biological therapeutic agents, such as Imatinib mesylate (Imatinib), a 2-phenylaminopyrimidine derivative that specifically inhibits the tyrosine kinase enzymes, c-Abl, PDGFR, and c-Kit [2, 3]. Imatinib was first developed to specifically inhibit Bcr-Abl, a fusion protein with kinase activity present in chronic myeloid leukemia (CML) patients. It has been widely used for treatment of patients with CML and gastrointestinal stromal tumors and is under investigation in several malignancies, including breast cancer [4, 5]. Also, potentiation of chemotherapy by Imatinib in breast carcinoma cell lines has recently been reported [6].

Recent evidence suggests that breast cancer originates from and is maintained by a sub-population of stem cell-like cells, named cancer-initiating cells or cancer stem cells (CSC), able to differentiate along different pathways.

Electronic supplementary material The online version of this article (doi:10.1007/s00280-010-1394-x) contains supplementary material, which is available to authorized users.

G. Rappa · F. Anzanello · A. Lorico (✉)
Department of Drug Development, Nevada Cancer Institute,
One Breakthrough Way, Las Vegas, NV 89135, USA
e-mail: alorico@nvcancer.org

The identification and characterization of breast CSC has important diagnostic, prognostic and therapeutic implications [4–7]. Originally, breast CSC were described as CD44⁺CD24^{−/low} cells.

In the present study, we have employed as breast cancer model the human MA-11 breast carcinoma cell line, established from bone marrow micrometastases of a breast cancer patient. We have previously reported [7] that in MA-11 cells CD24 expression ranges from 0 to 1,120,000 molecules/cell, with an average number of 214,000 molecules/cell. All cells in the MA-11 line express CD44, ranging from 900,000 to 8,800,000/cell, with an average number of 4,100,000 molecules/cell. Therefore, the MA-11 line is composed of both CD44⁺CD24^{−/low} and CD44⁺CD24^{high} sub-populations. Upon in vitro culture as mammospheres under stem cell-like conditions, MA-11 cells acquire increased tumorigenicity and display reduced levels of CD24 compared with parental adherent cells [22]. However, the level of CD24 expression is neither a stable feature of mammosphere-forming cells nor confers tumorigenic potential to MA-11 cells; in fact, upon passage of MA-11 mammospheres into serum-supplemented cultures, CD24 expression is restored; the rapid increase in CD24 expression is consistent with up-regulation of the antigen, and not with in vitro selection of CD24⁺ cells [7]. Also, the tumorigenicity of sorted CD44⁺CD24[−] and CD44⁺CD24^{high} MA-11 cells is equal. Single cell-sorted CD24[−] and CD24^{high} MA-11 give rise in vitro to cell populations with heterogeneous CD24 expression; subcutaneous tumors derived from sorted CD24[−] sub-populations and single-cell clones have levels of CD24 expression similar to the unsorted cells, suggesting that the level of expression of CD24 is associated with but does not contribute to tumorigenicity [7]. We have also previously reported [7] that CSC-enriched MA-11 mammospheres have activated specific signal transduction pathways, including MAPK and Wnt.

Here, we have studied the effects of Imatinib on MA-11 cells in vitro and in vivo and report that treatment with Imatinib results in increased tumorigenicity and malignancy of MA-11 cells.

Materials and methods

Cell culture

MA-11 cells were cultured in RPMI-1640 (Mediatech Inc. Herndon, VA) added with 10% fetal bovine serum (Atlanta Biologicals, Lawrenceville, GA) at 37°C in a 5% CO₂ humidified incubator. Cells were used between passages 10 and 20 and tested routinely for mycoplasma contamination.

Tumor implantation

Animal studies were done under a protocol approved by the Institutional Animal Care and Use Committee. For s.c. injection, cells were resuspended in 100 µl PBS and injected subcutaneously into the right flank of the animals.

Wound healing assay

To perform the wound healing assay, a wound in a confluent MA-11 cell monolayer was introduced using a pipette tip. The monolayers were observed over a time course of 3–24 h.

Flow cytometric analysis

Mouse anti-human CD24 (clone ML5) and anti-human CD44 (clone G44-26) monoclonal antibodies were from BD Biosciences (San Jose, CA). Cell surface CD24 expression was quantitated using the quantum simply cellular system (Bangs Laboratories, Fishers, IN), as previously described [8]. And 5×10^5 cells/sample were incubated with saturating concentrations (10 µg/ml) of antibodies for 30 min at 4°C. Standard curves of beads with fixed antibody-binding capacity and samples were analyzed on a FACS Vantage flow cytometer (BD Biosciences, San Jose, CA). Antibody-binding capacity for each cell population was calculated using the QuickCal v.2.3. software (Bangs Laboratories), employing median histogram values and linear regression analyses.

Microarray analysis

Total RNA was isolated from Imatinib- and mock-treated MA-11 cells using trizol phenol-based extraction. The quality of the RNA was assessed by gel electrophoresis and A₂₆₀/A₂₈₀ ratio. Microarray analyses were performed by Ocean Ridge Biosciences (ORB, Palm Beach Gardens, FL) using human exonic evidence-based oligonucleotide (HEEBO) microarrays (<http://alizadehlab.stanford.edu/>) containing approximately 50,000 70-mer probes complementary to constitutive exons of most human genes, as well as alternatively spliced exons, ESTs, and control sequences. Biotinylated UTP complementary RNA (cRNA) probes were prepared, fragmented and hybridized to the microarrays for 16–18 h with constant rotation. The microarray slides were washed under stringent conditions, stained with Streptavidin-Alexa-647 (Invitrogen), and scanned using an Axon GenePix 4000B scanner. For data analyses, the local background was subtracted, and the spot intensities were log₂-transformed. The spot intensities were then normalized by subtracting the 70th percentile of spot intensity of the probes against human constitutive exons and adding

back a scaling factor (grand mean of 70th percentiles). After removing data for low quality spots, control sequences, and non-human probes, 41,213 human probe intensities remained. The human probes intensities were filtered to identify all probes with intensity above a normalized threshold [$\log 2(3 \times \text{standard deviation of raw local background}) + \text{mean of } \log 2\text{-transformed negative controls}$], to arrive at 23,291 probes above threshold in both samples from at least one group. For statistical analysis, samples were binned in two groups (Imatinib treated and control). The $\log 2$ -transformed and normalized spot intensities for the detectable probes were examined for differences between the treatment groups by one-way ANOVA using National Institute of Aging (NIA) Array Analysis software. This ANOVA was conducted using the Bayesian Error Model and 20 degrees of freedom. Gene ontology analysis was performed using GenMAPP software (Gladstone Institute, San Francisco, CA) for all detectable probes with current Entrez Gene IDs. Specifically, the MAPPfinder module of GenMAPP was first used to map all detectable probes, based on their gene targets, to GO and Local MAPP categories. Then, MAPPfinder compared the relative representation in each functional group of genes associated with probes meeting one of five differential expression criteria to the relative representation of genes associated with the full set of detectable probes. Significance was determined by permutation of Z scores with correction for multiple comparisons as described in the GenMAPP software manual. To shed light on the Imatinib-specific signaling pathways, all differentially expressed genes linked to GO terms identified in the microarray were subjected to pathway analysis using Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis.

Transduction of MA-11 cells with lentiviral reporter gene constructs

MA-11 cells were transduced with VSV-pseudotyped lentiviral particles expressing inducible reporter constructs encoding the luciferase gene under the control of a basal promoter element, joined to tandem repeats of a specific transcriptional response element (signal finder lenti cancer 10-pathway reporter array kit, from SABiosciences, Frederick, MD). For transduction of MA-11 cells, lentiviral supernatants were preloaded onto recombinant fibronectin (Retronectin, Takara Shuzo, Japan)-coated plates and centrifuged at 950g for 30 min at 4°C. The operation was repeated a second time with fresh supernatant. The supernatant was then removed, and the plates washed with PBS before addition of MA-11 cells in their complete culture medium. After transduction, stable cell lines expressing the specific transcriptional response elements were isolated via selection with 2 µg/ml puromycin. Pathway-specific

MA-11 cells were seeded at a density of 2×10^4 per well in 24-well plates and grown to 50–60% confluence. Cells were then treated with Imatinib or with the solvent for 24 h or for 9 days. The preparation of cell extracts and measurement of luciferase activities were carried out using the Steady-Glo Luciferase Reporter Assay System according to recommendations by the manufacturer (Promega, Madison, WI). The assays for luciferase activity were done in a 20/20 luminometer (Turner Biosystems, Sunnyvale, CA). The ratio of luciferase activity between Imatinib- and mock-treated samples was then calculated.

Results

Effects of Imatinib on MA-11 xenografts

To investigate the activity of Imatinib on breast cancer xenografts, we implanted 30,000 MA-11 cells s.c. in athymic nude mice and treated the animals i.p. with 60 mg/kg Imatinib diluted in phosphate-buffered saline for nine consecutive days, starting the day after the implantation of the tumor cells. Imatinib did not have an anti-tumor effect, rather increased the tumorigenic potential of MA-11 cells and in 30% of animals caused the formation of large tumor masses in the intraperitoneal cavity. Table 1 shows the take rates in control and Imatinib-treated animals (30.8 and 84.6%, respectively). The average time to reach 1 cm³ was 83.5 days for control animals and 41.1 days for animals exposed to Imatinib. To investigate whether a similar effect on the growth of MA-11 xenografts occurred with a different anti-cancer agent, we employed the chemotherapeutic drug, paclitaxel, on an additional cohort of mice. Mice were exposed to nine doses of 10 mg/kg paclitaxel for 9 days. As shown in Table 1, no difference in the take rate was observed, with only a minor increase in time required to

Table 1 In vivo effects of i.p. Imatinib and paclitaxel on the formation of MA-11 tumors in athymic s.c. implanted nude mice

Parameter	Control	Imatinib	Paclitaxel
Take rate (%)	30.8 (<i>n</i> = 13)	84.6* (<i>n</i> = 13)	33.3 (<i>n</i> = 6)
Time to 1 cm ³ tumor (days)	83.5 ± 24.5 ^a	41.1* ± 4.7 ^a	101
Presence of an abdominal tumor mass (%)	0 (<i>n</i> = 13)	30* (<i>n</i> = 13)	0 (<i>n</i> = 6)

Mice were implanted with 3×10^4 MA-11 cells s.c. and treated daily i.p. with 60 mg/kg Imatinib or 10 mg/kg paclitaxel for 9 days starting the day after the implantation

* *P* < 0.05, Wilcoxon two-tailed test

^a SD




Fig. 1 In vivo effects of Imatinib treatment on MA-11 tumors. **a** Athymic nude mice implanted s.c. with MA-11 cells and treated with nine daily doses of 60 mg/kg Imatinib or with the solvent alone. **b** Hematoxylin-eosin staining of an abdominal tumor mass from a nude mouse injected s.c. with Imatinib-surviving cells

reach 1 cm³ (83.5 vs. 101 days). A representative experiment with two control and two Imatinib-treated mice is shown in Fig. 1a. An hematoxylin-eosin-stained section from an abdominal MA-11 mass, showing a very cellular tumor, is represented in Fig. 1b.




Fig. 2 Imatinib-induced changes in CD24 expression on the surface of MA-11 cells. MA-11 cells were exposed to 25 μM Imatinib for 72 h, followed by incubation in drug-free medium for up to 48 h. After labeling of MA-11 cells with an anti-CD24-PE monoclonal antibody,

Reversible changes in the CD44⁺CD24^{−/low} phenotype of MA-11 cells by Imatinib

Originally, breast CSC were described as CD44⁺CD24^{−/low} cells. We have previously reported [7] that in MA-11 cells the level of CD24 expression is neither a stable feature of CSC-enriched cells grown as mammospheres nor confers tumorigenic potential to the cells. We now found that upon treatment of MA-11 cells with Imatinib, CD24 expression on the plasma membrane was almost completely lost in 3 days (Fig. 2). The average number of CD24 molecules on the surface of MA-11 cells, measured by flow cytometry, employing phycoerythrin-conjugated anti-human CD24 monoclonal antibodies and the quantum simply cellular system, decreased from 214,000 to 7,000 molecules/cell. Instead, CD44 did not change significantly with Imatinib treatment, with the average number of CD44 molecules/cell going from 4,100,000 to 4,000,000 after 72 h exposure to Imatinib. But 24 and 48 h after removal of Imatinib from the culture medium, the average number of CD24 molecules/cell increased very rapidly to 21,000 and 45,000, respectively (Fig. 2). Microarray analysis (data not shown) confirmed that 4 days after removal of Imatinib from the culture medium, CD24 expression was similar to that of control mock-treated cells.

In vitro study of the Imatinib-surviving fraction of MA-11 cells

To investigate the mechanism of the effects of Imatinib on MA-11 cells, we simulated the in vivo treatment by exposing MA-11 cells in vitro for 9 days to 25 μM Imatinib. After 9 days, the great majority of cells died. After growth and expansion in culture of the surviving cells for four additional days in the absence of Imatinib, we compared

cells were analyzed by flow cytometry for CD24 expression. P2 gates include only the CD24⁺ cells based on their respective isotype controls

Imatinib-treated cells and the parental cell line for biological properties, chemosensitivity and gene expression pattern.

First, we noticed that the motility of the cells increased, as measured by a conventional wound healing assay [9]. After 24 h, MA-11 cells pre-treated with Imatinib polarized toward the wound, initiated protrusion, migrated, and closed the wound faster than mock-treated cells (Suppl. Fig. 1).

We then investigated the pattern of chemosensitivity of Imatinib-surviving cells by an MTS/PMS microtiter plate assay. Cells were exposed to the investigational drugs for 72 h. The level of resistance to Imatinib was twofold, while no changes in sensitivity to the cytotoxic activity of etoposide, paclitaxel, doxorubicin, antimony potassium tartrate and chlorambucil were observed (Table 2 and Suppl. Fig. 2).

To investigate whether specific genes and/or signaling pathways were involved in the increased tumorigenicity and invasiveness of Imatinib-surviving MA-11 cells, we analyzed Imatinib-induced changes in gene expression profile. By HEEBO microarray analysis of the gene expression profiles of two independent cultures each for control and Imatinib-surviving MA-11 cells, we found 877 probes showing significant differences with $P < 0.05$ and FDR (false discovery rate) $< 5\%$ (Suppl. Table 1). Excluding the non-annotated genes and multiple probes, only 301 and 241 genes were found to be significantly up- and down-regulated, respectively (Suppl. Table 2). In all cases where multiple probes were available for the 542 genes, the changes in expression were very similar. First, we searched for Imatinib target genes, whose over-expression could cause resistance to Imatinib. We found that c-kit and pdgfr-alpha were not expressed in MA-11 cells, abl and pdgfr-beta were equally expressed between parental and Imatinib-treated cells, while CACNA1G, a T-type calcium channel, recently

identified as an additional target of Imatinib, unrelated to the inhibition of tyrosine kinases [10], was threefold over-expressed in Imatinib-treated cells (Suppl. Table 3).

Although no difference in the cytotoxic activity of several chemotherapeutic agents was observed, 35 transporters were found differentially regulated in Imatinib-surviving versus control cells, including ABCC3 (MRP3), ABCC6 (MRP6), and several SLC (solute carrier) genes. The over-expression of SLC22A3 and ABCC3 was confirmed by RT-PCR (Table 3).

To shed light on Imatinib-induced or repressed signaling pathways, all genes differentially expressed in the microarray were subjected to pathway analysis and assigned to pathway networks. In particular, eight genes of the Wnt pathway were up-regulated, and five and six genes of the MAPK pathway were respectively up- and down-regulated in Imatinib-surviving cells (Suppl. Tables 4, 5). We then looked for genes associated with tumorigenicity, invasiveness, and/or the breast cancer stem cell phenotype, and we found that Imatinib-treated cells overexpress many genes associated with malignancy of breast cancer cells, including the mitochondrial benzodiazepine receptor TSPO, neurotensin, MMP-9, the inhibitor of cell death, COP1, and the aldehyde dehydrogenase gene, ALDH3A1, associated with the breast cancer stem cell phenotype [11] (Suppl. Table 6). We then employed RT-PCR to confirm the differential expression of ten representative over-expressed genes. A good correlation between microarray and RT-PCR data was observed for all ten genes (Table 3).

To further investigate the effects of Imatinib on the WNT/ β -catenin and the MAPK pathways, we analyzed the transcriptional activation of promoters driven by Wnt and MAPK proteins, using luciferase as reporter. To this aim, we transduced MA-11 cells with TCF/LEF and SRE lentiviral reporters, as well as with lentiviral reporters for eight additional cancer-related pathways, namely Notch, p53, TGF β ,

Table 2 Drug sensitivity profile of Imatinib-surviving MA-11 cells (Imatinib)

Drug	IC50 control (mean $\pm$ SD)	IC50 Imatinib (mean $\pm$ SD)
Imatinib	20 $\pm$ 4	40 $\pm$ 5*
Etoposide	1.2 $\pm$ 0.3	1.3 $\pm$ 0.2
Paclitaxel	0.05 $\pm$ 0.01	0.06 $\pm$ 0.02
Doxorubicin	0.2 $\pm$ 0.07	0.15 $\pm$ 0.06
Antimony	6 $\pm$ 1	5 $\pm$ 1.5
Chlorambucil	500 $\pm$ 150	600 $\pm$ 100

MA-11 cells were exposed to 25 μ M Imatinib for 9 days. After 4 days in drug-free medium, their sensitivity to a panel of drugs was compared with that of mock-treated cells by an MTT assay. IC50 values are expressed in μ M concentrations, except for Imatinib (μ g/ml)


SD standard deviation

* $P > 0.05$, ANOVA test

Table 3 Fold-increase in expression for 10 genes over-expressed in Imatinib-surviving MA-11 cells by microarray and RT-PCR

Entrez gene ID	Entrez symbol	Microarray fold change	RT-PCR fold change
22943	NTS	6.9	10.0
4804	DACT1	4.0	5.0
4897	FAIM2	2.5	10.0
7045	SLC22A3	4.9	2.5
28513	ABCC3	2.2	3.3
1000	COP1	2.2	2.5
558	CASP9	1.9	2.0
3725	PCDHGA11	5.2	5.0
4035	CACNA1G	2.2	1.7
51339	MMP9	1.6	1.3

Fig. 3 Imatinib enhances promoter activation driven by Wnt and MAPK proteins. MA-11 cells were transduced with ten different lentiviral reporters (signal finder lenti cancer 10-pathway reporter array kit, from SABiosciences). After puromycin selection, each reporter cell line was exposed to 25 μ M Imatinib for 9 days, followed by a 4-day recovery period (**a**) or for 24 h (**b**). The activity of the different pathways was analyzed by a luciferase assay and expressed as the ratio between the luciferase activity of Imatinib-treated and mock-treated MA-11 cells. Data points are the mean of three separate experiments; bars SD; * $P < 0.05$ (ANOVA test)



pRB-E2F, NFkB, Myc, Hypoxia, and JNK. After puromycin selection, we exposed each reporter cell line to a short treatment (24 h) or a long treatment (9 + 4 days recovery) with 25 μ M Imatinib and analyzed the activity of the different pathways by a luciferase assay. Figure 3 shows that in Imatinib-surviving cells only the Wnt/ β -catenin, the MAPK/Erk1-2, and the Myc pathways were activated (Fig. 3a), while after 24-h exposure to Imatinib no significant changes could be observed for any of the ten pathways investigated (Fig. 3b).

Discussion

Despite novel therapeutic strategies for advanced and metastatic breast cancer, many patients relapse after an initial

response to conventional and/or experimental drug treatments. In most cases, local and distant recurrences are associated with a more aggressive phenotype and resistance to therapy. It has been suggested that a small subpopulation of “tumor-initiating cells” or “cancer stem cells” (CSC) are resistant to therapy and hence may reinitiate tumor growth after treatment [1]. Chemotherapy-induced selection of CSC has been reported in several studies [12, 13]. However, the CSC that have survived first-line therapy should re-establish a recurrent cancer that is similar to the original cancer both in biological behavior and susceptibility to first-line therapy.

We have found that treatment of MA-11 cells in vitro with Imatinib results in a surviving population of cells with increased motility, that over-express a set of genes

associated with tumorigenic and invasive behavior. In vivo, nude mice subcutaneously implanted with MA-11 cells and treated with Imatinib developed more tumors at low levels of inoculum and developed with high frequency abdominal tumor masses. There are three possible non-mutually exclusive explanations for our findings: (1) Imatinib selects a sub-population of breast CSC; (2) Imatinib induces a more malignant gene expression pattern; (3) Imatinib exerts an immunosuppressive activity in vivo, thus facilitating tumor engraftment.

While selection of a CSC population could explain our in vitro data, it is not sufficient to justify our data of increased tumorigenicity and malignancy in immunodeficient mice. Also, microarray analysis showed that stem cell markers or cancer stem cell markers were either not expressed in MA-11 cells or not differentially expressed: in particular, ALDH1, Notch3, CD133, Rex-1, Oct-4, Nanog were not expressed, and Sox-2 was equally expressed in Imatinib-treated and control MA-11 cells. In vitro treatment with Imatinib resulted in decreased expression of CD24, a marker negatively associated with the breast CSC phenotype. This is in agreement with observations by other groups of a decrease in CD24 expression in patients treated with neoadjuvant chemotherapy [14], and that the CD44⁺CD24^{-low} phenotype can be induced by genetic events or epigenetic events, such as exposure to TGF- β , a factor secreted by tumor-associated stroma [15]. However, the rapidity of changes in CD24 expression observed in the present study was consistent with changes in the expression level of the antigen, not with selection of a CD24^{-low} sub-population. This is in agreement with our previous findings that the CD44⁺CD24^{-low} phenotype of MA-11 cells is reversible and unrelated to tumorigenicity and metastatic potential [7].

Our in vivo findings may be attributed to epigenetic changes induced by the drug treatment, i.e. Imatinib-surviving cells activate some cell machinery that results in expression of a set of pro-malignant genes and consequently become endowed with pro-malignant properties. Alternatively, the in vivo effects of Imatinib on MA-11 xenografts could be mediated, at least in part, by an immunosuppressive effect of the drug cannot be excluded. Imatinib can adversely affect the immune system by blocking immune surveillance of the malignancy and permitting disease recurrence. In fact, Imatinib reportedly inhibits activation, differentiation, and proliferation of normal T lymphocytes [16], antigen presentation by dendritic cells [17], and TNF α secretion [18]. Also, Cristofanilli et al. [4] recently reported Imatinib-mediated inhibition of interferon-gamma production by TCR-activated CD4⁺ T cells in patients with PDGFR- β -expressing advanced breast cancer.

By microarray analysis, we found that resistance to Imatinib of the surviving MA-11 cell population could be

attributed to increased expression of the CACNA1G gene, a T-type calcium inhibitor recently identified as an additional target of Imatinib, and/or over-expression of the MRP3 transporter [19, 20]. The observed changes in expression for 35 SLC genes, which typically mediate uptake of nutrients (glucose, amino acids, organic anions, and peptides), may be related to increased energy and nutrition needs of cells with increased malignancy.

Interestingly, several genes potentially associated with the malignant phenotype of Imatinib-surviving cells were found to be over-expressed. Among these, TSPO, a mitochondrial permeability transition-pore protein involved in the regulation of apoptosis and cell proliferation that potentiates proliferation, motility, and transmigration capabilities of cancer cells [21]; COP1 inhibitor of cell death that binds caspase 1 and caspase 4, blocking apoptosis [22]; MMP-9, involved in the breakdown of the extracellular matrix and in the metastatic process; and Neurotensin, a trophic and anti-apoptotic factor associated with breast cancer progression that interacts with the neurotensin receptor on the cell membrane and stimulates Rho GTPases [23, 24]. Interestingly, many RhoGTPases were also over-expressed in Imatinib-surviving cells, and a link between neurotensin and MMP-9 has recently been demonstrated in breast cancer cells [23]. We have also found several signal transduction pathways, potential targets for anti-breast cancer therapy, differentially expressed in the Imatinib-surviving population versus the parental cell line. Among these, the Wnt/ β -Catenin and the MAPK pathway, both recently found by our group to be altered in CSC-enriched MA-11 mammospheres compared with parental MA-11 cells [7]. The association of significant changes in the Wnt/ β -Catenin and the MAPK pathway with increased malignancy of breast cancer cells is supported by recent studies showing that aberrant activation of Wnt signaling induces mammary tumors from stem/progenitor cells [25], and that activation of the classical MAPK/ERK pathway, leading to c-ras activation, is associated with breast cancer stem cells and breast tumor epithelial-mesenchymal transition [26, 27]. Our data suggest caution in the clinical use of Imatinib in breast cancer patients; the comparison of Imatinib-surviving and parental breast cancer cells may help understand the role of differentially expressed genes in tumorigenicity and invasiveness and define the regulatory pathways involved in the increased malignancy of residual tumor cells that survive therapy, ultimately providing important therapeutic targets.

Acknowledgments The authors thank Dr. Willoughby for help with the interpretation of microarray data. The project described was supported by Award Number R01CA133797 (G.R.) from the National Cancer Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Cancer Institute or the National Institutes of Health.

References

- Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ (2007) Cancer statistics, 2007. *CA Cancer J Clin* 57:43–66
- Druker BJ, Tamura S, Buchdunger E, Ohno S, Segal GM, Fanning S, Zimmermann J, Lydon NB (1996) Effects of a selective inhibitor of the Abl tyrosine kinase on the growth of Bcr-Abl positive cells. *Nat Med* 2:561–566
- Joensuu H, Dimitrijevic S (2001) Tyrosine kinase inhibitor imatinib (STI571) as an anticancer agent for solid tumours. *Ann Med* 33:451–455
- Cristofanilli M, Morandi P, Krishnamurthy S, Reuben JM, Lee BN, Francis D, Booser DJ, Green MC, Arun BK, Puzstai L, Lopez A, Islam R, Valero V, Hortobagyi GN (2008) Imatinib mesylate (Gleevec) in advanced breast cancer-expressing C-Kit or PDGFR-beta: clinical activity and biological correlations. *Ann Oncol* 19:1713–1719
- Chew HK, Barlow WE, Albain K, Lew D, Gown A, Hayes DF, Gralow J, Hortobagyi GN, Livingston R (2008) A phase II study of imatinib mesylate and capecitabine in metastatic breast cancer: Southwest oncology group study 0338. *Clin Breast Cancer* 8:511–515
- Sims JT, Ganguly S, Fiore LS, Holler CJ, Park ES, Plattner R (2009) STI571 sensitizes breast cancer cells to 5-fluorouracil, cisplatin and camptothecin in a cell type-specific manner. *Biochem Pharmacol* 78:249–260
- Rappa G, Lorico A (2010) Phenotypic characterization of mammosphere-forming cells from the human MA-11 breast carcinoma cell line. *Exp Cell Res* 316(9):1576–1586
- Rappa G, Fodstad O, Lorico A (2008) The stem cell-associated antigen CD133 (Prominin-1) is a molecular therapeutic target for metastatic melanoma. *Stem Cells* 26:3008–3017
- Lampugnani MG (1999) Cell migration into a wounded area in vitro. *Methods Mol Biol* 96:177–182
- Cataldi M, Gaudino A, Lariccia V, Russo M, Amoroso S, di Renzo G, Annunziato L (2004) Imatinib-mesylate blocks recombinant T-type calcium channels expressed in human embryonic kidney-293 cells by a protein tyrosine kinase-independent mechanism. *J Pharmacol Exp Ther* 309:208–215
- Moreb JS (2008) Aldehyde dehydrogenase as a marker for stem cells. *Curr Stem Cell Res Ther* 3:237–246
- Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong C, Huang Y, Hu X, Su F, Lieberman J, Song E (2007) Let-7 regulates self renewal and tumorigenicity of breast cancer cells. *Cell* 131:1109–1123
- Levina V, Marrangoni AM, DeMarco R, Gorelik E, Lokshin AE (2008) Drug-selected human lung cancer stem cells: cytokine network, tumorigenic and metastatic properties. *PLoS One* 3:e3077
- Li X, Lewis MT, Huang J, Gutierrez C, Osborne CK, Wu MF, Hilsenbeck SG, Pavlick A, Zhang X, Chamness GC, Chamness GC, Wong H, Rosen J, Chang JC (2008) Intrinsic resistance of tumorigenic breast cancer cells to chemotherapy. *J Natl Cancer Inst* 100:672–679
- Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, Brooks M, Reinhard F, Zhang CC, Shipitsin M, Campbell LL, Polyak K, Briskin C, Yang J, Weinberg RA (2008) The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 133:704–715
- Cwynarski K, Laylor R, Macchiarulo E, Goldman J, Lombardi G, Melo JV, Dazzi F (2004) Imatinib inhibits the activation and proliferation of normal T lymphocytes in vitro. *Leukemia* 18:1332–1339
- Seggewiss R, Price DA, Purbhoo MA (2008) Immunomodulatory effects of imatinib and second-generation tyrosine kinase inhibitors on T cells and dendritic cells: an update. *Cytotherapy* 10:633–641
- Wolf AM, Wolf D, Rumpold H, Ludwiczek S, Enrich B, Gastl G, Weiss G, Tilg H (2005) The kinase inhibitor imatinib mesylate inhibits TNF- α production in vitro and prevents TNF-dependent acute hepatic inflammation. *Proc Natl Acad Sci USA* 102:13622–13627
- Hu S, Franke RM, Filipinski KK, Hu C, Orwick SJ, de Bruijn EA, Burger H, Baker SD, Sparreboom A (2008) Interaction of imatinib with human organic ion carriers. *Clin Cancer Res* 14:3141–3148
- Radich JP, Dai H, Mao M, Oehler V, Schelter J, Druker B, Sawyers C, Shah N, Stock W, Willman CL, Friend S, Linsley PS (2006) Gene expression changes associated with progression and response in chronic myeloid leukemia. *Proc Natl Acad Sci USA* 103:2794–2799
- Rechichi M, Salvetti A, Chelli B, Costa B, Da Pozzo E, Spinetti F, Lena A, Evangelista M, Rainaldi G, Martini C, Gremigni V, Rossi L (2008) TSPO over-expression increases motility, transmigration and proliferation properties of C6 rat glioma cells. *Biochim Biophys Acta* 1782:118–125
- Wang X, Narayanan M, Bruey JM, Rigamonti D, Cattaneo E, Reed JC, Friedlander RM (2006) Protective role of Cop in Rip2/caspase-1/caspase-4-mediated HeLa cell death. *Biochim Biophys Acta* 1762:742–754
- Souaze F, Dupouy S, Viardot-Foucault V, Bruyneel E, Attoub S, Gespach C, Gompel A, Forgez P (2006) Expression of neurotensin and NT1 receptor in human breast cancer: a potential role in tumor progression. *Cancer Res* 66:6243–6249
- Dupouy S, Viardot-Foucault V, Alifano M, Souaze F, Plu-Bureau G, Chaouat M, Lavour A, Hugol D, Gespach C, Gompel A, Forgez P (2009) The neurotensin receptor-1 pathway contributes to human ductal breast cancer progression. *PLoS One* 4:e4223
- Lindvall C, Bu W, Williams BO, Li Y (2007) Wnt signaling, stem cells, and the cellular origin of breast cancer. *Stem Cell Rev* 3:157–168
- Dreesen O, Brivanlou AH (2007) Signaling pathways in cancer and embryonic stem cells. *Stem Cell Rev* 3:7–17
- Liu M, Casimiro MC, Wang C, Shirley LA, Jiao X, Katiyar S, Ju X, Li Z, Yu Z, Zhou J, Johnson M, Fortina P, Hyslop T, Windle JJ, Pestell RG (2009) p21CIP1 attenuates Ras- and c-Myc-dependent breast tumor epithelial mesenchymal transition and cancer stem cell-like gene expression in vivo. *Proc Natl Acad Sci USA* 106:19035–19039