

A multi-institution phase II study of gemcitabine/S-1 combination chemotherapy for patients with advanced biliary tract cancer

Masashi Kanai · Kenichi Yoshimura · Takehiko Tsumura · Masanori Asada · Chihiro Suzuki · Miyuki Niimi · Shigemi Matsumoto · Takafumi Nishimura · Takashi Nitta · Kentaro Yasuchika · Kojiro Taura · Yukiko Mori · Akihiko Hamada · Naoya Inoue · Shinsuke Tada · Kazuhiro Yanagihara · Shujiro Yazumi · Yukio Osaki · Tsutomu Chiba · Iwao Ikai · Masanori Fukushima · Shinji Uemoto · Etsuro Hatano

Received: 25 May 2010 / Accepted: 20 August 2010 / Published online: 2 September 2010
© Springer-Verlag 2010

Abstract

Purpose We aimed to evaluate the efficacy and safety of gemcitabine/S-1 combination chemotherapy for the treatment of patients with advanced biliary tract cancer.

Methods Patients with histologically or cytologically confirmed unresectable or recurrent biliary tract cancer were eligible for inclusion. The primary endpoint was overall survival. Gemcitabine was administered intravenously at a dose of 1,000 mg/m² over 30 min on days 1 and 8, and oral S-1 was administered daily at a dose of 60 mg/m² on days

1–14. This schedule was repeated every 3 weeks until disease progression or patient refusal.

Results Twenty-five patients were enrolled between October 2007 and January 2009. Eleven patients (44%) had extrahepatic bile duct cancer, 5 (20%) had intrahepatic bile duct cancer, 8 had gallbladder cancer (32%), and 1 (4%) had ampulla of Vater cancer. The median overall survival time was 12.7 months (95% CI, 8.4–23.5 months), and the 1-year survival rate was 52.0% (95% CI, 31.2–69.2%). Of the 23 patients with evaluable target regions, seven patients experienced a partial response, and an overall response rate was 30.4%. The following grade 3–4 hematological toxicities occurred: neutropenia (56%), leukopenia (24%), anemia (8%) and thrombocytopenia (4%). In spite of the high incidence of grade 3–4 neutropenia, no patients developed febrile neutropenia in the present study. The major grade 3–4 non-hematological toxicities were fatigue (8%), anorexia (8%) and diarrhea (4%).

Conclusions Gemcitabine/S-1 combination chemotherapy offered a promising survival benefit with acceptable toxicity in patients with advanced biliary tract cancer.

M. Kanai (✉) · S. Matsumoto · T. Nishimura · Y. Mori · K. Yanagihara · T. Chiba
Outpatient Oncology Unit, Kyoto University Hospital,
54 Shogoin-Kawahara-cho Sakyo-ku, Kyoto, Japan
e-mail: kanai@kuhp.kyoto-u.ac.jp

K. Yoshimura · C. Suzuki · M. Niimi · M. Fukushima
Translational Research Center,
Kyoto University Hospital, Kyoto, Japan

T. Tsumura · Y. Osaki
Osaka Red Cross Hospital, Osaka, Japan

M. Asada · S. Yazumi
Kitano Hospital, Osaka, Japan

T. Nitta · K. Yasuchika · K. Taura · I. Ikai · S. Uemoto · E. Hatano
Department of Surgery, Graduate School of Medicine,
Kyoto University Hospital, Kyoto, Japan

A. Hamada
Kyoto Katsura Hospital, Kyoto, Japan

N. Inoue
Kansai Denryoku Hospital, Osaka, Japan

S. Tada · T. Chiba
Department of Gastroenterology and Hepatology,
Kyoto University Hospital, Kyoto 606-8507, Japan

Keywords Biliary tract cancer · Gemcitabine · S-1 · Chemotherapy

Introduction

Biliary tract cancer is one of the most lethal malignancies worldwide, with surgery representing the only potentially curative treatment for this disease. However, many patients are diagnosed too late for curative resection, and even if surgery can be performed, the likelihood of relapse is very high [7, 13]. Over the past decade, gemcitabine has been widely used to treat unresectable or recurrent biliary tract

cancer [3, 4, 9, 17, 18, 23, 27], although no phase III trials have established this drug as a standard treatment for advanced biliary tract cancer. We have previously evaluated the outcome of consecutive 22 patients with advanced biliary tract cancer who received gemcitabine monotherapy as first line and reported that median survival time (MST) was 8.3 months (95% CI 6.4–11.2 months) [9].

In the ABC-02 study, the first prospective multicenter phase III study in this field, gemcitabine/cisplatin combination chemotherapy was compared with gemcitabine monotherapy. The study found that the combination regimen significantly prolonged MST (from 8.1 to 11.7 months; $P < 0.001$) [26]. The superiority of gemcitabine/cisplatin combination chemotherapy over gemcitabine monotherapy was also demonstrated in a randomized phase II study conducted in Japan (the BT-22 study) [6]. Given these findings, gemcitabine/cisplatin combination chemotherapy is now becoming accepted as a new standard regimen for advanced biliary tract cancer.

S-1 is an oral fluoropyrimidine prodrug that has confirmed efficacy against various solid tumors, both alone and in combination with other cytotoxic drugs [1, 12, 14, 19, 29]. S-1 monotherapy has yielded good results against advanced biliary tract cancer [5, 24], and gemcitabine/S-1 combination therapy has yielded promising results with acceptable toxicity levels for patients with advanced pancreatic cancer [15, 16, 28]. At the time of planning this clinical trial in 2007, there had been no reports on gemcitabine/S-1 combination chemotherapy for patients with advanced biliary tract cancer, so we designed this clinical trial to determine its efficacy and safety in this context.

Patients and methods

Eligibility criteria

Patients with advanced biliary tract cancer that was not amenable to potentially curative surgery or that had recurred after surgery were eligible for inclusion if they met the following criteria: histologically or cytologically confirmed biliary tract cancer; Eastern Cooperative Oncology Group performance status of 0–2; age ≥ 20 years; adequate bone marrow function (neutrophil count $\geq 1,500/\text{mm}^3$, and platelet count $\geq 100,000/\text{mm}^3$), liver function (total bilirubin ≤ 3 times the upper limit of normal (ULN) and aspartate aminotransferase [AST]/alanine aminotransferase [ALT] ≤ 5 times ULN) and renal function (creatinine ≤ 1.5 mg/dL); adequate oral intake; life expectancy ≥ 3 months. All patients provided written informed consent. Exclusion criteria included a history of chemotherapy or radiotherapy (patients who had undergone adjuvant chemotherapy were not excluded if at least 6 months had passed since the last

administration), pregnancy or lactation, a history of severe drug allergy and other severe comorbid diseases. This phase II study (UMIN ID 000000792) was conducted in five institutions in Japan. The protocol was approved by the institutional review board at each institution, and patient registration and data management were conducted at an independent data center at Translational Research Center, Kyoto University Hospital. All procedures were performed in accordance with the 1964 Declaration of Helsinki.

Treatment

Gemcitabine was infused at a dose of $1,000 \text{ mg/m}^2$ over 30 min on days 1 and 8. S-1 was given orally twice a day for 14 consecutive days. Doses of S-1 were calculated according to body surface area (BSA) as follows: $\text{BSA} < 1.25 \text{ m}^2$, 60 mg/day; $1.25 \text{ m}^2 \leq \text{BSA} < 1.5 \text{ m}^2$, 80 mg/day; $\text{BSA} \geq 1.5 \text{ m}^2$, 100 mg/day. The gemcitabine and S-1 treatment regimen was repeated every 3 weeks. Doses were reduced in response to adverse effects (graded according to the Common Terminology criteria for Adverse Events v 3.0) [22].

Chemotherapy was started if on day 1 the neutrophil count was $\geq 1,500/\text{mm}^3$, platelet count was $\geq 100,000/\text{mm}^3$, total bilirubin was ≤ 3 times the ULN, AST/ALT was ≤ 5 times ULN, and there were no non-hematological toxicities of grade 3 or higher (except for abnormal blood test results not relevant to the chemotherapy regimen). Chemotherapy was continued if on day 8 the neutrophil count was $\geq 1,000/\text{mm}^3$, platelet count was $\geq 75,000/\text{mm}^3$, total bilirubin was ≤ 3 times the ULN, AST/ALT was ≤ 5 times the ULN, and there were no non-hematological toxicities of grade 3 or higher. If the patient did not meet the above criteria, chemotherapy was delayed by 1 week. If neutropenia (grade 3–4), thrombocytopenia (grade 3–4), febrile neutropenia or non-hematological toxicity associated with gemcitabine (grade 3) occurred, the subsequent gemcitabine dose was reduced to 800 mg/m^2 . If further toxicity occurred with the reduced dose, it was further reduced to 600 mg/m^2 . If a further dose reduction was necessary, the subsequent gemcitabine dose was reduced by 20%. If diarrhea or stomatitis (grade 3–4) associated with S-1 occurred, S-1 was discontinued, and patients were withdrawn from the study. No dose reescalation was allowed. The treatment regimen was continued until disease progression, unacceptable toxicity including non-hematological toxicity of grade 4 or patient refusal occurred.

Pretreatment and follow-up evaluation

Pretreatment evaluation included obtaining the patient's medical history and performing a physical examination, imaging using contrast-enhanced computed tomography

(CT) or magnetic resonance imaging (MRI), a complete blood cell count, serum biochemical tests, an electrocardiogram and chest X-rays. During the treatment cycles, physical examinations and blood tests were scheduled on days 1 and 8. Carcinoembryonic antigen (CEA) and carbohydrate antigen 19-9 (CA19-9) were measured at the time patients were enrolled in the study and every month thereafter. Toxicity was evaluated using the Common Terminology criteria for Adverse Events v3.0.

Statistical analysis

The primary endpoint was overall survival. The secondary endpoints were toxicity and response rate. Twenty-five patients were enrolled, a sample size that would allow rejection of a null hypothesis of a 30% 1-year survival rate and acceptance of an alternative hypothesis of a 50% 1-year survival rate, with a significance level of 0.05 and a power of 80%. Overall survival was calculated using the Kaplan–Meier method and was defined as the time from initiation of therapy to death from any cause or the final follow-up. Among patients with measurable target lesions, the objective response rate was evaluated according to Response Evaluation Criteria in Solid Tumors (RECIST) version 1.0 [21]. Patients were enrolled between October 2007 and January 2009, and the final analysis was conducted in January 2010 after a 1-year follow-up period. All analyses were conducted on an intention-to-treat basis and were performed using SAS version 9.2 (SAS Institute, Cary, NC, USA).

Results

Patient characteristics

Twenty-five patients were enrolled between October 2007 and January 2009. The patient characteristics are shown in Table 1. The median age was 63 years (range 32–78 years), and 18 patients (72%) were men. Four of the 25 patients (16%) experienced recurrent disease after undergoing curative surgery. Out of the 21 patients with unresectable disease, distant metastasis was reported in 13 patients at the time of enrollment. Eleven patients (44%) had extrahepatic bile duct cancer, 8 had gallbladder cancer (32%), 5 (20%) had intrahepatic bile duct cancer, and 1 (4%) had ampulla of Vater cancer.

Efficacy

Seventeen patients (68%) died during the study period. The median overall survival time was 12.7 months (95% CI, 8.4–23.5 months), and the 1-year overall survival rate was

Table 1 Characteristics of patients with advanced biliary tract cancer ($n = 25$)

Sex	
Male	18 (72.0%)
Female	7 (28.0%)
Median age (years)	63 (range 32–78)
Primary lesion	
Intrahepatic	5 (20.0%)
Extrahepatic	11 (44.0%)
Gallbladder	8 (32.0%)
Ampulla of Vater	1 (4.0%)
Disease status	
Unresectable	21 (84.0%)
Recurrent*	4 (16.0%)
Target lesion	
Primary	18
Liver	7
Lymph node	3
Peritoneum	2
Local recurrence	2
Lung	1
None	2
Median no. treatment cycles	7 (range 1–20)
Median CEA (ng/mL)	4.5 (range 0.3–468)
Median CA19-9 (U/mL)	167 (range 1–6,373)

CEA carcinoembryonic antigen, CA 19-9 carbohydrate antigen

* One patient had a history of adjuvant chemotherapy using gemcitabine

52% (95% CI, 31.2–69.2%, $P = 0.02$ under a null hypothesis of 30%). Of the 23 patients with target regions that were evaluable according to RECIST, 7 (30.4%) experienced a partial response, and 13 (56.5%) had stable disease, with an overall disease control rate of 87.0%.

Toxicity

In total, 229 cycles of gemcitabine/S-1 combination chemotherapy were delivered, with a median of 7 cycles per patient (range 1–20 cycles; Table 1). The mean relative dose intensities of gemcitabine and S-1 were 75% and 84%, respectively. The incidence rates of hematological and non-hematological adverse events are summarized in Tables 2 and 3, respectively. The most common grade 3–4 hematological toxicity was neutropenia (56%); however, no instances of febrile neutropenia were observed in this study. The incidence rates of grade 3–4 anemia and thrombocytopenia were 8% and 4%, respectively. Grade 3–4 hyperbilirubinemia and ALT were observed in 16% and 8% of patients, respectively, mostly associated with obstructive jaundice caused by the primary disease. Other grade 3–4

Table 2 Hematological adverse events among patients with advanced biliary tract cancer treated with gemcitabine/S-1 combination chemotherapy ($n = 25$)

	Grade 1	Grade 2	Grade 3	Grade 4	Incidence of grade 3–4 events (%)
Neutropenia	0	5	12	2	56
Leukopenia	1	9	6	0	24
Anemia	5	7	1	1	8
Thrombocytopenia	1	4	1	0	4
Febrile neutropenia	–	–	0	0	0

Table 3 Non-hematological adverse events among patients with advanced biliary tract cancer treated with gemcitabine/S-1 combination chemotherapy ($n = 25$)

	Grade 1	Grade 2	Grade 3	Grade 4	Incidence of grade 3–4 events (%)	Incidence of grade 1–4 events (%)
Fatigue	8	3	2	0	8	52
Anorexia	3	2	2	0	8	28
Diarrhea	1	4	1	0	4	24
Constipation	1	6	0	0	0	28
Rash	9	3	0	0	0	48
Fever	8	3	0	0	0	44
Hand-foot rash	7	3	0	–	0	40
Infection-other	8	2	0	0	0	40
Nausea	3	2	0	0	0	20
Stomatitis	5	1	0	0	0	24
Allergic reaction	4	1	0	0	0	20
Hyperpigmentation	8	0	–	–	0	32
Alopecia	3	0	–	–	0	12
Injection site reaction	2	0	0	–	0	8
Vomiting	1	0	0	0	0	4
Hyperbilirubinemia	3	1	4	0	16	32
AST	11	5	0	0	0	64
ALT	8	4	2	0	8	56
Creatinine	3	0	0	0	0	12

AST aspartate aminotransferase,
ALT alanine aminotransferase

non-hematological adverse events were fatigue (8%), anorexia (8%) and diarrhea (4%).

Discussion

In our population of patients with advanced biliary tract cancer, gemcitabine/S-1 combination chemotherapy achieved an MST of 12.7 months and a 1-year survival rate of 52%. The MST for patients with gall bladder cancer ($n = 8$) was shorter (7.6 months) than that for patients with other cancer types (16.0 months), which is consistent with the findings of previous studies and possibly reflects the more aggressive nature of gall bladder cancer [8, 10, 20]. The proportion of patients with gall bladder cancer in our study (32%) was comparable with the proportions in previous randomized trials (26–39%) [6, 25, 26], so the good MST observed in the current study was unlikely to be simply due

to tumor-type selection bias. Furthermore, this was a multi-institution trial, and the eligibility criteria were almost identical to the indications used for administering chemotherapy in daily clinical practice; both these factors are likely to have contributed to reducing selection bias. Although comparing single-arm phase II studies can be problematic, our current results are comparable to those of Sasaki et al., who observed an MST of 11.6 months and a 1-year survival rate of 44% among patients with advanced biliary tract cancer treated with gemcitabine/S-1 combination chemotherapy [20] (Table 4). Their treatment schedule differed slightly from ours: it consisted of 1,000 mg/m² gemcitabine on days 1 and 15 and 80 mg/m² S-1 daily for 14 consecutive days every 4 weeks. In this study, grade 3–4 neutropenia was observed in 56% of patients, and this often caused suspension of chemotherapy on day 8. In fact, planned chemotherapy administration on day 8 needed to be suspended in 28.5% of cycles. Meanwhile, Sasaki et al.

Table 4 Results of clinical trials of gemcitabine and oral fluoropyrimidine combination chemotherapy for the treatment of advanced biliary tract cancer

	Present study	Sasaki et al. [20]	Knox et al. [10]	Cho et al. [2]	Koeberle et al. [11]
Oral fluoropyrimidine	S-1	S-1	Capecitabine	Capecitabine	Capecitabine
MST (months)	12.7	11.6	14	14	13.2
1-year survival rate (%)	52	44	49	58	N/A
Prevalence of gall bladder cancer (%)	32	40	49	16	18
Incidence of grade 3–4 neutropenia (%)	56	34	34	11	11 ^a
Incidence of grade 3–4 anorexia (%)	8	3	N/A	2	7
Incidence of grade 3–4 fatigue (%)	8	N/A	4	0	11
Sample size	25	35	45	44	44

MST median survival time, N/A not available

^a These subjects had leukopenia

reported grade 3–4 neutropenia was 34%, and their regimen might have an advantage of avoiding suspension of chemotherapy due to neutropenia because gemcitabine administration was scheduled on day 1 and 15, not on day 8 with 2-week interval. Interestingly, a previous study of gemcitabine/cisplatin combination therapy in Japanese patients (BT-22 study) yielded a 56.1% incidence rate of grade 3–4 neutropenia, whereas in ABC-02 study involving the same regimen, the rate was only 22.6% among Caucasian patients [6, 26]. Although we need to take into account the difference of treatment duration between 2 studies (up to 24 weeks in ABC-02 study versus up to 48 weeks in BT-22 study), it is tempting to speculate that ethnic differences exist between patients with biliary tract cancer in terms of susceptibility to gemcitabine-related neutropenia. In spite of the high incidence of grade 3–4 neutropenia in the present study, no patients developed febrile neutropenia, probably due to the short duration of neutropenia caused by this combination therapy. Aside from AST/ALT elevation, the most common non-hematological toxicity was fatigue (52%); however, the incidence rates of grade 3–4 toxicities were relatively low, showing that this regimen was generally well tolerated in an outpatient setting. The grade 3–4 hyperbilirubinemia observed in this study was associated with obstructive jaundice caused by the primary disease and so was unlikely to be relevant to the combination therapy regimen. In vitro study also demonstrated the advantage of gemcitabine/S-1 combination. Yoshizawa et al. tested the combination of S-1 with other anticancer drugs (gemcitabine, cisplatin, irinotecan, mitomycin C, adriamycin and paclitaxel) and reported that synergic effect was most evident in gemcitabine/S-1 combination [30]. The combination of gemcitabine and another oral fluoropyrimidine, capecitabine, was found to be similarly efficacious in previous single-arm phase II studies [2, 10, 11]. In their respective studies, Cho et al. [2] observed an MST of 14 months and a 1-year survival rate of 58%, and Knox

et al. [10] also observed an MST of 14 months and a 1-year survival rate of 49%. Koeberle et al. [11] found similar results, with an MST of 13.2 months (Table 4). Koeberle et al. also highlighted the importance of maintaining a balance between treatment efficacy and quality of life in palliative chemotherapy for advanced biliary tract cancer. From the point of view of quality of life, combination therapy using oral fluoropyrimidines has the major advantage of being very convenient to administer. Clearly, we must be cautious about the interpretation of data from single-arm phase II studies; however, the combination of gemcitabine and oral fluoropyrimidines can be used for patients with advanced biliary tract cancer in situations that preclude the use of cisplatin (e.g., allergy to cisplatin). In summary, gemcitabine/S-1 combination chemotherapy yielded a promising survival benefit with acceptable toxicity in patients with advanced biliary tract cancer. We believe that this regimen would be a good candidate for the experimental arm of a future phase III trial of gemcitabine/cisplatin combination therapy.

Acknowledgments This work was supported by the Smoking Research Foundation. We thank Hiroe Tada for her valuable support.

References

1. Boku N, Yamamoto S, Fukuda H, Shirao K, Doi T, Sawaki A, Koizumi W, Saito H, Yamaguchi K, Takiuchi H, Nasu J, Ohtsu A (2009) Fluorouracil versus combination of irinotecan plus cisplatin versus S-1 in metastatic gastric cancer: a randomised phase 3 study. *Lancet Oncol* 10:1063–1069
2. Cho JY, Paik YH, Chang YS, Lee SJ, Lee DK, Song SY, Chung JB, Park MS, Yu JS, Yoon DS (2005) Capecitabine combined with gemcitabine (CapGem) as first-line treatment in patients with advanced/metastatic biliary tract carcinoma. *Cancer* 104:2753–2758
3. Eckel F, Schmid RM (2007) Chemotherapy in advanced biliary tract carcinoma: a pooled analysis of clinical trials. *Br J Cancer* 96:896–902

4. Eng C, Ramanathan RK, Wong MK, Remick SC, Dai L, Wade-Oliver KT, Mani S, Kindler HL (2004) A Phase II trial of fixed dose rate gemcitabine in patients with advanced biliary tree carcinoma. *Am J Clin Oncol* 27:565–569
5. Furuse J, Okusaka T, Boku N, Ohkawa S, Sawaki A, Masumoto T, Funakoshi A (2008) S-1 monotherapy as first-line treatment in patients with advanced biliary tract cancer: a multicenter phase II study. *Cancer Chemother Pharmacol* 62:849–855
6. Furuse J, Okusaka T, Miyazaki M, Tainai H, Nimura Y (2009) A randomized study of gemcitabine/cisplatin versus single-agent gemcitabine in patients with biliary tract cancer. *Proc Am Soc Clin Oncol* 27: abstract no: 4579
7. Gores GJ (2003) Cholangiocarcinoma: current concepts and insights. *Hepatology* 37:961–969
8. Jarnagin WR, Ruo L, Little SA, Klimstra D, D'Angelica M, DeMatteo RP, Wagman R, Blumgart LH, Fong Y (2003) Patterns of initial disease recurrence after resection of gallbladder carcinoma and hilar cholangiocarcinoma: implications for adjuvant therapeutic strategies. *Cancer* 98:1689–1700
9. Kiba T, Nishimura T, Matsumoto S, Hatano E, Mori A, Yasumi S, Doi R, Ikai I, Kitano T, Nishimura T, Yoshikawa K, Ishiguro H, Yanagihara K, Doi E, Teramukai S, Fukushima M (2006) Single-agent gemcitabine for biliary tract cancers. Study outcomes and systematic review of the literature. *Oncology* 70:358–365
10. Knox JJ, Hedley D, Oza A, Feld R, Siu LL, Chen E, Nematollahi M, Pond GR, Zhang J, Moore MJ (2005) Combining gemcitabine and capecitabine in patients with advanced biliary cancer: a phase II trial. *J Clin Oncol* 23:2332–2338
11. Koeberle D, Saletti P, Borner M, Gerber D, Dietrich D, Caspar CB, Mingrone W, Beretta K, Strasser F, Ruhstaller T, Mora O, Herrmann R (2008) Patient-reported outcomes of patients with advanced biliary tract cancers receiving gemcitabine plus capecitabine: a multicenter, phase II trial of the Swiss Group for Clinical Cancer Research. *J Clin Oncol* 26:3702–3708
12. Koizumi W, Akiya T, Sato A, Sakuyama T, Sasaki E, Tomidokoro T, Hamada T, Fujimori M, Kikuchi Y, Shimada K, Mine T, Yamaguchi K, Sasaki T, Kurihara M (2010) Phase II study of S-1 as first-line treatment for elderly patients over 75 years of age with advanced gastric cancer: the Tokyo Cooperative Oncology Group study. *Cancer Chemother Pharmacol* 65:1093–1099
13. Leonard GD, O'Reilly EM (2005) Biliary tract cancers: current concepts and controversies. *Expert Opin Pharmacother* 6:211–223
14. Morizane C, Okusaka T, Furuse J, Ishii H, Ueno H, Ikeda M, Nakachi K, Najima M, Ogura T, Suzuki E (2009) A phase II study of S-1 in gemcitabine-refractory metastatic pancreatic cancer. *Cancer Chemother Pharmacol* 63:313–319
15. Nakamura K, Yamaguchi T, Ishihara T, Sudo K, Kato H, Saisho H (2006) Phase II trial of oral S-1 combined with gemcitabine in metastatic pancreatic cancer. *Br J Cancer* 94:1575–1579
16. Oh DY, Cha Y, Choi IS, Yoon SY, Choi IK, Kim JH, Oh SC, Kim CD, Kim JS, Bang YJ, Kim YH (2010) A multicenter phase II study of gemcitabine and S-1 combination chemotherapy in patients with unresectable pancreatic cancer. *Cancer Chemother Pharmacol* 65:527–536
17. Okusaka T, Ishii H, Funakoshi A, Yamao K, Ohkawa S, Saito S, Saito H, Tsuyuguchi T (2006) Phase II study of single-agent gemcitabine in patients with advanced biliary tract cancer. *Cancer Chemother Pharmacol* 57:647–653
18. Penz M, Kornek GV, Raderer M, Ulrich-Pur H, Fiebigler W, Lenauer A, Depisch D, Krauss G, Schneeweiss B, Scheithauer W (2001) Phase II trial of two-weekly gemcitabine in patients with advanced biliary tract cancer. *Ann Oncol* 12:183–186
19. Sakuramoto S, Sasako M, Yamaguchi T, Kinoshita T, Fujii M, Nashimoto A, Furukawa H, Nakajima T, Ohashi Y, Imamura H, Higashino M, Yamamura Y, Kurita A, Arai K (2007) Adjuvant chemotherapy for gastric cancer with S-1, an oral fluoropyrimidine. *N Engl J Med* 357:1810–1820
20. Sasaki T, Isayama H, Nakai Y, Ito Y, Kogure H, Togawa O, Toda N, Yasuda I, Hasebe O, Maetani I, Sasahira N, Hirano K, Tsujino T, Tada M, Omata M (2010) Multicenter, phase II study of gemcitabine and S-1 combination chemotherapy in patients with advanced biliary tract cancer. *Cancer Chemother Pharmacol* 65:1101–1107
21. Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van Glabbeke M, van Oosterom AT, Christian MC, Gwyther SG (2000) New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 92:205–216
22. Trotti A, Colevas AD, Setser A, Rusch V, Jaques D, Budach V, Langer C, Murphy B, Cumberlin R, Coleman CN, Rubin P (2003) CTCAE v3.0: development of a comprehensive grading system for the adverse effects of cancer treatment. *Semin Radiat Oncol* 13:176–181
23. Tsavaris N, Kosmas C, Gouveris P, Gennatas K, Polyzos A, Mouratidou D, Tsipras H, Margaris H, Papastratis G, Tzima E, Papadoniou N, Karatzas G, Papalambros E (2004) Weekly gemcitabine for the treatment of biliary tract and gallbladder cancer. *Invest New Drugs* 22:193–198
24. Ueno H, Okusaka T, Ikeda M, Takezako Y, Morizane C (2004) Phase II study of S-1 in patients with advanced biliary tract cancer. *Br J Cancer* 91:1769–1774
25. Valle JW, Wasan H, Johnson P, Jones E, Dixon L, Swindell R, Baka S, Maraveyas A, Corrie P, Falk S, Gollins S, Locks F, Evans L, Meyer T, Anthoney A, Iveson T, Highley M, Osborne R, Bridgewater J (2009) Gemcitabine alone or in combination with cisplatin in patients with advanced or metastatic cholangiocarcinomas or other biliary tract tumours: a multicentre randomised phase II study—The UK ABC-01 Study. *Br J Cancer* 101:621–627
26. Valle J, Wasan H, Palmer DH, Cunningham D, Anthoney A, Maraveyas A, Madhusudan S, Iveson T, Hughes S, Pereira SP, Roughton M, Bridgewater J, ABC-02 Trial Investigators (2010) Cisplatin plus gemcitabine versus gemcitabine for biliary tract cancer. *N Engl J Med* 362:1273–1281
27. von Delius S, Lersch C, Schulte-Frohlinde E, Mayr M, Schmid RM, Eckel F (2005) Phase II trial of weekly 24-hour infusion of gemcitabine in patients with advanced gallbladder and biliary tract carcinoma. *BMC Cancer* 5:61
28. Yamauchi J, Kanai M, Matsumoto S, Nishimura T, Yazumi S, Kami K, Kawaguchi Y, Yasuda H, Kitano T, Misawa A, Ishiguro H, Yoshikawa K, Yanagihara K, Fukushima M, Doi R, Chiba T (2008) Clinical outcome of gemcitabine/S-1 combination therapy for advanced pancreatic cancer. *Pancreas* 36:327–328
29. Yanagihara K, Yoshimura K, Niimi M, Yasuda H, Sasaki T, Nishimura T, Ishiguro H, Matsumoto S, Kitano T, Kanai M, Misawa A, Tada H, Teramukai S, Mio T, Fukushima M (2010) Phase II study of S-1 and docetaxel for previously treated patients with locally advanced or metastatic non-small cell lung cancer. *Cancer Chemother Pharmacol* 66:913–918
30. Yoshizawa J, Takizawa A, Takeuchi O, Hiraku O, Sasaki K, Morimoto Y, Atsuda K, Inoue G, Suzuki Y, Asanuma F, Yamada Y (2009) Experimental study of combination therapy with S-1 against pancreatic cancer. *Cancer Chemother Pharmacol* 64:1211–1219