

new applications for starch although many of these opportunities remain to be explored. Areas of interest are packaging, textiles, controlled release, cosmetics, pharmaceuticals and flocculation.

Carbohydrate Chemistry Focused on Agrochemical Application

Amélia P. Rauter

Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Edifício C1, 5º Piso, Campo Grande, 1700 Lisboa, Portugal

In the last few years we have been devoted to the synthesis of biomolecules potentially interesting as agrochemicals. We now describe the preparation of the sugar moiety of Amipurimycin, of Miharamycin and of its analogues. These two molecules are antibiotics known to inhibit *Pyricularia oryzae*, responsible for the rice blast disease.

Pseudo-C-nucleosides are another group of compounds which have been investigated by us, in order to obtain new structures possessing bioactivity. In this work we also describe the synthesis and bioactivity of some thiazoles, tetrazoles, triazoles and pyrazoles, obtained by chain elongation of some sugar precursors.

The α,β -unsaturated- γ -lactone unit is known from the literature to confer a great diversity of biological effects. We present the synthesis of fungitoxic sugar molecules, containing this unit in their structure, via Reformatsky-type reaction of the appropriate carbonyl compounds with ethyl α -bromo-methyl acrylate and zinc. The relationship between structure, conformation, configuration of the molecules and the bioactivity detected will be evaluated.

Carbohydrate Liquid Crystals

George A. Jeffrey

Department of Crystallography, University of Pittsburgh, Pittsburgh, PA 15206, USA

Emil Fischer reported the *double melting points* of some long chain n-alkyl pyranosides, but failed to connect the observation with liquid crystal formation. It was not until nearly forty years later that these compounds were shown to form liquid crystals. This observation seemed to be overlooked, since some thirty years later none of the text books or reviews of the subject mentioned carbohydrates. The revival of interest came in 1984 when 17 examples were reported. Now a recent Liquid Crystal Data Base lists over 2000 Carbohydrates.

Liquid crystals fall into two classes; thermotropic and lyotropic. The thermotropic may be calamatic, discotic or chiral. The lyotropic phases, formed on contact with water, may be laminar, cubic or hexagonal.

Carbohydrates which illustrate these phases will be described. The potential commercial uses will be discussed.

Carbohydrate Polymers as Wound Management Aids

J.F. Kennedy*, P. Methacanon, L.L. Lloyd, M. Paterson and C.J. Knill

Birmingham Carbohydrate and Protein Technology Group, School of Chemistry, The University of Birmingham, Birmingham, B15 2TT, UK

Polymeric materials composed of carbohydrate units, i.e. polysaccharides, are a diverse group of biological macromolecules that are showing increasing application in areas of wound management. A variety of neutral polysaccharides, e.g. dextran and starch, basic polysaccharides, e.g. chitin and chitosan, acidic polysaccharides, e.g. alginic acid and hyaluronic acid, and glycosaminoglycans, e.g. dermatan sulphate and heparin, and their respective derivatives, have been the focus of much interest with respect to biomedical and particularly woundcare applications over recent years, however there has been no directive that any one chemical structure is more efficacious than any other. To be suitable as a wound management aid a dressing material should exhibit a number of properties including abilities to: maintain high humidity at the wound-dressing interface provide protection against secondary infection, be able to remove excess wound exudate and toxic components, and maintain its strength when sterile and wet.

More recent investigations have examined some of the more unusual polysaccharides isolated from plant, bacterial and animal sources which possess potentially useful biological properties that may make them suitable for woundcare applications, e.g. arabinoxylan and β -D-(1 \rightarrow 3)-glucan derivatives. The precise structures of such unusual polysaccharide based materials are often unknown and detailed compositional and structural characterisation is required in order to satisfy regulatory authorities. The carbohydrate structure of a novel polysaccharide based material, namely Sterigel[®] (Seton Healthcare), used as a wound management aid has been determined. Enzymic hydrolysis and methylation analysis have shown the carbohydrate structure to be a highly substituted β -D-(1 \rightarrow 4)-xylan. This polysaccharide backbone is substituted with α -L-arabinofuranoside residues and α -D-glucopyranosyluronic acid residues. The total amino acid content of the Sterigel, as determined after acid hydrolysis, is 4.4% w/w with the amino acid hydroxyproline accounting for 0.22%. The cinnamic acid derivative ferulic acid has been identified in both alkaline (0.40%) and enzymic (0.25%) hydrolysates of the polysaccharide.

Carbohydrate Substituted Porphyrins. Synthesis, Characterization and Lipoprotein Binding Properties

Hermann K. Hombrecher* and Christian Schell

Institut für Chemie der Medizinischen Universität zu Lübeck, Ratzeburger Allee 160, D-23538 Lübeck, Germany

The development of new photosensitisers for photodynamic therapy of tumours (PDT) is one of the most important fields of porphyrin chemistry today. PDT is a treatment that is based on the selectivity of porphyrinic compounds to tumour tissue and on the production of singlet oxygen by irradiation of the sensitiser with visible light. Thus formation of singlet oxygen in tumour cells causes cell death and tumour necrosis. Besides long wavelength absorption solubility and water and high selectivity to tumour cells are requirements a new photosensitiser has to fulfill. Therefore a number of different carbohydrate substituted porphyrins were synthesised. Although the mechanism of sensitiser uptake is not yet clarified, there is evidence that amphiphilic porphyrins associate to LDL and are introduced into the tumour cell via receptor mediated endocytosis.