

Short communication

Characterization and production of the exopolysaccharide (EPS) from
Pseudomonas aeruginosa G1 and *Pseudomonas putida* G12 strainsGokcen Yuvali Celik ^{a,*}, Belma Aslim ^b, Yavuz Beyatli ^b^a Nigde University, Faculty of Science and Arts, Department of Biology, Campus, 51200 Nigde, Turkey^b Gazi University, Faculty of Science and Arts, Department of Biology, 06500 Teknikokullar, Ankara, Turkey

Received 11 July 2007; received in revised form 19 October 2007; accepted 14 November 2007

Available online 22 November 2007

Abstract

In this study, exopolysaccharides (EPSs) production was investigated by growing strains of *Pseudomonas aeruginosa* G1 and *Pseudomonas putida* G12 in medium containing various carbon sources such as glucose, mannose, fructose, and xylose. EPS production (192 and 182 mg/L, respectively) of these strains grown in PAP medium with 2% glycerol (v/v) was used as control. The highest EPS production of the two strains was found in the xylose containing medium. The effect of different concentrations [2–6% (w/v)] of xylose on EPS production of both strains was also studied. The maximum EPS yield (368 mg/L⁻¹) of the strain G1 was recorded in 3% (w/v) xylose, while the highest yield EPS yield (262 mg/L⁻¹) of the strain G12 was recorded in 2% (w/v) xylose. The monosaccharide compositions of EPS produced by the two strains were analyzed by HPLC. Strain G1 [2% (w/v) glycerol] was found to compose of neutral sugars (92.0%), acetylated amino sugars (8.0%), while strain G1 [3% (w/v) xylose] contained neutral sugars (99.2%), acetylated amino sugars (0.8%). However, the composition of strain G12 [2% (w/v) glycerol] was neutral sugars (96.8%), acetylated amino sugars (3.2%) and the strain G12 [2% (w/v) xylose] was found to contain neutral sugars (96.1%), acetylated amino sugars (3.9%).

© 2007 Elsevier Ltd. All rights reserved.

Keywords: Exopolysaccharide (EPS); *P. aeruginosa*; *P. putida*; Different carbon sources

1. Introduction

Microbial exopolysaccharides (EPSs) are produced by various genera of bacteria and yeasts have found a wide range of applications in the food, pharmaceutical and petroleum industries and in medical fields and are widely accepted products of biotechnology (Lee et al., 1997; Sutherland, 1998). A wide range of chemical structures of homopolymeric or heteropolymeric type, made up of sugar and non-sugar components, is possible and the range of monosaccharide combinations, together with non-carbohydrate substituents and varied linkage types, makes the EPS an excellent agent and attributes diversity in bacteria (Keene & Lindberg, 1983).

Often bacterial EPS production is favored by temperatures lower than those optimal for growth, a high carbon to nitrogen ratio in the growth medium and nutrient deprivation (Fett, 1993). EPS characteristics and amounts can be influenced by several factors such as composition of the medium (carbon and nitrogen sources), as well as incubation conditions (temperature, pH, time, etc.) (De Vuyst & Degeest, 1999; Looisjesteijn, Boels, Kleerebezem, & Hugenholtz, 1999; Tallon, Bressollier, & Urdaci, 2003). That explains the diversity of the contradictory results on exopolysaccharide production. There is a considerable interest in finding new EPSs that are suitable for special applications, or that have potential industrial relevance, either by applying different culture conditions or by using novel bacterial strains (Looisjesteijn, Van Casteren, Tuinier, Doeswijk-Voragen, & Jhugenholtz, 2000).

Increasing attention is being paid to these molecules because of their bioactive role and their wide range of in the biotechnology and biopharmaceutical industries

* Corresponding author. Tel. +90 388 2252116; fax: +90 388 2250180.
E-mail addresses: gycelik@nigde.edu.tr, gokcenyc@hotmail.com (G.Y. Celik).

(Sutherland, 1990). Examples of industrially important microbial exopolysaccharides are dextrans, xanthan, gellan, pullulan, yeast glucans and bacterial alginates (De Vuyst & Degeest, 1999). Bacterial alginates were secreted by *Pseudomonas* species, *Azotobacter vinelandi* and *Azotobacter chroococcum*. Also, production of gellan by *Pseudomonas* species on a laboratory scale was reported by (Banik, Kanari, & Upadhyay, 2000).

In this work, we aimed to study the EPS synthesis by *Pseudomonas aeruginosa* and *Pseudomonas putida* strains during growth on various carbon sources and their different concentrations. We also aimed at describing the monosaccharide composition of EPS produced by these strains grown in both *Pseudomonas* Agar P (PAP) medium with concentrations of carbon source determined the highest EPS production and *Pseudomonas* Agar P medium with 2% (v/v) glycerol (control).

2. Experimental

2.1. Isolation and identification

Pseudomonas aeruginosa G1 and *P. putida* G12 strains used in this research were isolated from polluted soil in the Turkey. Initially, they were tested from Gram stains and catalase reactions, cell shape (Sneath, 1986). Then their carbohydrate fermentation characteristics were determined using Analytical Profile Index (API) 20 NE (Biomerieux, Marcy l'Etoile, France) identification system. Also, G1 and G12 strains identified by the 16S rRNA gene sequence analysis and the biochemically identification of these strains were confirmed by 16S rRNA gene sequence analysis. The sequences obtained were searched against The GenBank DNA database using the blast function (Widmer, Seidler, Gillevet, Watrud, & Di Giovanni, 1998). All isolates were stored on Nutrient Agar Medium (Difco) slopes at 4 °C and stock cultures were maintained at –20 °C in 0.5% (v/v) glycerol.

2.2. Culture conditions for EPS production

PAP medium with 2% (v/v) glycerol was used for the production of EPS. The glycerol in PAP medium was taken out to study the abilities EPS producing of both strains at different carbon sources. After then, glucose, mannose, fructose, and xylose were added into the PAP medium in amounts equivalent to the glycerol concentration. All media were autoclaved 15 min at 121 °C. Sugars were autoclaved separately.

To study the influence of xylose carbon source on EPS production by strains G1 and G12 were added to PAP medium at concentrations of 2–6% (w/v).

2.3. Isolation and quantification of EPS

Isolation of bacterial EPS was done as described (Cérantola, Bounéry, Segonds, Marty, & Montrozier,

2000) by Cérantola et al. (2000). Strains were grown on *Pseudomonas* Agar P (Difco) medium, supplemented with 2% (v/v) glycerol for 3 days at appropriate temperature (30 or 37 °C). Agar plate cultures were then washed with saline (0.9% NaCl w/v) using a glass rod and the resulting suspensions were stirred with glass beads in order to detach EPS associated with the bacterial cells. Cells were then removed by centrifugation at 10,000g for 30 min. The resulting supernatants were precipitated overnight at 4 °C with six volumes of 95% ethanol. Precipitated EPS were recovered by centrifugation and the ethanol precipitation step was repeated once again. After centrifugation (12,000g for 30 min at 4 °C), pellets were dissolved in distilled water. Total EPS (expressed as mg L⁻¹) was estimated in each sample by phenol-sulphuric method (Dubois, Gilles, Hamilton, Rebers, & Smith, 1956) using glucose as a standard (Torino, Taranto, Sesma, & de Valdez, 2001).

2.4. Determination of the EPS monomer composition

The isolated EPS was lyophilized. For the monomer analysis, 2 mg of dried EPS was dissolved in 2 ml of double distilled water. The solution was incubated at 100 °C in a water bath for 7 h. After being cooled and neutralized, the clear solution was collected and kept at –20 °C until analysis. Determination of monosaccharides was performed by HPLC (Varian ProStar) with Carbohydrates Ca (300 × 6.5 mm) CP28351 column using water as mobile phase at a flow rate of 0.5 mL min⁻¹ and the eluted were detected with a detector (VARIAN 350 RI).

Determination of (acetylated) amino sugars was performed by HPLC (Varian ProStar) with MetaCarb 87H (300X 7.8 mm) (5220) column using 0.0008 N H₂SO₄ as mobile phase at a flow rate of 0.5 mL min⁻¹ and the eluted were detected with a detector (VARIAN 350 RI). The analyses were accomplished by the METU – AR-GE (Ankara – TURKEY) and studied as two replications.

3. Results and discussion

3.1. Isolation and quantitative determination of EPS

Carbohydrate sugars, such as glucose, xylose, lactose, galactose and sucrose, are preferred carbon sources for the production of EPS. *P. aeruginosa* G1 and *P. putida* G12 strains were cultivated in the PAP medium containing various carbon sources (glucose, mannose, fructose and xylose) to find a suitable carbon source for the EPS production. It was used EPS production (192 and 182 mg/L, respectively) of these strains grown in PAP medium with 2% glycerol (v/v) as control. When strains G1 and G12 were grown in the xylose containing medium, the EPS productions (335, 262 mg L⁻¹, respectively) were the highest among those tested and their control. The EPS produced (267 mg L⁻¹) from fructose and other carbon sources by strain G11 was much lower than that from xylose. Simi-

larly, the EPSs produced (208, 196, 242 mg L⁻¹) from glucose, mannose, fructose by strain G12 were lower than that from xylose. The results were indicated in Table 1.

The composition of the medium plays an important role in the production of EPS (Drevet, Monot, Ballerni, Lecourtier, & Choplin, 1994). Manresa et al. reported that *Pseudomonas* sp. strain GSP-910 was able to utilize all the tested carbon source (glucose, sucrose, mannose, galactose, fructose, xylose, lactose and maltose) to synthesize the EPS however, glucose, sucrose and xylose allowed maximum EPS production (Manresa, Espuny, Guinea, & Comelles, 1987).

Also, the influence of different concentrations of xylose [2–6% (v/v)] on EPS production of both strains was studied. As shown in Table 2, maximum EPS productions (368 mg/L⁻¹) of strain G1 was recorded with 3% (w/v) xylose. On the other hand, the highest yield EPS production (262 mg/L⁻¹) by strain G12 was detected with 2% (w/v) xylose. Also, these EPS amounts (368 and 262 mg/L⁻¹) were higher than their controls. Several researches determined that EPS production by *Pseudomonas* sp. was enhanced in presence of suitable carbon sources in excess (Congregado et al., 1985). From our data, when strain G1 was cultivated in the PAP medium with 3% xylose, the production of EPS was stimulated. However, high concentrations of xylose inhibited EPS production of both strains.

3.2. Monosaccharide composition of EPS

The monosaccharide compositions of EPS produced by the two strains showing the highest and lowest EPS production were analyzed by HPLC. The results showed that the contents of neutral sugars and amino sugars of these strains. Glycerol dominated in the EPS produced by strain G1 and strain G12 in PAP medium with 2% (v/v) glycerol (92.0–88.4%). Besides glycerol, in PAP medium with 2% (v/v) glycerol by strain G1 was found to contain *N*-acetyl D-glucose amine (8.0%) while in PAP medium with 2%

Table 2

Influence of different concentrations in media containing xylose on EPS production

Strains	Concentration (%)	EPS (mg L ⁻¹)
<i>P. aeruginosa</i> G1	2	335 ± 1
	3	368 ± 0 ^a
	4	204 ± 2
	5	124 ± 3
	6	110 ± 1
<i>P. putida</i> G12	2	262 ± 0 ^a
	3	208 ± 1
	4	189 ± 0
	5	172 ± 2
	6	97 ± 1

^a The highest EPS production.

(v/v) glycerol by strain G12 was found to contain mannose (8.4%) and *N*-acetyl D-glucose amine (3.2%). The polysaccharides produced by strain G1 in PAP medium with 2% (w/v) xylose, was composed of mannose (88.0%), mannitol (11.2%) and *N*-acetyl D-glucose amine (0.8%). Also, The polysaccharides produced by strain G12 in PAP medium with 3% (w/v) xylose, was composed of galactose (78.7%), glycerol, (11.8%), glucose (3.9%), *N*-acetyl D-glucose amine (3.9%) and ribose (1.7%) (Table 3).

Most EPS are composed of the neutral sugars glucose, galactose and/or rhamnose (Faber, Zoon, Kamerling, & Vliegthart, 1998; Lemoine et al., 1997; Yamamoto, Nunome, Yamauchi, Kato, & Sone, 1995). Some of them also consist of (acetylated) amino sugars (Robijn et al., 1996; Stinglee, Neeser, & Mollet, 1996). The strain, the culture conditions, and the type of carbon source influence the amount and the composition of microbial EPS that is produced by a certain species (Looisjesteijn et al., 1999). With glucose as a carbon source, *P. putida* and *Pseudomonas fluorescens* synthesized an EPS composed of glucose, galactose and pyruvate (Read & Costerton, 1987). Hung et al. found that EPS produced by *P. fluorescens* Biovar II was composed of galactose, mannose, rhamnose, glucose, fucose, ribose, arabinose and xylose (Hung, Santschi, & Gillow, 2005). Kachlany et al. revealed that the purified EPS from *P. putida* G7 contained the monosaccharides, glucose, rhamnose, ribose, *N*-acetylgalactosamine and glucuronic acid (Kachlany et al., 2001). Similarly, some researches showed that EPS were isolated from *Pseudomonas caryophylli* CFR 1705 grown on lactose containing medium was composed of rhamnose, mannose and glucose in the ratio 1:3.26:4.97, respectively (Sudhamani, Tharanathan, & Prasad, 2004). Also, EPS production and its monomeric sugar composition were dependent on the carbon source, nitrogen source and carbon/nitrogen ratio of the growth medium (Cerning et al., 1994). Therefore, the results reported here demonstrate that our strains likely produces as dependent on the carbon source similar common polysaccharides as other *Pseudomonas*.

In conclusion, the present study describes the influence of various carbon sources and different concentrations of

Table 1
Influence of different carbon sources on EPS production

Strains	Carbon source (2%)	EPS (mg L ⁻¹) ^c
<i>P. aeruginosa</i> G1	Glucose	225 ± 0
	Mannose	262 ± 3
	Fructose	267 ± 1
	Xylose	335 ± 1 ^b
	Control ^a	192 ± 5
<i>P. putida</i> G12	Glucose	208 ± 0
	Mannose	196 ± 1
	Fructose	242 ± 0
	Xylose	262 ± 0 ^b
	Control ^a	182 ± 3

^a It was used *Pseudomonas* Agar P medium with 2% glycerol (v/v) as control.

^b The highest EPS production.

^c Values are means ± standard deviations of triplicate measurements.

Table 3
Carbohydrate compositions of EPSs produced by two strains

Strains	Medium	Monosaccharide (%)									
		Glucose	Xylose	Galactose	Rhamnose	Mannose	Arabinose	Mannitole	Glycerol	Ribose	Galactronic asit
<i>P. aeruginosa</i> G1	PAP with 2% glycerol	–	–	–	–	–	–	–	92.0	–	–
<i>P. aeruginosa</i> G1	PAP with 3% xylose	–	–	–	–	88.0	–	11.2	–	–	–
<i>P. putida</i> G12	PAP with 2% glycerol	–	–	–	–	8.4	–	–	88.4	–	–
<i>P. putida</i> G12	PAP with 2% xylose	3.9	–	78.7	–	–	–	–	11.8	1.7	–

Not determined.

carbon source on EPS production and monomeric sugar composition of two *Pseudomonas* strains isolated from waste soil in Turkey.

References

- Banik, R. M., Kanari, B., & Upadhyay, S. N. (2000). Exopolysaccharide of the gellan family: Prospects and potential. *World Journal of Microbiology*, 16, 407–414.
- Cérantola, S., Bounéry, J. D., Segonds, C., Marty, N., & Montrozier, H. (2000). Exopolysaccharide production by mucoid and non-mucoid strains of *Burkholderia cepacia*. *FEMS Microbiology Letters*, 185, 243–246.
- Cerning, J., Renard, C. M. G. C., Thibault, J. F., Bouillanne, C., Landon, M., Desmazeaud, M., et al. (1994). Carbon source requirements for exopolysaccharide production by *Lactobacillus casei* CG11 and partial structure analysis of the polymer. *Journal of Applied Microbiology*, 60, 3914–3919.
- Congregado, F., Estañol, I., Espuny, M. J., Fust, M. C., Manresa, M. A., Marqués, A. M., et al. (1985). Preliminary studies on the production and composition of the extracellular polysaccharide synthesized by *Pseudomonas* sp EPS-5028. *Biotechnology Letters*, 7, 883–888.
- De Vuyst, L., & Degeest, B. (1999). Heteropolysaccharides from lactic acid bacteria. *FEMS Microbiology Reviews*, 23, 153–177.
- Drevet, E., Monot, F., Ballerni, D., Lecourtier, J., & Choplin, L. (1994). Effect of mixing and mass transfer conditions on gellan production by *Pelodea*. *Journal of Fermentation and Bioengineering*, 77, 642–649.
- Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A., & Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, 28, 350–356.
- Faber, E. J., Zoon, P., Kamerling, J. P., & Vliegthart, J. F. G. (1998). The exopolysaccharides produced by *Streptococcus thermophilus* Rs and Sts have the same repeating unit but differ in viscosity of their milk cultures. *Carbohydrate Research*, 310, 269–276.
- Fett, W. F. (1993). Bacterial exopolysaccharides: Their nature, regulation and role in host–pathogen interactions. *Current Topics in Botanical Research*, 1, 367–390.
- Hung, C.-C., Santschi, P. H., & Gillow, J. B. (2005). Isolation and characterization of extracellular polysaccharides produced by *Pseudomonas fluorescens* Biovar II. *Carbohydrate Polymers*, 61, 141–147.
- Kachlany, S. C., Levery, S. B., Kim, J. S., Reuhs, B. L., Lion, L. W., & Ghiorse, W. C. (2001). Structure and carbohydrate analysis of the exopolysaccharide capsule of *Pseudomonas putida* G7. *Environmental Microbiology*, 3, 774–784.
- Keene, L., & Lindberg, B. (1983). Bacterial polysaccharide. In G. O. Aspinall (Ed.), *In the polysaccharides* (Vol. 2, pp. 287–363). London: Academic Press.
- Lee, I. Y., Seo, W. T., Kim, G. J., Kim, M. K., Ahn, S. G., & Kwon, G. S. (1997). Optimization of fermentation conditions for production of exopolysaccharide by *Bacillus polymyxa*. *Bioprocess and Biosystems Engineering*, 16, 71–75.
- Lemoine, J., Chirat, F., Wieruszkeski, J.-M., Strecker, G., Favre, N., & Neeser, J.-N. (1997). Structural characterization of the exocellular polysaccharides produced by *Streptococcus thermophilus* SFi39 and SFi12. *Applied and Environmental Microbiology*, 63, 3512–3518.
- Looisjesteijn, P. J., Boels, I. C., Kleerebezem, M., & Hugenholtz, J. (1999). Regulation of exopolysaccharide production by *Lactococcus lactis* subsp. cremoris by the sugar source. *Applied and Environmental Microbiology*, 65, 5003–5008.
- Looisjesteijn, P. J., Van Casteren, W. H. M., Tuinier, R., Doeswijk-Voragen, C. H. L., & Hugenholtz, J. (2000). Influence of different substrate limitations on the yield, composition and molecular mass of exopolysaccharides produced by *Lactococcus lactis* subsp. cremoris in continuous cultures. *Journal of Applied Microbiology*, 89, 116–122.
- Manresa, A., Espuny, M. J., Guinea, J., & Comelles, F. (1987). Characterization and production of a new extracellular polymer from

- Pseudomonas* sp. GSP-910. *Applied Microbiology and Biotechnology*, 26, 347–351.
- Read, R. R., & Costerton, J. W. (1987). Purification and characterisation of adhesive exopolysaccharides from *Pseudomonas putida* and *Pseudomonas fluorescens*. *Canadian Journal of Microbiology*, 33, 1080–1090.
- Robijn, G. W., Wienk, H. L. J., van den Berg, D. J. C., Haas, H., Kamerling, J. P., & Vliegthart, J. F. G. (1996). Structural studies of the exopolysaccharide produced by *Lactobacillus paracasei* 34-1. *Carbohydrate Research*, 285, 129–139.
- Sneath, P. H. A. (1986). Bergey's Manual of Systematic bacteriology, Sneath, P. H. A., Mair, N. S., Sharp, M. E., Holt, J. G. (Eds.), Williams and Wilkins, Vol. 2, 1 (pp. 141–199). Baltimore.
- Stingle, F. J.-R., Neeser, J.-R., & Mollet, B. (1996). Identification and characterization of the eps (exopolysaccharide) gene cluster from *Streptococcus thermophilus* Sfi6. *Journal of Bacteriology*, 178, 1680–1690.
- Sudhamani, S. R., Tharanathan, R. N., & Prasad, M. S. (2004). Isolation and characterization of an extracellular polysaccharide from *Pseudomonas caryophylli* CFR 1705. *Carbohydrate Polymers*, 56, 423–427.
- Sutherland, I. W. (1990). *Biotechnology of microbial exopolysaccharides*. Cambridge: Cambridge University Press.
- Sutherland, I. W. (1998). Novel and established applications of microbial polysaccharides. *Trends in Biotechnology*, 16, 41–46.
- Tallon, R., Bressollier, P., & Urdaci, M. C. (2003). Isolation and characterization of two exopolysaccharides produced by *Lactobacillus plantarum* EP56. *Research Microbiology*, 154, 705–712.
- Torino, M. I., Taranto, M. P., Sesma, F., & de Valdez, G. F. (2001). Heterofermentative pattern and exopolysaccharide production by *Lactobacillus helveticus* ATCC 15807 in response to environmental pH. *Journal of Applied Microbiology*, 91, 846–852.
- Widmer, F., Seidler, R. J., Gillevet, P. M., Watrud, L. S., & Di Giovanni, G. D. (1998). A highly selective PCR Protocol for detecting 16 SrRNA genes of the genus *Pseudomonas* (Sensu Stricto) in environmental samples. *Applied and Environmental Microbiology*, 64, 2545–2553.
- Yamamoto, Y., Nunome, T., Yamauchi, R., Kato, K., & Sone, Y. (1995). Structure of an exocellular polysaccharide of *Lactobacillus helveticus* TN-4, a spontaneous mutant strain of *Lactobacillus helveticus* TY1-2. *Carbohydrate Research*, 275, 319–332.