SUBJECT INDEX

- 2-Acetamido-6-*O*-(5-acetamido-3,5-dideoxy-β-Dglycero-D-galacto-2-nonulopyranosylonic acid)-2-deoxy-D-glucose, synthesis of, 19
- 2-Acetamido-2-deoxy-α-D-glucopyranoside,preparation of the 3,4-di-O-benzyl and 3,4-di-O-(4-phenylbenzyl) derivatives of benzyl, 168
- Acetates, n.m.r. (1H and 13C) studies of some carbohydrate, 266
- 6-O-(N-Acetyl-β-D-neuraminyl)-2-deoxy-D-glucose, synthesis of 2-acetamido-, 19
- Aldonolactones, conformational studies on, by n.m.r. spectroscopy, 111, 131
- 2-Amino-2,4-dideoxy-DL-pentopyranosides, methyl, synthesis and conformation of, 9
- 5-Aminofluorescein, labeling of glycosaminoglycuronans with, 69
- 3-Amino-2,3,6-trideoxy-3-C-methyl-L-xylo-hexopyranose, synthesis of derivatives of, c1
- Antigen from Escherichia coli LP 1092, structure of the 3-deoxy-D-manno-octulosonic acid-containing polysaccharide K6, 45
- Beef-lung heparan sulphates with specific selfassociations, structural differences between, 215
- Benzyl 2-acetamido-2-deoxy-α-D-glucopyranoside, preparation of the 3,4-di-O-benzyl and 3,4-di-O-(4-phenylbenzyl) derivatives of, 168
- Calorimetric determination of the conformational transition of kappa carrageenan, 227
- Capsular polysaccharide from *Klebsiella* serotype K54; location of the *O*-acyl groups, and a revised structure, 189
- Carbohydrate acetates, n.m.r. (¹H and ¹³C) studies of some, 266
- Carrageenan, calorimetric determination of the conformational transition of kappa, 227
- Cell recognition, and cell-surface interactions, the phytochemistry of (book review), c11
- Cellulose under water-limiting conditions, the role of water in the hydrolysis of, 288
- Chemical features of aggregatable heparan sulphates from human-lung fibroblasts, 205
- Conformational studies on aldonolactones by n.m.r. spectroscopy. Conformations of p-glucono-, p-manno-, p-gulono-, and p-galactono-1,4-lactone in solution, 131

- Conformational studies on aldonolactones by n.m.r. spectroscopy. Conformations of p-pentono-1,4-lactones in solution, 111
- Conformational transition of kappa carrageenan, calorimetric determination of the, 227
- Conformations of the p-glucarolactones and pglucaric acid in solution, 95
- Contribution to the sequential analysis of oligosaccharides by mass spectrometry, 251
- Cyclodextrin-nitroxide-radical complexes, determination by ¹H-n.m.r. of, 154
- Cytotoxic activity of (triazeno)dextrans, 57
- Deoxy sugars, formation from halogenated carbohydrates of, by α -hydroxyalkyl radicals, 276
- Dextrans, triazeno-, synthesis of, 57
- Fluorescent glycosaminoglycuronans obtained by labeling with 5-aminofluorescein, 69
- 1,6:2,3-Dianhydro-4-*O*-benzyl-β-D-mannopyranose, side-product formation during the hydrogenolysis of, 269
- 2,6-Dideoxy-4-O-methyl-L-ribo-hexose, a facile synthesis of derivatives of, 165
- Formation of deoxy sugars from halogenated carbohydrates by α-hydroxyalkyl radicals, optimisation of the reaction conditions for the, 276
- Formation of superoxide, and in vitro inactivation of viruses, by hexopyranosid-3-uloses, c5
- p-Galacto-p-mannans, from yeasts, ¹³C-n.m.r. spectra of, 283
- Glomerulonephritis glycopeptide, synthesis of compounds related to, 271
- p-Glucaric acid, conformations of, in solution, 95
- D-Glucarolactones, conformations of, in solution, 95
- D-Glucitol, indirect anodic oxidation of, in aqueous calcium iodide, 158
- α-D-Glucopyranosylamine, N-(L-glutam-5-oyl)and α-D-Glcp-(1 \rightarrow 6)-N-(L-glutam-5-oyl)-, synthesis of, 271
- D-[1-14C]Glucosamine, low-molecular-weight metabolites in cultured, normal, and carcinomatous, human-mammary cells, 292

c16 Subject index

- N-L-(Glutam-5-oyl)-x-D-glucopyranosylamine, synthesis of, 271
- N-L-(Glutam-5-oyl)- α -D-glucopyranosylamine, α -D-Glcp-(1 \rightarrow 6)- β -D-Glcp-(1 \rightarrow 6)-, synthesis of, 271
- Glycosaminoglycuronans, fluorescent, labeled with 5-aminofluorescein, 69
- Halogenated carbohydrates, formation of deoxy sugars from, by α-hydroxyalkyl radicals, 276 Hemicellulose, obtained from the fiber of
- Hemicellulose, obtained from the fiber of Sansevieria trifasciata leaves, investigation on the structure of the, 247
- Heparan sulphates from human-lung fibroblasts, chemical features of aggregatable, 205
- Heparan sulphates with specific self-associations, structural differences between beef-lung, 215
- Hexopyranosid-3-uloses, formation of superoxide by, and *in vitro* inactivation of viruses by,
- Human-lung fibroblasts, chemical features of aggregatable heparan sulphates from, 205
- Human-mammary cells, normal and carcinomatous, low-molecular weight metabolites of p-[1-14C]glucosamine in cultured, 292
- Hydrogenolysis of 1,6:2,3-dianhydro-4-O-benzyl-β-D-mannopyranose, side-product formation during the, 269
- Hydrolysis of cellulose under water-limiting conditions, the role of water in the, 288
- Indirect, anodic oxidation of p-glucitol in aqueous calcium iodide, 158
- Investigation on the structure of the hemicellulose obtained from the fiber of Sansevieria trifasciata leaves, 247
- Isolation of bronchial mucins from cystic fibrosis sputum by use of citraconic anhydride, 87
- Kijanimicin, a facile synthesis of derivatives of the 2,6-dideoxy-4-O-methyl-L-ribo-hexose component of, 165
- 1,4-Lactones (D-glucono-, D-mannono-, D-gulono-, and D-galactono-), conformations of, in solution, 131
- (1→4)-Linked α-L-rhamnans, synthesis of, 173
 D-Lyxose and D-xylose, synthesis of 2-C-methyl derivatives of, 181
- D-Mannans, -D-galacto-, ¹³C-n.m.r. spectra of yeast, 283
- cis- and 1,2-trans-p-Mannopyranosides, stereoselective syntheses of 1,2-, 33
- Mass spectrometry, a contribution to the sequential analysis of oligosaccharides by, 251

- Metabolites of D-[i-14C]glucosamine, low-molecular-weight, in cultures of normal and carcinomatous, human-mammary cells, 292
- Methyl 2-amino-2,4-dideoxy-DL-pentopyranose, synthesis and conformation of, 9
- 2-C-Methyl-D-lyxose and 2-C-methyl-D-xylose, synthesis of, 181
- Methyl O-methyl-α-L-rhamnopyranosides in their complexes with tin(II) chloride, n.m.r. spectra of, 258
- 3-C-Methyl-L-xylo-hexopyranose, synthesis of derivatives of 3-amino-2,3,6-trideoxy-, c1
- Mucins, bronchial, from cystic fibrosis sputum, isolation of, by use of citraconic anhydride, 87
- Nitroxide radical-cyclodextrin, determination by ¹H-n.m.r. of complexes of, 154
- ¹H-N.m.r. determination of complexes of cyclodextrin with nitroxide radicals, 154
- ¹³C-N.m.r. spectra of D-galacto-D-mannans from yeasts, 283
- N.m.r. spectra of methyl O-methyl-α-L-rhamnopyranosides in their complexes with tin(II) chloride, 258
- N.m.r. (¹H and ¹³C) studies of some carbohydrate acetates, 266
- Oligosaccharides, a contribution to the sequential analysis of, by mass spectrometry, 251
- Oxidation of p-glucitol in aqueous calcium iodide, indirect anodic, 158
- D-Pentono-1,4-lactones, conformation of, in solution, 111
- DL-Pentopyranosides, methyl 2-aminc-2,4-dideoxy-, synthesis and conformation of, 9
- Phytochemistry of cell recognition and cell surface interactions (book review), c11
- Polysaccharide, alkali-soluble, from the fruit body of *Ganoderma japonicum* Lloyd, structure of a, 237
- Polysaccharide, capsular, from *Klebsiella* serotype K54, location of the *O*-acyl groups, and revised structure, 189
- Polysaccharide (K6 antigen) from Escherichia coli LP 1092, structure of the 3-deoxy-p-manno-octulosonic acid-containing, 45
- Preparation of the 3,4-di-O-benzyl and 3,4-di-O-(4-phenylbenzyl) derivatives of benzyl 2-acetamido-2-deoxy-α-D-glucopyranoside, 168
- α-L-Rhamnans, synthesis of (1→4)-linked, 173
 Role of water in the hydrolysis of cellulose under water-limiting conditions, 288
- Separation, by g.l.c., of enantiomeric sugars as diastereoisomeric dithioacetals, I

SUBJECT INDEX c17

- Side-product formation during the hydrogenolysis of 1,6:2,3-dianhydro-4-*O*-benzyl-β-D-mannopyranose, 269
- Stereoselective syntheses of 1,2-cis- and 1,2-trans-p-mannopyranosides, 33
- Structural differences between beef-lung heparan sulphates with specific self-associations, 215
- Structure of an alkali-soluble polysaccharide from the fruit body of *Ganoderma japonicum* Lloyd, 237
- Synthesis of derivatives of 2,6-dideoxy-4-O-methyl-L-ribo-hexose (a component of kijanimicin), a facile, 165
- Structure of the 3-deoxy-D-manno-octulosonic acid-containing polysaccharides (K6 antigen) from Escherichia coli LP 1092, 45
- Sugars, enantiomeric, separation of, by g.l.c. of the diastereoisomeric dithioacetals, 1
- Sugars, tautomeric equilibria of some, by partially relaxed, ¹³C pulse Fourier-transform,

- nuclear magnetic resonance spectroscopy, 145 Synthesis of 2-acetamido-6-O-(5-acetamido-3,5-dideoxy-β-D-glycero-D-galacto-2-nonulopyranosylonic acid)-2-deoxy-D-glucose, 19
- Synthesis of derivatives of 3-amino-2,3,6-trideoxy-3-C-methyl-L-xylo-hexopyranose, c1 Synthesis of (1 \rightarrow 4)-linked α -L-rhamnans, 173
- Synthesis of 2-C-methyl-D-lyxose and 2-C-methyl-D-xylose, 181
- Tautomeric equilibria of some sugars by partially relaxed, ¹³C pulse Fourier-transform, nuclear magnetic resonance spectroscopy, 145
- Tin(II) chloride, n.m.r. spectra of methyl Omethyl-α-L-rhamnopyranosides in their complexes with, 258
- (Triazeno)dextrans, synthesis of, 57
- Yeast D-galacto-D-mannans, ¹³C-n.m.r. spectra of, 283