

SUBJECT INDEX

- 2-*O*-(2-Acetamido-2-deoxy- β -D-glucopyranosyl)-5a-carba- α -D-mannopyranose and 5a-carba-2-*O*- and 3,6-di-*O*-(α -D-mannopyranosyl)- α -D-mannopyranoses, synthesis of, 117
- 2-*C*-Acetamidomethyl-2-deoxy- α -D-glucopyranoside and its *manno* isomer, synthesis of methyl, 161
- 3-(1-Acetamidopropyl)pyrazolo[3,4-*b*]quinoxaline (flavazole) derivatives, preparation of, 89
- Allofuranose derivatives, NMR spectroscopy and conformational analysis of substituted 1,2:5,6-di-*O*-isopropylidene- α -D-, 245
- Amadori compounds, influence of malondialdehyde on the Maillard degradation of, 307
- Anionic surfactant for biomedical use, synthesis of D-glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates] as a new type of, 323
- Anticoagulant and antithrombin activities of oversulfated fucans, 355
- Binding studies on immunodeterminants: synthesis of β -(1 \rightarrow 6)-linked oligosaccharide methyl glycosides having 1–4 internal D-galactopyranosyl residues flanked by gentiobiose residues, 271
- Building blocks for carba-oligosaccharides related to cell-surface glycans, preparation of, 173
- 5a-Carba-2-*O*- and 3,6-di-*O*-(α -D-mannopyranosyl)- α -D-mannopyranoses, synthesis of 2-*O*-(2-acetamido-2-deoxy- β -D-glucopyranosyl)-5a-carba- α -D-mannopyranose and, 117
- Carba-oligosaccharides related to cell-surface glycans, preparation of building blocks for, 173
- Carbon–proton coupling constants for the C–S–C–H pathway in 1-thioglycosides, Karplus-type equation for vicinal, 225
- Chiral polysubstituted *cis*-fused decalin containing an angular *C*-methyl group, synthesis by an intramolecular Diels–Alder reaction, 167
- Chlamydia* lipopolysaccharides, an NMR spectroscopic investigation of Kdo-containing oligosaccharides related to the genus-specific epitope of, 213
- Conformational analysis of substituted 1,2:5,6-di-*O*-isopropylidene- α -D-allofuranose derivatives, NMR spectroscopy and, 245
- Conformational study of α -L-Rhap-(1 \rightarrow 2)- α -L-Rhap-(1 \rightarrow OMe) by NMR nuclear Overhauser effect spectroscopy (NOESY) and molecular dynamics calculations, 195
- p*-Coumaroylated and feruloylated methyl glycosides, synthesis of, 183
- Crystal and molecular structures of the hexitol hexa-acetates, 17
- Crystal structure of α -D-glucopyranosyl potassium hydrogenphosphate, 33
- Crystal structure of konjac glucomannan in the mannan II polymorphic form, molecular and, 41
- Crystal structure of the inclusion complex of cyclomaltoheptaose (β -cyclodextrin) with 4-*tert*-butyltoluene, 1
- Crystal structure of the regenerated form of (1 \rightarrow 3)- α -D-mannan, molecular and, 57
- Cyclomaltoheptaose (β -cyclodextrin), crystal structure of the inclusion complex of 4-*tert*-butyltoluene with, 1
- Decalin containing an angular *C*-methyl group, synthesis of a chiral polysubstituted *cis*-fused, by an intramolecular Diels–Alder reaction, 167
- Deoxyfluorocellulose acetate, synthesis and characterization of highly substituted, 131
- 1-Deoxynojirimycin-containing glycans related to the Lewis X and sialyl-Lewis X epitopes recognized by LEC-CAMs, synthesis of, c1
- Diels–Alder reaction, synthesis of a chiral polysubstituted *cis*-fused decalin containing an angular *C*-methyl group by an intramolecular, 167

- 2,6-Dioxabicyclo[3.1.1]heptanes, substituted: synthesis and conformational analysis of 1,3-anhydro-2,4-di-*O*-benzyl-6-deoxy- β -D-glucopyranose, 75
- Disaccharides in aqueous sodium hydroxide, epimerization of reducing terminal groups of (1 \rightarrow 2)-linked D-gluco- and D-manno-, 369
- Epimerization of reducing terminal groups of (1 \rightarrow 2)-linked D-gluco- and D-manno-disaccharides in aqueous sodium hydroxide, 369
- Feruloylated and *p*-coumaroylated methyl glycosides, synthesis of, 183
- Fucans, anticoagulant and antithrombin activities of oversulfated, 355
- D-Galactopyranosyl residues flanked by gentiobiose residues, synthesis of β -(1 \rightarrow 6)-linked oligosaccharide methyl glycosides having 1–4 internal, 271
- Garlic (*Allium sativum*) bulbs, new mannose-specific lectin from, 347
- β -D-Glucopyranose, 1,3-anhydro-2,4-di-*O*-benzyl-6-deoxy, synthesis, and conformational analysis by ¹H-NMR spectroscopy and molecular mechanics calculation, 75
- α -D-Glucopyranosyl potassium hydrogenphosphate, the crystal structure of, 33
- D-Glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates], synthesis of a new type of anionic surfactant for biomedical use, 323
- β -D-Glycobiosides, 1-thio-, 4-nitrophenyl, phase transfer catalyzed synthesis of, 149
- Glycosides, methyl, synthesis of feruloylated and *p*-coumaroylated, 183
- Hexitol hexa-acetates, the crystal and molecular structures of the, 17
- Inclusion complex of cyclomaltoheptaose (β -cyclodextrin) with 4-*tert*-butyltoluene, crystal structure of the, 1
- Isopropylidene and benzylidene derivatives of sugars: transacetalation, a convenient non-aqueous method for effecting the deprotection of, 141
- Karplus-type equation for vicinal carbon–proton coupling constants for the C–S–C–H pathway in 1-thioglycosides, 225
- Kdo-containing oligosaccharides related to the genus-specific epitope of *Chlamydia* lipopolysaccharides, an NMR spectroscopic investigation of, 213
- Konjac glucomannan in the mannan II polymorphic form, molecular and crystal structure of, 41
- Lectins from bulbs, mannose-specific, interaction with linear manno-oligosaccharides, 337
- Lewis X and sialyl-Lewis X epitopes, synthesis of related 1-deoxynojirimycin-containing glycans, C1
- Linked D-gluco- and D-manno-disaccharides in aqueous sodium hydroxide, epimerization of reducing terminal groups of (1 \rightarrow 2)-, 369
- Maillard degradation of Amadori compounds, influence of malondialdehyde on the, 307
- Malondialdehyde, influence on the Maillard degradation of Amadori compounds, 307
- (1 \rightarrow 3)- α -D-Mannan, molecular and crystal structure of the regenerated form of, 57
- Mannose-specific bulb lectins, interaction with linear manno-oligosaccharides and comparison with mannose/glucose-binding lectins, 337
- Mannose-specific lectins from garlic and ramsons bulbs, 347
- Methyl 3-*O*-(2-acetamido-2-deoxy-6-thio- β -D-glucopyranosyl)- β -D-galactopyranoside: a slow reacting acceptor-analogue which inhibits glycosylation by UDP-D-galactose-*N*-acetyl-D-glucosamine-(1 \rightarrow 4)-D-galactosyl-transferase, c5
- Methyl 2-C-acetamidomethyl-2-deoxy- α -D-glucopyranoside and its *manno* isomer, synthesis of, 161
- Methyl 3-deoxy-3-C-formyl- α -D-arabino-pentofuranoside, synthesis of derivatives of, 293
- Molecular and crystal structure of konjac glucomannan in the mannan II polymorphic form, 41
- Molecular and crystal structures of the hexitol hexa-acetates, 17
- Molecular and crystal structure of the regenerated form of (1 \rightarrow 3)- α -D-mannan, 57
- Neocarrabiose oligosaccharides having different length and sulphate substitution, use as model for ¹H-NMR spectroscopy, 233
- 4-Nitrophenyl 1-thio- β -D-glycobiosides, phase transfer synthesis of, 149
- NMR NOESY spectroscopy and molecular dynamics calculations, conformational study of α -L-Rhap-(1 \rightarrow 2)- α -L-Rhap-(1 \rightarrow OMe) by, 195
- NMR spectroscopic investigation of Kdo-containing oligosaccharides related to the genus-specific epitope of *Chlamydia* lipopolysaccharides, 213

- NMR spectroscopy and conformational analysis of substituted 1,2:5,6-di-*O*-isopropylidene- α -D-allofuranose derivatives, 245
- NMR spectroscopy, the use of neocarribose oligosaccharides having different length and sulphate substitution as model compounds for ^1H -, 233
- Oligosaccharide fragments of *Shigella dysenteriae* type 1 O-antigen, stereoselective syntheses of, 103
- Oligosaccharide methyl glycosides having 1–4 internal D-galactopyranosyl residues flanked by gentiobiose residues, synthesis of β -(1 \rightarrow 6)-linked, for immunodeterminant binding studies, 271
- Oligosaccharides having different length and sulphate substitution as model compounds for ^1H -NMR spectroscopy, the use of neocarribose, 233
- Oversulfated fucans, anticoagulant and antithrombin activities of, 355
- Pentasaccharide fragment of the capsular polysaccharide of *Streptococcus pneumoniae* type 18C, synthesis of a repeating, 155
- Pentofuranoside, synthesis of derivatives of methyl 3-deoxy-3-C-formyl- α -D-arabino-, 293
- Phenylhydrazone derivatives, addition and cyclisation reactions of sugar, via a 1,4-elimination process, 89
- Polysaccharide of *Streptococcus pneumoniae* type 18C, synthesis of a repeating pentasaccharide fragment of the capsular, 155
- Potassium hydrogenfluoride, reaction of 2-deoxy-6-*O*-[2,3-dideoxy-4,6-*O*-isopropylidene-2,3-(*N*-tosylepimino)- α -D-mannopyranosyl]-4,5-*O*-isopropylidene-1,3-di-*N*-tosylstreptamine with, 363
- Preparation of building blocks for carba-oligosaccharides related to cell-surface glycans, 173
- Quinoxalinone derivatives, preparation of 3-(2-acetamido-1-phenylhydrazonobutyl)-2(1*H*)-, 89
- Ramsons (*Allium ursinum*) bulbs, new mannose-specific lectin from, 347
- Reaction of 2-deoxy-6-*O*-[2,3-dideoxy-4,6-*O*-isopropylidene-2,3-(*N*-tosylepimino)- α -D-mannopyranosyl]-4,5-*O*-isopropylidene-2,3-di-*N*-tosylstreptamine with potassium hydrogenfluoride, 363
- α -L-Rha *p*-(1 \rightarrow 2)- α -L-Rha *p*-(1 \rightarrow OMe), conformational study by NMR NOESY spectroscopy and molecular dynamics calculations, 195
- Shigella dysenteriae* type 1 O-antigen, stereoselective syntheses of a di-, tri-, and tetra-saccharide fragment of, 103
- Sialyl-Lewis X and Lewis X epitopes, synthesis of related 1-deoxynojirimycin-containing glycans, C1
- Streptococcus pneumoniae* type 18C, synthesis of a repeating pentasaccharide fragment of the capsular polysaccharide of, 155
- Surfactant for biomedical use, synthesis of D-glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates] as a new type of anionic, 323
- Synthesis and characterization of highly substituted deoxyfluorocellulose acetate, 131
- Synthesis of 2-*O*-(2-acetamido-2-deoxy- β -D-glucopyranosyl)-5a-carba- α -D-mannopyranose and 5a-carba-2-*O*- and 3,6-di-*O*-(α -D-mannopyranosyl)- α -D-mannopyranoses, 117
- Synthesis of derivatives of methyl 3-deoxy-3-C-formyl- α -D-arabino-pentofuranoside, 293
- Synthesis of methyl 2-C-acetamidomethyl-2-deoxy- α -D-glucopyranoside and its *manno* isomer, 161
- 1-Thio- β -D-glycobiosides, 4-nitrophenyl, phase transfer catalyzed synthesis of, 149
- 1-Thioglycosides, Karplus-type equation for vicinal carbon–proton coupling constants for the C–S–C–H pathway in, 225
- Tosylepimino- α -D-mannopyranosyl]-4,5-*O*-isopropylidene-1,3-di-*N*-tosylstreptamine with potassium hydrogenfluoride, reaction of 2-deoxy-6-*O*-[2,3-dideoxy-4,6-*O*-isopropylidene-2,3-(*N*-, 363
- Transacetalation: a convenient nonaqueous method for effecting the deprotection of isopropylidene and benzylidene derivatives of sugars, 141
- 3,4,6-Tri-*O*-acetyl-2-azido-2-deoxy- α -D-glucopyranosyl chloride, use as a glycosyl donor in stereoselective syntheses of fragments of *Shigella dysenteriae* type 1 O-antigen, 103