

Subject index

- N*-Acetylated chitosans as substrates, a convenient assay for chitinase that uses partially, 373
- N*-Acetylneuraminic acid, use as a temporary blocking group in a combined chemical-enzymic oligosaccharide synthesis, C7
- N*-Acetylneuraminic acids, deoxy, incorporation into ganglioside GM₃ analogs by synthesis, 273
- N*-Acetylneuraminic acids, deoxy, synthesis of methyl thioglycosides of, for use as glycosyl donors, 257
- Affinity ligands from D-myo-inositol 1,4,5-triphosphate, synthesis of, 63 Amadori intermediates. GC-MS/SPECMA data bank identification of volatile aroma compounds, kinetics and thermal degradation of the fructose-methionine, 15
- Amino acid, C-glycosyl analogue of *O*-(β-D-xylopyranosyl)serine, synthesis of, 197
- 6-Amino-6-deoxy-D-gluconic acid, preparation of derivatives that are precursors for the synthesis of polyamides, 191
- 3-Amino-3,6-dideoxy-D-mannose (mycosamine), a precursor to the β-pyranosides of, 79
- Assay for chitinase that uses partially *N*-acetylated chitosans as substrates, 373
- Assay for oligo-(1 → 4) → (1 → 4)-glucantransferase activity in the glycogen debranching enzyme system by using HPLC with a pulsed amperometric detector, 179
- Branched-chain sugars with push-pull functionality, syntheses of, 319
- Bromination of cellulose with *N*-bromosuccinimide–triphenylphosphine under homogeneous conditions in lithium bromide–organic solvent systems, 165
- Bromosuccinimide–triphenylphosphine under homogeneous conditions in lithium bromide–organic solvent systems, bromination of cellulose with *N*-, 165
- Carbohydrate–protein conjugates, synthesis of ω-(methoxycarbonyl)alkyl and 9-(methoxycarbonyl)-3,6-dioxanonyl glycosides for use in preparation of, 117
- Cellulose with *N*-bromosuccinimide–triphenylphosphine under homogeneous conditions in lithium bromide–organic solvent systems, bromination of, 165
- Cellulose with *N*-chlorosuccinimide–triphenylphosphine under homogeneous conditions in lithium chloride–*N,N*-dimethylacetamide, 151
- Cervus nippon* Temminck, structure of the complement-activating proteoglycan from the pilose antler of, 361
- Chitinase, a convenient assay that uses partially *N*-acetylated chitosans as substrates, 373
- Chlorination of cellulose with *N*-chlorosuccinimide–triphenylphosphine under homogeneous conditions in lithium chloride–*N,N*-dimethylacetamide, 151
- Chlorosuccinimide–triphenylphosphine under homogeneous conditions in lithium chloride–*N,N*-dimethylacetamide, chlorination of cellulose with *N*-, 151
- Complement-activating proteoglycan from the pilose antler of *Cervus nippon* Temminck, structure of, 361
- Complexation with C₁₂ cyclic compounds, enhanced production of cyclomalto-octaose through selective, 343
- Conformational analysis of the anomeric forms of kojibiose, nigerose, and maltose using MM3, 223
- Conformational behaviour of the cardiac glycoside digoxin as indicated by NMR spectroscopy and molecular dynamics calculations, 201
- Crystal packing of *N*-(*n*-octyl)-D-gulonamide containing tail-to-tail sheets compared to its gluconamide diastereomer showing head-to-tail arrangement, 31
- Cyclomaltoheptaose, regioselectivity of the insertion reactions of some aromatic diazo compound complexes with, 289
- Cyclomalto-octaose through selective complexation with C₁₂ cyclic compounds, enhanced production of, 343

- 2,3-Dideoxy-2,3-(*N*-tosylepimino)- α -D-allopyranosides, and synthesis of 3'-deoxy-3'-fluorokanamycin B and 3',4'-dideoxy-3'-fluorokanamycin B, study on fluorination of, 89
- Digoxin, conformational behaviour as indicated by NMR spectroscopy and molecular dynamics calculations, 201
- Enhanced production of cyclomaltooctaose through selective complexation with C₁₂ cyclic compounds, 343
- Esterase from rabbit serum, partially pivaloylated D-glucopyranose substrates for the, 107
- Fluorination of 2,3-dideoxy-2,3-(*N*-tosylepimino)- α -D-allopyranosides, and synthesis of 3'-deoxy-3'-fluorokanamycin B and 3',4'-dideoxy-3'-fluorokanamycin B, study on, 89
- Fluorokanamycin B, study on fluorination of 2,3-dideoxy-2,3-(*N*-tosylepimino)- α -D-allopyranosides, and synthesis of 3'-deoxy-3'-fluorokanamycin B and 3',4'-dideoxy-3'-, 89
- Fructose-methionine Amadori intermediates. GC-MS/SPECMA data bank identification of volatile aroma compounds, kinetics and thermal degradation of the, 15
- Furanose conformer, a locked ⁴T₃. Synthesis and X-ray crystal and solution structures of 2,5-anhydro-3,4-*O*-(1,2-ethanediyl)-D-mannitol, 213
- Ganglioside GM₃ analogs containing a deoxy-*N*-acetylneuraminic acid residue, synthesis of a series of, 273
- Glucantransferase activity in the glycogen debranching enzyme system, assay for oligo-(1 → 4) → (1 → 4)-, by using HPLC with a pulsed amperometric detector, 179
- Gluconamide diastereomer showing head-to-tail arrangement, the crystal packing of *N*-(*n*-octyl)-D-gulonamide containing tail-to-tail sheets compared to its, 31
- D-Gluconic acid, preparation of derivatives of 6-amino-6-deoxy-, as precursors for the synthesis of polyamides, 191
- D-Glucopyranoses, syntheses of partially pivaloylated, 107
- Glycogen debranching enzyme system, assay for oligo-(1 → 4) → (1 → 4)-glucantransferase activity by using HPLC with a pulsed amperometric detector, 179
- C-Glycosyl amino acid analogue of *O*-(β -D-xylopyranosyl)serine, synthesis of, 197
- Hyaluronic acid esters and salts, NMR studies of hydrogen-bond patterns and conformational behaviour for solutions in methyl sulfoxide, 1
- Hydrogen-bond patterns and conformational behaviour of salts and esters of hyaluronic acid in solution in methyl sulfoxide, NMR studies of, 1
- Hydroxyl → halogen exchange reaction in the presence of triphenylphosphine, *N*-bromosuccinide, and *N,N*-dimethylformamide, the mechanism of the, 327
- Inhibitors of neural cell division, synthesis of oligosaccharide, 185
- D-*myo*-Inositol 1,4,5-triphosphate affinity ligands, synthesis of, 63
- Insertion reactions of some aromatic diazo compound complexes with cyclomaltoheptaose, regioselectivity of the, 289
- Intramolecular hydrogen bonds in methyl β -cellobioside, a ¹H-NMR and MD study, 41
- Kinetics and thermal degradation of the fructose-methionine Amadori intermediates. GC-MS/SPECMA data bank identification of volatile aroma compounds, 15
- Kojibiose, nigerose, and maltose, conformational analysis of the anomeric forms of using MM3, 223
- Maltose, kojibiose, and nigerose, conformational analysis of the anomeric forms of using MM3, 223
- D-Mannitol, 2,5-anhydro-*O*-(1,2-ethanediyl)-, a locked ⁴T₃ furanose conformer: synthesis and X-ray crystal and solution structures of, 213
- α -D-Mannopyranosyl and α -D-mannopyranosyl-(1 → 2)- α -D-mannopyranosyl linked to L-serine and L-threonine, synthesis and conformational and NMR studies of, 299
- (Methoxycarbonyl)alkyl and 9-(methoxycarbonyl)-3,6-dioxanonyl glycosides for use in preparation of carbohydrate-protein conjugates, synthesis of ω -, 117
- Methyl β -cellobioside, a ¹H-NMR and MD study of intramolecular hydrogen bonds in, 41
- MM3, conformational analysis of the anomeric forms of kojibiose, nigerose, and maltose using, 223
- Mucin-type core unit, synthesis of neoglycolipids containing a, C11
- Mycosamine (3-amino-3,6-dideoxy-D-mannose), a precursor to the β -pyranosides of, 79
- Neoglycolipids containing a mucin-type core unit, synthesis, C11

- Neural cell division, synthesis of oligosaccharide inhibitors of, 185
- Nigerose, kojibiose, and maltose, conformational analysis of the anomeric forms of using MM3, 223
- NMR and MD study of intramolecular hydrogen bonds in methyl β -cellobioside, ^1H -, 41
- NMR spectroscopy and molecular dynamics calculations, conformational behaviour of the cardiac glycoside digoxin as indicated by, 201
- NMR studies of solutions of hyaluronic acid esters and salts in methyl sulfoxide, comparison of hydrogen-bond patterns and conformational behaviour, 1
- Oligosaccharide inhibitors of neural cell division, synthesis of, 185
- Pivaloylated D-glucopyranoses, syntheses of partially, 107
- Polyamides, preparation of derivatives of 6-amino-6-deoxy-D-gluconic acid that are precursors for the synthesis of, 191
- Precursor to the β -pyranosides of 3-amino-3,6-dideoxy-D-mannose (mycosamine), 79
- Proteoglycan from the pilose antler of *Cervus nippon* Temminck, structure of the complement-activating, 361
- Regioselectivity of the insertion of some aromatic diazo compound complexes with cyclo-maltoheptaose, 289
- L-Serine and L-threonine, synthesis and conformational and NMR studies of α -D-mannopyranosyl and α -D-mannopyranosyl-(1 \rightarrow 2)- α -D-mannopyranosyl linked to, 299
- Sialyl dimeric Lewis^x heptasaccharide, combined chemical-enzymic synthesis, C7
- Sialyl X ceramide consisting of a pentasaccharide recognized by the selectin family, stereocontrolled synthesis of, C1
- Structures, X-ray crystal and solution, and synthesis of 2,5-anhydro-3,4-O-(1,2-ethanediyl)-D-mannitol, a locked 4T_3 furanose conformer, 213
- Sugars with push-pull functionality, syntheses of branched-chain, 319
- Sulfated alkyl glycosides, synthesis and properties, 245
- Syntheses of branched-chain sugars with push-pull functionality, 319
- Synthesis and conformational and NMR studies of α -D-mannopyranosyl and α -D-mannopyranosyl-(1 \rightarrow 2)- α -D-mannopyranosyl linked to L-serine and L-threonine, 299
- Synthesis and properties of sulfated alkyl glucosides, 245
- Synthesis of ω -(methoxycarbonyl)alkyl and 9-(methoxycarbonyl)-3,6-dioxanonyl glycosides for use in preparation of carbohydrate-protein conjugates, 117
- Synthesis of sialyl X ceramide consisting of a pentasaccharide recognized by the selectin family, stereocontrolled, C1
- Thioglycosides of deoxy-N-acetylneuraminic acids for use as glycosyl donors, synthesis of the methyl, 257
- Triphenylphosphine, N-bromosuccinimide, and N,N-dimethylformamide, the mechanism of the hydroxyl \rightarrow halogen exchange reaction in the presence of, 327
- Vilsmeier-type reagent, application in carbohydrate chemistry of a new, 327
- VIM-2 hexasaccharide determinant, combined chemical-enzymic synthesis, C7