

Iterative one-pot syntheses of chitotetroses

Lijun Huang,^a Zhen Wang,^a Xiaoning Li,^a Xin-shan Ye^b and Xuefei Huang^{a,*}

^aDepartment of Chemistry, The University of Toledo, 2801 W. Bancroft Street, MS 602, Toledo, OH 43606, USA

^bThe State Key Laboratory of Natural and Biomimetic Drugs, School of Pharmaceutical Sciences, Peking University, Xue Yuan Road 38, Beijing 100083, China

Received 7 November 2005; received in revised form 3 January 2006; accepted 8 January 2006

Available online 26 January 2006

Dedicated to Professor Koji Nakanishi on the occasion of his 80th birthday

Abstract—Rapid syntheses of chitotetrose derivatives were achieved in good yields using the newly developed reactivity independent iterative one-pot strategy. The protective groups on donors and acceptors were independently evaluated allowing matching of the two partners in glycosylation. No anomeric reactivity adjustments or intermediate purification were necessary thus significantly improving the overall synthetic efficiency. Only near stoichiometric amounts of building blocks were required for the assembly of target molecules further highlighting the potential of the iterative one-pot method in complex oligosaccharide synthesis. © 2006 Elsevier Ltd. All rights reserved.

Keywords: Iterative; One-pot synthesis; Chitotetrose; Reactivity independent

1. Introduction

With the recognition of multifaceted biological properties of oligosaccharides and glyco-conjugates, the development of novel methods to expedite oligosaccharide synthesis has been a major focus of modern synthetic carbohydrate chemistry.^{1,2} A popular synthetic strategy is the reactivity based chemoselective glycosylation method, where a more reactive armed glycosyl donor is preferentially activated in the presence of a less reactive disarmed acceptor.^{2,3} Many innovative methods have been developed for tuning anomeric reactivities.^{3–5} However, the very need to achieve precise anomeric reactivity values necessitates extensive and tedious protecting group and/or aglycon adjustments, which in turn detracts from its broad application and restricts the possibility of matching⁶ the glycosyl donor with the acceptor.

Recently, we have developed a new iterative one-pot glycosylation method, by which multiple sequential glycosylations of a thioglycosyl donor by a thioglycosyl

acceptor can be carried out *independent* of the anomeric reactivities of donors and acceptors.⁷ Medium sized oligosaccharides have been assembled in a few hours using this methodology without the need of intermediate purification and aglycon adjustments. Our reactivity independent one-pot approach represents a significant advancement in chemoselective glycosylation as the time-consuming anomeric reactivity adjustment is unnecessary. To explore the scope of this method, we have embarked on application of this strategy in assembly of complex oligosaccharides. Herein, we report our syntheses of chitotetroses, that is, tetrasaccharides containing β -(1→4)-linked glucosamines.

2. Results and discussion


Chitotetroses and their derivatives have diverse biological functions such as immunostimulatory,⁸ antitumor,⁹ anticoagulant,¹⁰ and anti-HIV activities.¹¹ Chemical synthesis of this class of compounds presents a significant challenge due to the well known low nucleophilicity of 4-hydroxyl group of glucosamine building blocks. To date, there have been a few reports of step-wise assembly

* Corresponding author. Tel.: +1 419 530 1507; fax: +1 419 530 4033; e-mail: xuefei.huang@utoledo.edu

of these molecules, which are often characterized by multiple protecting group manipulations and aglycon leaving group adjustments on oligosaccharide intermediates.^{12,13} A single one-pot synthesis was accomplished by our group using the reactivity based chemoselective glycosylation approach.⁴ Even though building blocks with multiple levels of anomeric reactivities were conveniently generated through post-synthetic modification of the aglycon of a common intermediate, the need for reactivity adjustment of our previous one-pot synthesis prompted us to explore the utility of our reactivity independent iterative one-pot method.⁷


It is known that the low nucleophilicity of 4-hydroxyl group of protected *N*-acetyl glucosamine is mainly due to the steric hindrance around this secondary hydroxyl group and the intermolecular hydrogen bonding with the 2-acetamido moiety.¹⁴ The *N*-protecting group on a glycosyl acceptor can significantly affect its nucleophilicity, with the following order of activities: N_3 , $NH\text{-Troc} > N\text{-Phth} > NH\text{-Ac}$.^{14,15} Recently, Crich and Vinod have demonstrated that the usage of oxazolidinone protected *N*-acetyl glucosamine O-glycoside acceptor **1** considerably enhanced the glycosylation yield compared with the corresponding O-glycoside without the oxazolidinone moiety.^{16,17} We anticipated that oxazolidinone protected thioglycosyl acceptor **2** may lead to higher glycosylation yield and the resulting disaccharide can function as a glycosyl donor.¹⁸ To investigate the influence of *N*-protecting groups on our glycosylation methodology, glycosyl acceptors **2**, **3**, and **4** were prepared. The azido moiety is unsuitable as an *N*-protecting group for one-pot synthesis of β -linked oligoglucosamines because it is a well known non-participatory group favoring the formation of α -anomer when used in a donor.¹⁹

The synthesis of oxazolidinone protected acceptor **2** started from the benzylidene protected glucosamine derivative **5** (Scheme 1).²⁰ Removal of the phthalimido group followed by oxazolidinone formation and acetylation provided the fully protected glucosamine **6**. Interest-



ingly, regioselective reductive opening of the benzylidene ring by treatment of NaCNBH_3 and HCl ⁴ yielded a mixture of α - and β -thioglycoside **2**, with **2 α** as the major product (40%). This is in line with the observation by the Crich group where the oxazolidinone bearing O-glycoside **7** was epimerized at the anomeric position under similar reaction conditions.¹⁶ Thioglycosyl acceptors **3**²¹ and **4**²² were synthesized in a straightforward manner from building block **5** (Scheme 1b and c).


Glycosylations of thioglycosyl donor **8**²⁰ with acceptors **2** to **4** were carried out following our previously established pre-activation condition.⁷ The disarmed glycosyl donor **8** was cleanly activated in the absence of any acceptors at -70°C by the powerful thiophilic promoter *p*-TolSOTf, formed in situ by reaction of *p*-TolSCl²³ with AgOTf.^{7,24} Subsequent addition of the glycosyl acceptor (0.9 equiv) to the reaction mixture led to the formation of thioglycosyl disaccharide. The reaction of *N*-Phth protected acceptor **3** with donor **8** produced the desired disaccharide **9** in 56% yield with trace amount ($<5\%$) of donor elimination product glycal **10** (Table 1, entry 1). It is noteworthy that acceptor **3** has a higher anomeric reactivity than donor **8**. This reversal of anomeric reactivity, that is, chemoselective glycosyl-



Scheme 1. Preparation of acceptors **2–4**. Reagents and conditions: (i) ethylene diamine, EtOH, reflux, overnight; (ii) 4-O₂NC₆H₄OCOC₂H₅, NaHCO₃, CH₃CN/H₂O, rt, overnight; (iii) AcCl, DIPEA, DCM, rt, overnight; (iv) NaCNBH₃, 1 M HCl in Et₂O, 0°C , 1.5 h; (v) NaH, TBAI, DMF, MS; BnBr, 2 h, rt; (vi) Troc-Cl, NaHCO₃, H₂O, THF.

Table 1. Evaluation of *N*-protective group on glycosylation of various thioglycosyl acceptors by donor **8**

Donor 8 (1 eq) + AgOTf		1) <i>p</i> -TolSCl, -70 °C 2) acceptor (0.9 eq)		Product
Entry #	Acceptor	Pdt.	Yield (%)	
1	3	9	56	
2	4	11	50	
3	12	13	50	
4	2a	14	39	



ation of an armed acceptor with a disarmed donor is not possible with the traditional reactivity based glycosylation methods.


The glycosylation of *N*-Troc protected acceptor **4** by donor **8** gave a yield of 50% of the disaccharide **11**, along with 20% of regenerated donor **8** even after complete consumption of the donor during pre-activation (Table 1, entry 2). This is the first time that a substantial amount of the regenerated donor was observed among all reactions we carried out so far using the pre-activation scheme. The donor regeneration is presumably due to the intermolecular aglycon transfer²⁵ from the acceptor to the activated donor via S-alkylation instead of the desired O-alkylation of the hydroxyl group. This further re-affirms the low nucleophilicity of 4-hydroxyl group of glucosamine acceptors. The usage of *p*-nitro phenyl thioglycosyl acceptor **12**⁴ suppressed the aglycon transfer side reaction, but the disaccharide product **13** was obtained in only 50% yield (Table 1, entry 3). We examined next the glycosylation of oxazolidinone thio-


glycosyl acceptor **2a** by donor **8**. Disappointingly, the disaccharide product **14** was produced in only 39% yield with 50% of acceptor **2a** recovered after reaction (Table 1, entry 4). The oxazolidinone moiety did not seem to improve the glycosylation and *N*-Phth was used as the *N*-protecting group in further studies.

The four-component one-pot assembly of fully protected tetraglucosamine derivative **16** was performed as outlined in Scheme 2. Pre-activation of donor **8** by *p*-TolSOTf was followed by addition of acceptor **3**. The reaction temperature was raised to -10 °C in 20 min. After acceptor **3** was completely consumed as judged by TLC analysis, the reaction mixture was cooled back down to -70 °C, followed by sequential addition of AgOTf, *p*-TolSCl, acceptor **3**, and warming up to -10 °C. Subsequently, the reaction temperature was lowered to -70 °C again, which was followed by addition of acceptor **15**⁴ and AgOTf/*p*-TolSCl. Slight excess of glycosyl donors was employed for the first two glycosylations to ensure complete consumption of the acceptor. The excess activated donor decomposed upon warming up thus not affecting the following reactions. The desired protected chitotetrose **16** was isolated from the one-pot reaction by flash column chromatography in 32% overall yield. It is noteworthy that the same acceptor **3** was used for the formation of the first and the second glycosidic linkages without resorting to anomeric reactivity adjustment.

To enhance the glycosylation yield, we next explored the protecting groups on the glycosyl donor. From the readily available alcohol **5**,²⁰ benzylation and silylation gave donor **17**²¹ and **18** in 90% and 85% yields, respectively (Scheme 3a and b). Regioselective opening of the benzylidene group with TMSOTf and borane THF²⁶ followed by benzylation produced the benzyl protected donor **19** in 80% yield for the two steps (Scheme 3c).

With various donors in hand, we began to evaluate their effects on glycosylation. The benzyl and benzylidene containing donor **17** reacted with acceptor **3** giving 61% of disaccharide **20** (Table 2, entry 1) together with 20% of recovered acceptor **3**. Exchange of the benzyl group in **17** with TBS moiety (donor **18**) led to a much

**Scheme 2.** One-pot synthesis of chitotetrose **16**.




Scheme 3. Preparation of donors **17–19**. Reagents and conditions: (i) NaH, TBAI, DMF, MS; BnBr, 2 h, rt; (ii) TBSOTf, 2,6-lutidine, DCM, 4 h, 0 °C; (iii) TMSOTf, BH₃·THF, 4 h, –20 to 0 °C.

cleaner reaction with complete consumption of the acceptor, producing disaccharide **21** in 74% yield (Table 2, entry 2). Both the TBS and benzylidene groups are important for high yield, because donor **19**, which is devoid of the benzylidene group, glycosylated acceptor **3** in a lower 50% yield (Table 2, entry 3). 1,1'-Linked disaccharide **23** and glycal **24** were isolated as the major side products (~10%) for this reaction. The enhancement effect of TBS group is currently under investigation.


One-pot sequential reactions of **18**, **3**, **25**,²⁷ and **26**¹³ promoted by *p*-TolSOTf led to the formation of tetrasaccharide **27** in 45% yield in less than 6 h (Scheme 4). This corresponds to an average of 85% yield for each step of the five synthetic steps carried out in one pot. The reactivity independent nature of our method allowed us to utilize the PMB containing building block **25** instead of acceptor **3** for the second glycosylation. To prevent the loss of acid labile PMB group, a steri-

Table 2. Evaluation of donors on glycosylation of acceptor **3**

Donor (1 eq) + AgOTf		1) <i>p</i> -TolSOTf, -70 °C 2) acceptor 3 (0.9 eq)	Product
Entry #	Donor	Pdt.	Yield (%)
1	17	20	61
2	18	21	74
3	19	22	50



cally hindered base TTBP²⁸ was added with each acceptor. In addition to the desired product **27** and final acceptor **26** (50%), we have isolated from the final reaction mixture, glucals **28** (18%), **29** (15%), and hemiacetal **30** (24%) as the major side products, which were formed through elimination or hydrolysis of activated glycosyl donors due to insufficient nucleophilicity of the acceptors. The glucals do not react with the promoter *p*-TolSOTf at low temperature, which do not affect further donor activation. In addition, these side products have sufficiently different polarities from the desired oligosac-



Scheme 4. One-pot synthesis of chitotetrose **27**.

charide **27**, thus not interfering with product purification by flash chromatography. The PMB and TBS groups in tetrasaccharide **27** can be selectively removed to allow for alkylations of alternating 3-hydroxyl groups in the tetraglucosamine, which are useful intermediates for peptidoglycan synthesis.²⁹

In conclusion, facile syntheses of chitotetrose derivatives were achieved in good yields using our newly developed reactivity independent iterative one-pot method. Unlike solid phase synthesis,³⁰ only near stoichiometric amounts of building blocks were required. No anomeric reactivity differentiations between donors and acceptors were necessary, thus allowing independent evaluation of protecting groups. As the *N*-protecting group, the phthalimido moiety gave the most consistent results, while the new oxazolidinone group did not improve the glycosylation. The introduction of benzylidene and TBS groups onto thioglycosyl donor significantly enhanced the yield. Further improvements on this challenging glycosylation reaction will depend upon the augmentation of the nucleophilicity of 4-hydroxyl group of glucosamine acceptors.

3. Experimental

3.1. General procedures

All reactions were carried out under nitrogen with anhydrous solvents in flame-dried glassware, unless otherwise noted. All glycosylation reactions were performed in the presence of molecular sieves, which were flame-dried right before the reaction under high vacuum. Glycosylation solvents were dried using an MBraun solvent purification system. Chemicals used were reagent grade as supplied except where noted. Analytical thin-layer chromatography was performed using silica gel 60 F254 glass plates (EM Science); compound spots were visualized by UV light (254 nm) and by staining with a yellow solution containing $\text{Ce}(\text{NH}_4)_2(\text{NO}_3)_6$ (0.5 g) and $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}\cdot 4\text{H}_2\text{O}$ (24.0 g) in 6% H_2SO_4 (500 mL). Flash column chromatography was performed on silica gel 60 (230–400 Mesh, EM Science). ^1H NMR, ^{13}C NMR, ^1H – ^1H COSY, and ^1H – ^{13}C HMQC spectra were recorded on a Varian VXR-400 or Inova-600 instrument and were referenced using Me_4Si (0 ppm), residual CHCl_3 (δ ^1H NMR 7.26 ppm, ^{13}C NMR 77.0 ppm). Optical rotations were measured at 25 °C. ESI mass spectra were recorded on ESQUIRE LC–MS operated in positive ion mode. High-resolution mass spectra were recorded on a Micromass electrospray ToF™ II (Micromass, Wythenshawe, UK) mass spectrometer equipped with an orthogonal electrospray source (Z-spray) operated in positive ion mode, which is located at the Mass Spectrometry and Proteomics Facility, the Ohio State University.

3.2. *p*-Tolyl 2-acetamido-4,6-*O*-benzylidene-2-*N*-3-*O*-carbonyl-2-deoxy-1-thio- β -D-glucopyranoside (**6**)

p-Tolyl 4,6-*O*-benzylidene-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside **5** (1.5 g, 2.98 mmol, 1 equiv) and ethylenediamine (9 g, 150 mmol, 50 equiv) were dissolved in dry ethanol (15 mL). The mixture was heated at reflux for 16 h. The reaction mixture was cooled and neutralized with 2 N HCl until slightly basic. After concentration, the crude product was dissolved in EtOAc (10 mL) and washed with H_2O , brine, then dried, and concentrated. Column chromatography (15:1, CH_2Cl_2 –MeOH) on silica gel afforded the free amine (0.98 g, 88%). The obtained amine (0.98 g, 2.62 mmol, 1 equiv) and NaHCO_3 (1.1 g, 13.09 mmol, 5 equiv) were dissolved in $\text{H}_2\text{O}/\text{CH}_3\text{CN}$ mixture (15 mL/15 mL) and cooled to 0 °C. *p*-Nitrophenyl chloroformate (1.59 g, 7.89 mmol, 3 equiv) in CH_3CN (5 mL) was added slowly and the mixture was stirred at 0 °C for 3 h. The resulting aqueous mixture was extracted with EtOAc and the combined extracts were washed with water, brine, then dried, and concentrated. Silica gel column chromatography (2.5:1, hexanes–EtOAc) afforded the oxazolidinone protected glucosamine as a white solid (0.88 g, 84%). The oxazolidinone protected glucosamine (0.42 g, 1.05 mmol, 1 equiv) and Hünig's base (0.92 mL, 5.29 mmol, 5 equiv) were dissolved in CH_2Cl_2 (6 mL). Acetyl chloride (0.38 mL, 5.34 mmol, 5.1 equiv) was added under N_2 . The mixture was stirred for 6 h and then poured into saturated aqueous NaHCO_3 . The mixture was diluted with CH_2Cl_2 (5 mL) and the organic layer was separated. The aqueous layer was extracted twice with CH_2Cl_2 and the combined extracts were washed with H_2O , brine, then dried, and concentrated. Column chromatography (1:1, hexanes–EtOAc) on silica gel afforded the product **11** as a white solid (0.39 g, 84%). ^1H NMR (600 MHz, CDCl_3): δ 2.33 (s, 3H), 2.58 (s, 3H), 3.50–3.54 (m, 1H), 3.92 (t, J = 10.2 Hz, 1H), 4.07 (t, J = 9.3 Hz, 1H), 4.13 (dd, J = 9.0, 10.8 Hz, 1H), 4.27 (dd, J = 4.8, 10.5 Hz, 1H), 4.33 (t, J = 0.5 Hz, 1H), 4.91 (d, J = 9.0 Hz, 1H), 5.59 (s, 1H), 7.11–7.12 (m, 2H), 7.38–7.35 (m, 5H), 7.43–7.44 (m, 2H); ^{13}C NMR (150 MHz, CDCl_3): δ 21.42, 25.17, 61.01, 68.46, 73.31, 78.60, 78.97, 88.95, 101.77, 126.35, 128.60, 129.65, 129.87, 130.02, 133.20, 136.45, 138.79, 153.81, 173.54; HRMS: $[\text{M}+\text{Na}]^+$ $\text{C}_{23}\text{H}_{23}\text{NNaO}_6\text{S}$ calcd 464.1144, obsd 464.1181.

3.3. *p*-Tolyl 2-acetamido-6-*O*-benzyl-2-*N*,3-*O*-carbonyl-2-deoxy-1-thio- α -D-glucopyranoside (**2 α**)

The mixture of compound **6** (0.11 g, 0.25 mmol, 1 equiv), NaCNBH_3 (0.19 g, 3.0 mmol, 12 equiv) in THF (5 mL) was cooled to 0 °C. One molar of HCl in Et_2O (~3 mL) was added under N_2 until the solution

was slightly acidic. The reaction mixture was stirred at 0 °C for 1.5 h and diluted with 10 mL of CH₂Cl₂ and then poured into aqueous NaHCO₃. The organic layer was separated and washed with H₂O, brine, then dried, and concentrated. Column chromatography (1:1, hexanes–EtOAc) on silica gel afforded the product **2a** as a white solid (44 mg, 40%). [α]_D +176.4 (*c*, 0.0056 g/mL, CH₂Cl₂); ¹H NMR (600 MHz, CDCl₃): δ 2.31 (s, 3H), 2.53 (s, 3H), 3.11 (br, 1H), 3.72–3.74 (m, 1H), 3.83–3.86 (m, 1H), 4.01–4.04 (m, 1H), 4.08–4.11 (m, 1H), 4.16–4.19 (m, 1H), 4.37 (dd, *J* = 12, 9.6 Hz, 1H), 4.53 (d, *J* = 12.0 Hz, 1H), 4.61 (d, *J* = 12 Hz, 1H), 6.07 (d, *J* = 4.2 Hz, 1H), 7.08–7.09 (m, 2H), 7.29–7.36 (m, 7H); ¹³C NMR (150 MHz, CDCl₃): δ 21.37, 24.06, 59.91, 69.72, 70.70, 72.51, 74.07, 78.41, 86.88, 128.06, 128.31, 128.82, 128.92, 130.21, 133.27, 137.55, 138.70, 153.32, 171.45; HRMS: [M+Na]⁺ C₂₃H₂₅NNaO₆S calcd 466.1300, obsd 466.1271.

3.4. *p*-Tolyl 3,6-di-*O*-benzyl-2-deoxy-2-*N*-phthalimido-1-thio- β -D-glucopyranoside (**3**)

The mixture of compound **17** (1 g, 1.68 mmol), NaC–NBH₃ (1.06 g, 17 mmol), and molecular sieves MS AW300 in THF (20 mL) was cooled to 0 °C. Two molar of HCl in Et₂O (~9 mL) was added under N₂ until no more gas evolved from the reaction. The reaction mixture was stirred at 0 °C for 30 min. The reaction was quenched with Et₃N and filtered. The filtrate was diluted with CH₂Cl₂ (30 mL) and extracted with saturated solutions of aqueous NH₄Cl (20 mL) and NaHCO₃ (20 mL). The organic layer was separated and dried with Na₂SO₄. Column chromatography (2:1, hexanes–EtOAc) on silica gel afforded the product **3** as a white solid in 90% yield (80% yield for the two steps from compound **5**). Comparison of the NMR data with those reported in the literature²¹ confirmed the identity of **3**.

3.5. *p*-Tolyl 3,6-di-*O*-benzyl-2-deoxy-1-thio-2-*N*-trichloroethoxycarbonyl- β -D-glucopyranoside (**4**)

Compound **17** (1.9 g, mmol) and ethylenediamine (11.8 mL, 175 mmol, 55 equiv) were dissolved in dry ethanol (30 mL). The mixture was heated at reflux for 16 h. The reaction mixture was evaporated and dissolved in EtOAc (30 mL), which was neutralized with 1 N H₂SO₄ until slightly basic. The resulting suspension was kept in ice-water for 1 h, and then filtered to afford the glucosamine sulfate salt (1.53 g, 85%) as a white solid. The solid (1.05 g, 1.86 mmol) was dissolved in a mixture of THF and saturated aqueous NaHCO₃ solution (30 mL) followed by addition of TrocCl (340 μ L, 2.51 mmol). The mixture was stirred for 1 h, and then extracted with EtOAc (60 mL) and saturated aqueous NaHCO₃ (20 mL). The organic layer was dried over Na₂SO₄ and purified by silica gel column to give 2-

deoxy-2-*N*-Troc glucopyranoside (1.19 g, 100%). The desired compound **4** (660 mg, 90%) was then obtained as a white solid from the 2-deoxy-2-*N*-Troc glucopyranoside (730 mg, 1.14 mmol) following the same procedure as in the synthesis of compound **3**. The overall yield for synthesis of compound **4** was 70% for the four steps starting from compound **5**. Comparison of the NMR data with those reported in the literature²² confirmed the identity of **4**.

3.6. General procedure for single step glycosylation

A solution of donor (0.067 mmol) and freshly activated molecular sieve MS–4 Å (300 mg) in a mixture of CH₂Cl₂ and Et₂O (1:1, 4 mL) (100% Et₂O was used for donors **18** and **19**) was stirred for 30 min, and cooled to –70 °C, which was followed by sequential additions of AgOTf (52 mg, 0.201 mmol) dissolved in Et₂O (1 mL) and *p*-TolSCl (10.5 μ L, 0.067 mmol) through syringes. After the donor was completely consumed according to TLC analysis (about 15 min at –70 °C), a solution of acceptor (0.060 mmol) in CH₂Cl₂ (0.5 mL) was injected via a syringe. The reaction mixture was warmed to –10 °C under stirring in 2 h. Then the mixture was diluted with CH₂Cl₂ (20 mL) and filtered. The filtrate was washed twice with saturated aqueous NaHCO₃ solution (20 mL) and once with brine (10 mL). The organic layer was collected and dried over MgSO₄. After removal of the solvent, the residue was applied to silica gel flash chromatography column using a mixture of hexanes and EtOAc to afford the desired disaccharide.

3.7. *p*-Tolyl 3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (**9**)

Compound **9** was synthesized from donor **8** and acceptor **3** in 56% yield following the general procedure of single step glycosylation together with trace amount of glycal **10**. Compound **9** [α]_D +3.5 (*c*, 0.058 g/mL, CH₂Cl₂); ¹H NMR (600 MHz, CDCl₃): δ 1.82 (s, 3H), 1.93 (s, 3H), 1.98 (s, 3H), 2.20 (s, 3H), 3.35 (dd, *J* = 3.0, 9.6 Hz, 1H), 3.44–3.48 (m, 2H), 3.58 (d, *J* = 10.8 Hz, 1H), 3.89 (dd, *J* = 2.4, 12.0 Hz, 1H), 4.08–4.22 (m, 4H), 4.30 (dd, *J* = 8.4, 10.2 Hz, 1H), 4.44 (d, *J* = 12.6 Hz, 1H), 4.48 (d, *J* = 11.4 Hz, 1H), 4.54 (d, *J* = 11.4 Hz, 1H), 4.79 (d, *J* = 12.6 Hz, 1H), 5.11 (t, *J* = 9.6 Hz, 1H), 5.26 (d, *J* = 9.6 Hz, 1H), 5.49 (d, *J* = 8.4 Hz, 1H), 5.78 (dd, *J* = 9.0, 10.2 Hz, 1H), 6.79–7.88 (m, 22 H); ¹³C NMR (150 MHz, CDCl₃): δ 20.68, 20.86, 20.86, 21.31, 54.93, 55.46, 60.64, 68.38, 68.96, 70.82, 71.70, 72.94, 74.63, 76.15, 77.89, 78.81, 83.50, 97.03, 127.27–138.64 (aromatic carbons), 169.75, 170.36, 170.93 (CH₃CO); HRMS: [M+Na]⁺ C₅₅H₅₂N₂NaO₁₅S calcd 1035.2986, obsd 1035.2985. Compound **10**: [α]_D –22.5

(*c*, 0.0067 g/mL, CH₂Cl₂); ¹H NMR (600 MHz, CDCl₃): δ 1.92 (s, 3H), 2.12 (s, 3H), 2.14 (s, 3H), 4.36 (dd, *J* = 4.2, 12.0 Hz, 1H), 4.47–4.58 (m, 2H), 5.31 (t, *J* = 4.2 Hz, 1H), 5.59 (d, *J* = 4.2 Hz, 1H), 6.76 (s, 1H), 7.72–7.76 (m, 2H), 7.83–7.88 (m, 2H); HRMS [M+Na]⁺ C₂₀H₁₉NNaO₉ calcd 440.0958, obsd 440.0961.

3.8. *p*-Tolyl 3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido-β-D-glucopyranosyl-(1→4)-3,6-di-*O*-benzyl-2-deoxy-1-thio-2-*N*-trichloroethoxycarbonyl-β-D-glucopyranoside (11)

Compound **11** was synthesized from donor **8** and acceptor **4** in 50% yield following the general procedure of single step glycosylation together with recovered acceptor **4** (20%). [α]_D −10.7 (*c*, 0.012 g/mL, CH₂Cl₂); ¹H NMR (600 MHz, CDCl₃): δ 1.82 (s, 3H), 1.94 (s, 3H), 1.98 (s, 3H), 2.25 (s, 3H), 3.26 (m, 2H), 3.40 (d, *J* = 10.2 Hz, 1H), 3.45 (dd, *J* = 3.6, 10.8 Hz, 1H), 3.56 (d, *J* = 10.2 Hz, 1H), 3.75–3.83 (m, 2H), 4.08 (t, *J* = 9.0 Hz, 1H), 4.11 (dd, *J* = 4.2, 12.6 Hz, 1H), 4.27 (dd, *J* = 9.0, 10.8 Hz, 1H), 4.42 (d, *J* = 12 Hz, 1H), 4.48 (d, *J* = 12 Hz, 1H), 4.62–4.65 (m, 2H), 4.70 (d, *J* = 12 Hz, 1H), 4.76 (d, *J* = 4.2 Hz, 1H), 4.91 (d, *J* = 11.4 Hz, 1H), 5.03 (d, *J* = 7.8 Hz, 1H), 5.10 (t, *J* = 9.6 Hz, 1H), 5.50 (d, *J* = 8.4 Hz, 1H), 5.74 (t, *J* = 10.2 Hz, 1H), 6.95–7.84 (18, 2H); ¹³C NMR (150 MHz, CDCl₃): δ 20.66, 20.84, 20.89, 21.33, 55.43, 56.30, 61.66, 68.44, 68.86, 70.79, 71.88, 73.00, 74.40, 74.59, 75.33, 78.80, 79.76, 85.62, 97.10, 127.63–138.42 (aromatic carbons), 153.71 (carbonyl of Troc), 169.70, 170.35, 170.88 (CH₃CO); HRMS: [M+Na]⁺ C₅₀H₅₁Cl₃N₂NaO₁₅S calcd 1079.1973, obsd 1079.2067.

3.9. *p*-Nitrophenyl 3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido-β-D-glucopyranosyl-(1→4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido-1-thio-β-D-glucopyranoside (13)

Compound **13** was synthesized from donor **8** and acceptor **12** in 50% yield following the general procedure of single step glycosylation. [α]_D +21.8 (*c*, 0.010 g/mL, CH₂Cl₂); ¹H NMR (600 MHz, CDCl₃): δ 1.84 (s, 3H), 1.94 (s, 3H), 1.99 (s, 3H), 3.49–3.55 (m, 3H), 3.61 (d, *J* = 9.6 Hz, 1H), 3.96 (dd, *J* = 2.4, 12.0 Hz, 1H), 4.19–4.23 (m, 3H), 4.25 (dd, *J* = 3.6, 12.0 Hz, 1H), 4.33 (dd, *J* = 8.4, 10.8 Hz, 1H), 4.44 (d, *J* = 12 Hz, 1H), 4.45 (d, *J* = 12.6 Hz, 1H), 4.53 (d, *J* = 11.4 Hz, 1H), 4.82 (d, *J* = 12.6 Hz, 1H), 5.13 (t, *J* = 10.2 Hz, 1H), 5.47 (d, 1H, *J* = 10.2 Hz, 1H), 5.50 (d, *J* = 8.4 Hz, 1H), 5.79 (dd, *J* = 9.0, 10.8 Hz, 1H), 6.79–7.94 (m, 22H); ¹³C NMR (150 MHz, CDCl₃): δ 20.68, 20.86, 20.87, 54.52, 55.45, 61.69, 68.43, 68.90, 70.77, 71.81, 73.06, 74.89, 76.42, 77.70, 78.79, 82.03, 97.25, 123.94–146.65 (aromatic carbons), 169.74, 170.36, 170.88 (CH₃CO); HRMS: [M+Na]⁺ C₅₄H₄₉N₃NaO₁₇S calcd 1066.2680, obsd 1066.2742.

3.10. *p*-Tolyl 3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido-β-D-glucopyranosyl-(1→4)-2-acetamido-6-*O*-benzyl-2-*N*-3-*O*-carbonyl-2-deoxy-1-thio-α-D-glucopyranoside (14)

Compound **14** was synthesized from donor **8** and acceptor **2α** in 39% yield following the general procedure of single step glycosylation together with recovered acceptor **2α** (50%). [α]_D +98.8 (*c*, 0.0093 g/mL, CH₂Cl₂); ¹H NMR (600 MHz, CDCl₃): δ 1.86 (s, 3H), 2.04 (s, 3H), 2.14 (s, 3H), 2.29 (s, 3H), 2.52 (s, 3H), 3.39 (m, 1H), 3.49 (dd, *J* = 4.8 Hz, 13.8 Hz, 1H), 3.88–4.25 (m, 8H), 4.33–4.37 (m, 2H), 4.48 (dd, *J* = 14.4 Hz, 18.6 Hz, 1H), 5.21 (t, *J* = 15.0 Hz, 1H), 5.56 (d, *J* = 13.2 Hz, 1H), 5.78 (dd, *J* = 13.2 Hz, 15.6 Hz, 1H), 6.05 (d, *J* = 6.6 Hz, 1H), 6.98–7.88 (m, 13H); ¹³C NMR (150 MHz, CDCl₃): δ 20.69, 20.89, 21.04, 21.35, 24.12, 54.94, 60.05, 61.93, 67.82, 68.80, 71.03, 72.26, 72.46, 73.20, 75.40, 77.45, 86.45, 97.75, 123.86–138.61 (aromatic carbons), 152.98, 169.75, 170.41, 171.12, 171.41; HRMS: [M+Na]⁺ C₄₃H₄₄N₂NaO₁₅S calcd 883.2360, obsd 883.2333.

3.11. Benzyl 3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido-β-D-glucopyranosyl-(1→4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido-β-D-glucopyranosyl-(1→4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido-β-D-glucopyranoside (16)

After the donor **8** (60 mg, 0.11 mmol) and activated molecular sieve MS—4 Å (500 mg) were stirred for 30 min at room temperature in the mixed solvents of CH₂Cl₂ and Et₂O (1:1) (6 mL), the solution was cooled to −70 °C followed by sequential additions of AgOTf (86 mg, 0.335 mmol) in Et₂O (1.5 mL) and *p*-TolSCl (17.3 μL, 0.11 mmol). The mixture was vigorously stirred for 15 min and a solution of acceptor **3** (59 mg, 0.099 mmol) in CH₂Cl₂ (0.5 mL) was added. The mixture was stirred for 5 min, then the flask was warmed to −10 °C in 15 min. Then the mixture was cooled down to −70 °C again followed by sequential additions of AgOTf (76 mg, 0.297 mmol) in Et₂O (1.5 mL) and *p*-TolSCl (15.5 μL, 0.099 mmol). After 15 min, a solution of acceptor **3** (53 mg, 0.089 mmol) in CH₂Cl₂ (0.5 mL) was injected through a syringe. The mixture was stirred for 5 min and warmed to −10 °C in 15 min by switching the cold bath. A solution of acceptor **15** (64 mg, 0.11 mmol) in CH₂Cl₂ (0.5 mL) was added to the reaction mixture, and the resulting mixture was cooled down to −70 °C followed by sequential additions of AgOTf (68 mg, 0.089 mmol) in Et₂O (1.5 mL) and *p*-TolSCl (14 μL, 0.089 mmol). The reaction mixture was stirred for 1.5 h from −70 to 10 °C and then was diluted with CH₂Cl₂ (30 mL) followed by filtration and extraction. The collected organic phase was dried over Na₂SO₄, and the residue was applied to flash column

chromatography with a mixed solvent system of CH_2Cl_2 , hexanes, and EtOAc to give the desired tetrasaccharide **16** (55 mg) in 32% yield as colorless foam. Its structure was confirmed by ^1H NMR, ^{13}C NMR, ^1H – ^1H COSY experiments, and HRMS. $[\alpha]_{\text{D}} -20.0$ (*c*, 0.038 g/mL, CH_2Cl_2); ^1H NMR (600 MHz, CDCl_3): (a, b, c, and d denote the four monosaccharide units in the sequence of its anomeric proton appearance in the ^1H NMR spectrum from the most downfield to the most upfield shifted. Monosaccharide unit a is the one at the non-reducing end) δ 1.82 (s, 3H, CH_3CO), 1.88 (s, 3H, CH_3CO), 1.96 (s, 3H, CH_3CO), 2.76 (broad d, $J = 10.2$ Hz, 1H, c/d- H_5), 2.89 (broad d, $J = 10.2$ Hz, 1H, b- H_5), 3.02 (dd, $J = 3.0$, 10.2 Hz, 1H, c/d- H_6), 3.15 (dd, $J = 2.4$, 10.2 Hz, 1H, b- H_6), 3.18 (dd, $J = 3.0$, 10.2 Hz, 1H, c/d- H_5), 3.27 (dd, $J = 3.6$, 10.2 Hz, 1H, c/d- H_6), 3.29 (d, $J = 10.2$ Hz, 1H, c/d- H_6), 3.37–3.42 (m, 2H, a- H_5 , b- H_6), 3.44 (broad d, $J = 10.2$ Hz, 1H, c/d- H_6), 3.88 (dd, $J = 1.8$, 12.0 Hz, 1H, a- H_6), 3.97–4.10 (m, 7H, b- H_2 , c- H_2 , c- H_3 , c- H_4 , d- H_2 , d- H_3 , d- H_4), 4.14 (dd, $J = 9.0$, 10.8 Hz, 1H, a- H_3), 4.16 (dd, $J = 8.4$, 12.0 Hz, 1H, a- H_6), 4.24 (d, $J = 9.0$ Hz, 1H, a- H_4), 4.27–4.44 (m, 10H, 9Ph CH_2 , a- H_2), 4.52 (d, $J = 11.4$ Hz, 1H, Ph CH_2), 4.62 (d, $J = 12.6$ Hz, 1H, Ph CH_2), 4.67 (d, $J = 12.6$ Hz, 1H, Ph CH_2), 4.81 (d, $J = 12.6$ Hz, 1H, Ph CH_2), 4.84 (d, 1H, $J = 12.6$ Hz, Ph CH_2), 4.85 (d, $J = 8.4$ Hz, 1H, d- H_1), 5.01 (m, 1H, c- H_1), 5.07 (d, 1H, $J = 9.0$ Hz, 1H, b- H_1), 5.10 (t, $J = 9.0$ Hz, a- H_4), 5.46 (d, $J = 8.4$ Hz, 1H, a- H_1), 5.75 (dd, $J = 9.0$, 10.8 Hz, 1H, a- H_3), 6.63–7.91 (m, 51 H); ^{13}C NMR (150 MHz, CDCl_3): δ 20.68, 20.82, 20.85, 55.49, 55.86, 56.79, 56.82, 61.57, 67.19, 67.27, 68.28, 68.87, 70.64, 70.91, 71.53, 72.35, 72.50, 72.72, 74.18, 74.36, 74.42, 74.60, 74.63, 74.73, 75.47, 75.57, 76.08, 76.54, 96.81, 96.81, 96.91, 97.21, 123.26–139.00 (aromatic carbons), 167.78–168.41 (phthalimido carbonyl carbons), 169.74, 170.36, 170.91 (CH_3CO); HRMS: $[\text{M}+\text{Na}]^+ \text{C}_{111}\text{H}_{102}\text{N}_4\text{NaO}_{28}$ calcd 1961.6578, obsd 1961.6393.

3.12. *p*-Tolyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (**17**)

To a mixture of compound **5** (1 g, 2 mmol), tetrabutylammonium iodide (0.15 g, 0.4 mmol), and freshly activated molecular sieve MS-4 Å (300 mg) in anhydrous DMF (10 mL) was added 95% NaH (80 mg, 3.2 mmol) at 0 °C. The mixture was stirred for 30 min and benzyl bromide (290 μL , 0.42 g, 2.4 mmol) was added via a syringe. After 90 min, the reaction was quenched with acetic acid and filtered. Et_2O (40 mL) was added to the filtrate, which was extracted with saturated aqueous solutions of NH_4Cl (20 mL) and NaHCO_3 (20 mL). The organic layer was separated and dried over Na_2SO_4 . Column chromatography on silica gel (3:1, hexanes–EtOAc) afforded the product **17**

as white foam (1.07 g, 90%). Comparison of the NMR data with those reported in the literature²¹ confirmed the identity of **17**.

3.13. *p*-Tolyl 4,6-*O*-benzylidene-3-*O*-*tert*-butyldimethylsilyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (**18**)

To a mixture of compound **5** (1 g, 2 mmol) and molecular sieve MS-4 Å (300 mg) in anhydrous CH_2Cl_2 (10 mL) were added 2,6-lutidine (460 μL , 4 mmol) and *tert*-butyldimethylsilyl triflate (550 μL , 2.4 mmol) at 0 °C. The mixture was stirred for 4 h. The reaction mixture was filtered and the filtrate was extracted with a satd aqueous solution of NH_4Cl (20 mL). The organic layer was separated and dried over Na_2SO_4 . Column chromatography on silica gel (15% EtOAc in hexanes) afforded the product **18** as white foam (1.04 g, 85%). ^1H NMR (600 MHz, CDCl_3): δ -0.32 (s, 3H), -0.16 (s, 3H), 0.56 (s, 9H), 2.28 (s, 3H), 3.55 (t, $J = 9.6$ Hz, 1H), 3.67 (dt, $J = 4.8$, 9.6 Hz, 1H), 3.80 (t, $J = 10.2$ Hz, 1H), 4.29 (t, $J = 10.2$ Hz, 1H), 4.37 (dd, $J = 4.8$, 10.2 Hz, 1H), 4.61 (t, $J = 9.6$ Hz, 1H), 5.51 (s, 1H), 5.57 (d, $J = 10.2$ Hz, 1H), 7.01–7.04 (m, 2H), 7.20–7.35 (m, 7H), 7.75–7.95 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3): δ -5.11, -3.93, 17.96, 21.39, 25.63, 57.02, 68.92, 70.79, 82.70, 84.86, 102.24, 123.47–138.47 (aromatic carbons), 167.64, 168.61 (carbonyl groups of phthalimido); ESI-MS $[\text{M}+\text{Na}]^+ \text{C}_{34}\text{H}_{39}\text{NNaO}_6\text{SSi}$ calcd 640.8, obsd 640.3.

3.14. *p*-Tolyl 4,6-di-*O*-benzyl-3-*O*-*tert*-butyldimethylsilyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (**19**)

To a solution of compound **18** (0.43 g, 0.7 mmol) and 1 M borane THF (7.2 mL) in anhydrous CH_2Cl_2 (10 mL) was added trimethylsilyl triflate (100 μL , 0.5 mmol) at -20 °C. After 2 h, trimethylsilyl triflate (200 μL , 1 mmol) was added again. The mixture was stirred for 2 h more. The reaction was quenched by the addition of Et_3N , and MeOH was added until no gas was formed. All solvents were evaporated and the desired primary alcohol (0.39 g) was purified through column chromatography on silica gel (25% EtOAc in hexanes). To a mixture of the primary alcohol (0.3 g, 0.5 mmol), tetrabutylammonium iodide (40 mg, 0.1 mmol) and freshly activated molecular sieve MS-4 Å (100 mg) in anhydrous DMF (5 mL) was added 95% NaH (20 mg, 0.8 mmol) at 0 °C. The mixture was stirred for 30 min and benzyl bromide (180 μL , 0.26 g, 1.5 mmol) was added via a syringe. After 90 min, the reaction was quenched by the addition of acetic acid and filtered. Et_2O (10 mL) was added to the filtrate, which was extracted with saturated aqueous solutions of NH_4Cl (10 mL) and NaHCO_3 (10 mL). The organic layer was separated and dried over Na_2SO_4 . Column

chromatography on silica gel (4:1, hexanes–EtOAc) afforded the product **19** (0.32 g) as white foam in overall 80% yield for the two steps. ^1H NMR (600 MHz, CDCl_3): δ –0.44 (s, 3H), –0.08 (s, 3H), 0.72 (s, 9H), 2.27 (s, 3H), 3.58 (t, $J = 9.6$ Hz, 1H), 3.64–3.69 (m, 1H), 3.72–3.80 (m, 2H), 4.27 (t, $J = 10.2$ Hz, 1H), 4.47 (t, $J = 9.0$ Hz, 1H), 4.53 (d, $J = 12.0$ Hz, 1H), 4.59 (d, $J = 12.0$ Hz, 1H), 4.65 (d, $J = 12.0$ Hz, 1H), 4.78 (d, $J = 12.0$ Hz, 1H), 5.52 (d, $J = 10.2$ Hz, 1H), 7.01–7.04 (m, 2H), 7.20–7.35 (m, 12H), 7.75–7.95 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3): δ –4.46, –3.94, 17.84, 21.35, 25.87, 56.99, 69.18, 73.64, 73.71, 74.70, 79.67, 80.01, 83.69, 123.39, 123.78, 127.25, 127.59, 127.74, 127.87, 127.90, 128.45–128.53, 129.75, 133.34, 134.38, 138.51, 138.52; ESI-MS $[\text{M}+\text{Na}]^+$, $\text{C}_{41}\text{H}_{47}\text{NNaO}_6\text{SSi}$ calcd 732.9, obsd 732.5.

3.15. *p*-Tolyl 3-*O*-benzyl-4,6-benzylidene-2-deoxy-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (20)

Compound **20** was synthesized from donor **17** and acceptor **3** in 61% yield following the general procedure of single step glycosylation. $[\alpha]_{\text{D}} +17.0$ (c, 0.024 g/mL, CH_2Cl_2); ^1H NMR (600 MHz, CDCl_3): δ 2.21 (s, 3H), 3.33–3.42 (m, 3H), 3.48–3.52 (m, 2H), 3.70 (t, $J = 9.0$ Hz, 1H), 4.10–4.22 (m, 5H), 4.37 (d, $J = 20$ Hz, 1H), 4.42 (dd, 1H, $J = 9.0, 10.2$ Hz, 1H), 4.43–4.48 (m, 3H), 4.77 (dd, $J = 4.8, 12.6$ Hz, 2H), 5.27 (d, $J = 10.2$ Hz, 1H), 5.35 (d, $J = 8.4$ Hz, 1H), 5.50 (s, 1H), 6.86–7.88 (m, 32 H); ^{13}C NMR (150 MHz, CDCl_3): δ 21.32, 54.91, 56.72, 65.95, 68.35, 68.96, 72.92, 74.32, 74.67, 74.80, 76.32, 78.00, 78.93, 83.35, 83.39, 97.95, 101.41, 126.29–138.58 (aromatic carbons); HRMS: $[\text{M}+\text{Na}]^+$ $\text{C}_{63}\text{H}_{56}\text{N}_2\text{NaO}_{12}\text{S}$ calcd 1087.3452, obsd 1087.3494.

3.16. *p*-Tolyl 4,6-*O*-benzylidene-3-*O*-tert-butyldimethylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (21)

Compound **21** was synthesized from donor **18** and acceptor **3** in 74% yield following the general procedure of single step glycosylation. $[\alpha]_{\text{D}} -4.4$ (c, 0.011 g/mL, CH_2Cl_2); ^1H NMR (600 MHz, CDCl_3): δ –0.28 (s, 3H), –0.14 (s, 3H), 0.56 (s, 9H), 2.21 (s, 3H), 3.32–3.38 (m, 2H), 3.42 (dd, $J = 7.2, 10.8$ Hz, 1H), 3.46 (t, $J = 9.0$ Hz, 1H), 3.52 (t, $J = 10.2$ Hz, 1H), 3.54 (t, $J = 10.2$ Hz, 1H), 4.08–4.24 (m, 5H), 4.46 (d, $J = 12.0$ Hz, 1H), 4.49 (d, $J = 12.0$ Hz, 1H), 4.53 (d, $J = 12.0$ Hz, 1H), 4.61 (t, $J = 9.6$ Hz, 1H), 4.77 (d, $J = 12.0$ Hz, 1H), 5.27 (d, $J = 10.2$ Hz, 1H), 5.34 (d, $J = 8.4$ Hz, 1H), 5.41 (s, 1H), 6.84–6.92 (m, 5H), 7.00–7.03 (m, 2H), 7.16–7.20 (m, 2H), 7.28–7.39 (m, 8H), 7.42–7.46 (m, 2H), 7.54–7.58 (m, 1H), 7.62–7.69 (m, 2H),

7.71–7.79 (m, 3H), 7.84–7.88 (m, 1H), 7.92–7.94 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3): δ –5.46, –4.12, 17.67, 21.07, 25.36, 54.67, 58.29, 65.92, 68.17, 68.75, 69.30, 72.71, 74.56, 76.03, 77.55, 78.74, 82.78, 83.10, 97.60, 101.85, 123.07, 123.19, 123.41, 123.78, 126.34, 127.06, 127.33, 127.50, 127.74, 127.91, 128.05, 128.11, 128.26, 129.03, 129.41, 131.59, 131.65, 131.66, 133.48, 133.61, 133.78, 134.28, 134.33, 137.15, 138.04, 138.33, 138.37, 167.21, 167.63, 167.86, 168.64; HRMS: $[\text{M}+\text{Na}]^+$ $\text{C}_{62}\text{H}_{64}\text{N}_2\text{NaO}_{12}\text{SSi}$ calcd 1111.3925, obsd 1111.3877.

3.17. *p*-Tolyl 4,6-di-*O*-benzyl-3-*O*-tert-butyldimethylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (22)

Compound **22** was synthesized from donor **19** and acceptor **3** in 50% yield following the general procedure of single step glycosylation together with disaccharide **23** and glycal **24** (10%). $[\alpha]_{\text{D}} +20.1$ (c, 0.035 g/mL, CH_2Cl_2); ^1H NMR (600 MHz, CDCl_3): δ –0.40 (s, 3H), –0.07 (s, 3H), 0.70 (s, 9H), 2.20 (s, 3H), 3.32–3.36 (m, 2H), 3.48 (dd, $J = 3.6$ Hz, 10.8 Hz, 1H), 3.58–3.70 (m, 4H), 4.10–4.24 (m, 5H), 4.47–4.56 (m, 6H), 4.69 (d, $J = 12.0$ Hz, 1H), 4.75 (d, $J = 11.4$ Hz, 1H), 4.88 (d, $J = 12.6$ Hz, 1H), 5.26 (d, $J = 10.3$ Hz, 1H), 4.30 (d, $J = 10.2$ Hz, 1H), 6.71–7.89 (m, 32 H); ^{13}C NMR (150 MHz, CDCl_3): δ –4.82, –3.88, 17.89, 21.32, 25.87, 55.05, 58.59, 68.10, 68.62, 72.21, 72.92, 73.33, 74.37, 74.86, 75.31, 75.46, 77.78, 79.09, 80.11, 83.58, 96.77, 123.24–138.79 (aromatic carbons), 167.51, 167.92, 168.01, 169.25 (carbonyl groups of phthalimido); HRMS: $[\text{M}+\text{Na}]^+$ $\text{C}_{69}\text{H}_{72}\text{N}_2\text{NaO}_{12}\text{SSi}$ calcd 1203.4473, obsd 1203.4505.

3.18. 4,6-di-*O*-benzyl-3-*O*-tert-butyldimethylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 1)-4,6-di-*O*-benzyl-3-*O*-tert-butyldimethylsilyl-2-deoxy-2-phthalimido-D-glucopyranoside (23)

$[\alpha]_{\text{D}} = 0$ (c, 0.013 g/mL, CH_2Cl_2); ^1H NMR (600 MHz, CDCl_3): δ –0.48 (s, 6H), –0.13 (s, 6H), 0.66 (s, 18H), 3.37 (m, 2H), 3.47 (m, 4H), 3.65 (m, 2H), 4.05 (d, $J = 18$ Hz, 2H), 4.07 (dd, $J = 13.2$ Hz, 15.6 Hz, 4H), 4.20 (d, $J = 18$ Hz, 2H), 4.43 (m, 2H), 4.54 (d, $J = 17.4$ Hz, 2H), 4.65 (d, $J = 7.4$ Hz, 2H), 5.41 (d, $J = 13.2$ Hz, 2H), 7.13–7.55 (m, 28H); ^{13}C NMR (150 MHz, CDCl_3): δ –4.55, –4.04, 17.78, 25.82, 57.48, 69.12, 72.20, 73.72, 74.41, 75.47, 79.62, 96.88, 127.17–138.69 (aromatic carbons); HRMS: $[\text{M}+\text{Na}]^+$ $\text{C}_{68}\text{H}_{80}\text{N}_2\text{NaO}_{13}\text{Si}_2$ calcd 1211.5096, obsd 1211.5143. Compound **24** was isolated as a mixture with compound **23**, which was characterized by ESI-MS. ESI-MS: $[\text{M}+\text{Na}]^+$ $\text{C}_{34}\text{H}_{39}\text{NNaO}_6\text{Si}$ calcd 608.2, obsd 608.3.

3.19. Methyl 4,6-*O*-benzylidene-3-*O*-*tert*-butyldimethylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 4)-6-*O*-benzyl-2-deoxy-3-*O*-*p*-methoxybenzyl-2-phthalimido- β -D-glucopyranosyl-(1 \rightarrow 4)-3,6-di-*O*-benzyl-2-deoxy-2-phthalimido- β -D-glucopyranoside (27)

Compound **27** was synthesized in 45% overall yield using building blocks **18**, **3**, **25**, and **26** in a similar manner as in the assembly of tetrasaccharide **16** with the exception that pure Et₂O was used to dissolve donor **18**, and TTBP (1 equiv) was added together with each acceptor. For reaction time, see Scheme 4. The desired product was purified by silica gel flash chromatography using the solvent systems of hexanes–EtOAc (1:1) and hexanes–CH₂Cl₂–EtOAc (5:5:3). Its structure was confirmed by ¹H NMR, ¹³C NMR, ¹H–¹H COSY, ¹H–¹³C HMQC experiments, and HRMS. [α]_D –31.4 (c, 0.013 g/mL, CH₂Cl₂); ¹H NMR (600 MHz, CDCl₃): (a, b, c, and d denote the four monosaccharide units in the sequence of its anomeric proton appearance in the ¹H NMR spectrum from the most downfield to the most upfield shifted) δ –0.31 (s, 3H, SiCH₃), –0.17 (s, 3H, SiCH₃), 0.54 (s, 9H, SiC(CH₃)₃), 2.76 (broad d, *J* = 9.0 Hz, 1H, c-H₅), 2.90 (broad d, *J* = 9.0 Hz, 1H, b-H₅), 3.01 (dd, *J* = 3.0, 10.8 Hz, 1H, c-H₆), 3.12 (dd, *J* = 2.4, 10.8 Hz, 1H, b-H₆), 3.18–3.24 (m, 1H, d-H₄), 3.21 (s, 3H, OCH₃), 3.24–3.32 (m, 3H, c-H₆, d-H₅, d-H₆), 3.37–3.47 (m, 4H, a-H₄, a-H₆, b-H₆, d-H₆), 3.54 (s, 3H, ArOCH₃), 3.96–4.03 (m, 4H, c-H₂, c-H₃, d-H₂, d-H₃), 4.04–4.12 (m, 3H, a-H₆, b-H₂, c-H₄), 4.13–4.21 (m, 3H, a-H₂, a-H₅, b-H₃), 4.23 (t, *J* = 8.4 Hz, 1H, b-H₄), 4.29 (d, *J* = 12.0 Hz, 1H, ArCH₂), 4.32–4.42 (m, 6H, 6ArCH₂), 4.47 (d, *J* = 12.0 Hz, 1H, ArCH₂), 4.50 (d, *J* = 12.0 Hz, 1H, ArCH₂), 4.59 (t, *J* = 9.0 Hz, 1H, a-H₃), 4.70 (d, *J* = 12.6 Hz, 1H, ArCH₂), 4.71 (d, *J* = 12.6 Hz, 1H, ArCH₂), 4.77 (d, *J* = 7.8 Hz, 1H, d-H₁), 4.82 (d, *J* = 12.0 Hz, 1H, ArCH₂), 5.01–5.04 (m, 1H, c-H₁), 5.07 (d, *J* = 7.8 Hz, 1H, b-H₁), 5.33 (d, *J* = 8.4 Hz, 1H, a-H₁), 5.38 (s, 1H, PhCH_{benzylidene}), 6.18–6.21 (m, 2H), 6.62–7.96 (m, 48H); ¹³C NMR (100 MHz, CDCl₃): –5.18 (SiCH₃), –3.85 (SiCH₃), 17.94 (SiC(CH₃)₃), 25.65 (SiC(CH₃)₃), 54.88 (ArOCH₃), 55.76, 56.60 (OCH₃), 56.85 (b-C₂), 56.90, 58.58 (a-C₂), 66.05, 67.33, 68.30, 68.97, 69.61, 72.45, 72.61, 72.74, 74.36, 74.46, 74.52, 74.72, 75.51, 75.80, 76.26, 76.58, 83.04 (a-C₄), 96.75 (b-C₁), 96.95 (c-C₁), 97.76 (a-C₁), 99.18 (d-C₁), 102.09 (PhCH_{benzylidene}), 113.25, 123.05, 123.35, 123.52, 123.79, 123.95, 126.59, 126.88, 127.16, 127.23, 127.48, 127.53, 127.56, 127.59, 127.87, 127.99, 128.11, 128.14, 128.22, 128.27, 128.33, 128.52, 129.25, 129.88, 131.33, 131.77, 131.99, 133.71, 133.73, 133.82, 134.02, 134.19, 134.53, 137.42, 138.53, 138.60, 138.63, 138.88, 138.97, 158.50, 167.82, 167.84, 168.27, 168.38, 168.75 (carbonyl groups of phthalimido); ESI-MS C₁₁₃H₁₁₂N₄NaO₂₆Si [M+Na]⁺ calcd 1991.7, found

1991.6; HRMS: [M+Na]⁺ C₁₁₃H₁₁₂N₄NaO₂₆Si calcd 1991.7231, obsd 1991.7139. Compounds **28**, **29**, and **30** were characterized by ESI-MS. Compound **28**, C₅₅H₅₆N₂NaO₁₂Si [M+Na]⁺ calcd 987.4, obsd 987.6. Compound **29**, C₈₄H₈₃N₃NaO₁₉Si [M+Na]⁺ calcd 1488.5, obsd 1488.7. Compound **30**, C₈₄H₈₅N₃NaO₂₀Si [M+Na]⁺ calcd 1506.5, obsd 1506.7.

Acknowledgment

We are grateful for the financial support from the University of Toledo, the National Institutes of Health (R01 GM 72667), and the Pardee Foundation.

Supplementary data

¹H and ¹³C NMR spectra of compounds **2a**, **6**, **9**, **11**, **13**, **14**, **16**, **20**, **21**, **22**, **23**, and **27**. Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.carres.2006.01.007](https://doi.org/10.1016/j.carres.2006.01.007).

References

- (a) Codee, J. D. C.; Litjens, R. E. J. N.; Van den Bos, L. J.; Overkleeft, H. S.; Van der Marel, G. A. *Chem. Soc. Rev.* **2005**, *34*, 769–782; (b) *Carbohydrates in Chemistry and Biology*; Wiley-VCH: Weinheim, 2000; Vols. 1–4; (c) Davis, B. G. *J. Chem. Soc., Perkin Trans. 1* **2000**, 2137–2160.
- Yu, B.; Yang, Z.; Cao, H. *Curr. Org. Chem.* **2005**, *9*, 179–194.
- Koeller, K. M.; Wong, C.-H. *Chem. Rev.* **2000**, *100*, 4465–4493.
- Huang, L.; Wang, Z.; Huang, X. *Chem. Commun.* **2004**, 1960–1961.
- (a) Pornsuriyasak, P.; Gangadharmath, U. B.; Rath, N. P.; Demchenko, A. V. *Org. Lett.* **2004**, *6*, 4515–4518; (b) Lahmann, M.; Oscarson, S. *Org. Lett.* **2000**, *2*, 3881–3882; (c) Geurtsen, R.; Holmes, D. S.; Boons, G.-J. *J. Org. Chem.* **1997**, *62*, 8145–8154.
- (a) Lopez, J. C.; Agocs, A.; Uriel, C.; Gomez, A. M.; Fraser-Reid, B. *Chem. Commun.* **2005**, 5088–5090; (b) Fraser-Reid, B.; Lopez, J. C.; Gomez, A. M.; Uriel, C. *Eur. J. Org. Chem.* **2004**, 1387–1395.
- Huang, X.; Huang, L.; Wang, H.; Ye, X.-S. *Angew. Chem., Int. Ed.* **2004**, *43*, 5221–5224.
- (a) Semenuk, T.; Krist, P.; Pavlicek, J.; Bezouska, K.; Kuzma, M.; Novak, P.; Kren, V. *Glycoconjugate J.* **2002**, *18*, 817–826; (b) Shibata, Y. *Rec. Res. Dev. Immun.* **1999**, *1*, 551–558; (c) Bezouska, K.; Sklenar, J.; Dvorakova, J.; Havlicek, V.; Pospisil, M.; Thiem, J.; Kren, V. *Biochem. Biophys. Res. Commun.* **1997**, *238*, 149–153.
- (a) Gorbach, V. I.; Kraiskova, I. N.; Lukyanov, P. A.; Loenko, Y. N.; Soloveva, T. F.; Ovodov, Y. S.; Deev, V. V.; Pimenov, A. A. *Carbohydr. Res.* **1994**, *260*, 73; (b) Suzuki, K.; Mikami, T.; Okawa, Y.; Tokoro, A.; Suzuki, S.; Suzuki, M. *Carbohydr. Res.* **1986**, *151*, 403–408.
- Vongchan, P.; Sajomsang, W.; Subyen, D.; Kongtawelert, P. *Adv. Chitin Sci.* **2002**, *5*, 650–655.

11. (a) Nishimura, S.-I.; Kai, H.; Shinada, K.; Yoshida, T.; Tokura, S.; Kurita, K.; Nakashima, H.; Yamamoto, N.; Uryu, T. *Carbohydr. Res.* **1998**, *306*, 427–433; (b) Katsuraya, K.; Ishikawa, S.; Ichikawa, T.; Goto, K.; Uryu, T. *Seisan Kenkyu* **1997**, *49*, 170–173.
12. (a) Yamago, S.; Yamada, T.; Maruyama, T.; Yoshida, J.-I. *Angew. Chem., Int. Ed.* **2004**, *43*, 2145–2148; (b) Aly, M. R. E.; Ibrahim, E. I.; El Ashry, E. S. H.; Schmidt, R. R. *Carbohydr. Res.* **2001**, *331*, 129–142; (c) Hinou, H.; Umino, A.; Matsuoka, K.; Terunuma, D.; Takahashi, S.; Esumi, Y.; Kuzuhara, H. *Bull. Chem. Soc. Jpn.* **2000**, *73*, 163–171; (d) Kanie, O.; Ito, Y.; Ogawa, T. *J. Am. Chem. Soc.* **1994**, *116*, 12073–12074; (e) Tailler, D.; Jacquinet, J.-C.; Beau, J.-M. *J. Chem. Soc., Chem. Commun.* **1994**, 1827–1828; (f) Ikeshita, S.; Sakamoto, A.; Nakahara, Y.; Nakahara, Y.; Ogawa, T. *Tetrahedron Lett.* **1994**, *35*, 3123–3126; (g) Wang, L. X.; Li, C.; Wang, Q. W.; Hui, Y. Z. *Tetrahedron Lett.* **1993**, *34*, 7763–7766; (h) Kuyama, H.; Nakahara, Y.; Nukada, T.; Ito, Y.; Nakahara, Y.; Ogawa, T. *Carbohydr. Res.* **1993**, *243*, C1–C7; (i) Nicolaou, K. C.; Bockovich, N. J.; Carcanague, D. R.; Hummel, C. W.; Even, L. F. *J. Am. Chem. Soc.* **1992**, *114*, 8701–8702.
13. Schwartz, D. A.; Lee, H.-H.; Carver, J. P.; Krepinsky, J. J. *Can. J. Chem.* **1985**, *63*, 1073–1079.
14. Crich, D.; Dudkin, V. *J. Am. Chem. Soc.* **2001**, *123*, 6819–6825.
15. Xue, J.; Khaja, S. D.; Locke, R. D.; Matta, K. L. *Synlett* **2004**, 861–865.
16. Crich, D.; Vinod, A. U. *J. Org. Chem.* **2005**, *70*, 1291–1296.
17. Crich, D.; Vinod, A. U. *Org. Lett.* **2003**, *5*, 1297–1300.
18. (a) Boysen, M.; Gemma, E.; Lahmann, M.; Oscarson, S. *Chem. Commun.* **2005**, 3044–3046; (b) Wei, P.; Kerns, R. J. *J. Org. Chem.* **2005**, *70*, 4195–4198.
19. (a) Tanaka, H.; Adachi, M.; Takahashi, S. *Tetrahedron Lett.* **2004**, *45*, 1433–1436; (b) Orgueira, H. A.; Bartolozzi, A.; Schell, P.; Litjens, R. E. J. N.; Palmacci, E. R.; Seeberger, P. H. *Chem. Eur. J.* **2003**, *9*, 140–169; (c) Paulsen, H.; Stenzel, W. *Angew. Chem., Int. Ed.* **1975**, *14*, 558–559.
20. Zhang, Z.; Ollmann, I. R.; Ye, X.-S.; Wischnat, R.; Baasov, T.; Wong, C.-H. *J. Am. Chem. Soc.* **1999**, *121*, 734–753.
21. Wang, Y.; Huang, X.; Zhang, L.-H.; Ye, X.-S. *Org. Lett.* **2004**, *6*, 4415–4417.
22. Mong, T. K. K.; Huang, C.-Y.; Wong, C.-H. *J. Org. Chem.* **2003**, *68*, 2135–2142.
23. Barrett, A. G. M.; Dhanak, D.; Graboski, G. G.; Taylor, S. J. *Org. Synth. Coll.* **1993**, *8*, 550.
24. (a) Crich, D.; Sun, S. *Tetrahedron* **1998**, *54*, 8321–8348; (b) Martichonok, V.; Whitesides, G. M. *J. Org. Chem.* **1996**, *61*, 1702–1706.
25. (a) Zhu, T.; Boons, G.-J. *Carbohydr. Res.* **2000**, *329*, 709–715; (b) Belot, F.; Jacquinet, J.-C. *Carbohydr. Res.* **1996**, *290*, 79–86.
26. Hernandez-Torres, J. M.; Liew, S.-T.; Achkar, J.; Wei, A. *Synthesis* **2002**, 487–490.
27. Kameyama, A.; Ehara, T.; Yamada, Y.; Ishida, H.; Kiso, M.; Hasegawa, A. *J. Carbohydr. Chem.* **1995**, *14*, 507–523.
28. Crich, D.; Smith, M.; Yao, Q.; Picione, J. *Synthesis* **2001**, 323–326.
29. Chowdhury, A. R.; Siriwardena, A.; Boons, G. J. *Tetrahedron Lett.* **2002**, *43*, 7805–7807.
30. (a) Seeberger, P. H. *Chem. Commun.* **2003**, 1115–1121; (b) Plante, O. J.; Palmacci, E. R.; Seeberger, P. H. *Science* **2001**, *291*, 1523–1527; (c) Seeberger, P. H.; Haase, W.-C. *Chem. Rev.* **2000**, *100*, 4349–4393.