

*Ammotragus* is derived from the ancestral caprine form by one centric fusion. Furthermore, the identity between *Ammotragus* and *Ovis vignei* karyotypes supports GEIST'S<sup>3</sup> view that an *Ammotragus*-like form was directly ancestral to *Ovis*. Finally, the chromosomal homologies in the series of 1 to 3 biarmed autosomal pairs in *Ovis* lend further strength to the argument that *Ovis vignei* ( $2n = 58$ ) is descended from the ancestral *Ovis* population, and that  $2n = 54$  *Ovis* are the most recently evolved of wild sheep<sup>9, 11</sup>.

The fossil record indicates that *Ovis* and *Capra* have been separated for more than a million years<sup>4, 5</sup>. During that time, their chromosomes exhibited structural stability, evolutionary change consisting merely of whole-arm rearrangements. Morphological<sup>2</sup> and behavioral<sup>3</sup> differences between sheep and goats are thus due to the accumulation of gene differences. The cytogenetic evidence suggests there are two main lineages among surviving caprines. Starting from a hypothetical rupicaprine-type ancestor with a primitive  $2n = 60$ ,  $FN = 60$  karyotype, one lineage evolved through an intermediate,




Fig. 4. Karyotype of a male *Ovis vignei* ( $2n = 58$ ). The biarmed autosomes conform to the pattern of pair I.

aoudad-like form to the true sheep, with reductions in diploid number. In the true goat (*Capra*) lineage, morphological differentiation proceeded while the karyotype remained conservative; in contrast, the thar (*Hemitragus*) has remained morphologically close to the hypothetical rupicaprine ancestor<sup>3</sup> while its chromosome number has been reduced ( $2n = 48$ )<sup>7</sup>. The 5th member of the tribe Caprini, the bharal (*Pseudois nayaur*) has a reduced chromosome number ( $2n = 54$ )<sup>13</sup> but morphologically exhibits convergence toward true sheep<sup>2, 3</sup>. G-bands in *Hemitragus* and *Pseudois* must be studied to confirm their chromosomal relationship to other caprines and it will be of interest to determine whether the biarmed chromosomes of *Pseudois* are homologous to those of  $2n = 54$  *Ovis*. If so, it would support the concept of convergent evolution at the chromosomal level<sup>9</sup>.

**ВЫВОДЫ.** Хромосомы и Г-полосы диких баранов, Африканских гривистых баранов, и безоаровых козлов изучали. *Capra aegagrus* из Ирана показало  $2n = 60$ , и кариотип содержит 29 пары акроцентрических аутосом. Шесть из этих аутосомов одинаковый с трёх пар двуплечных аутосом у диких баранов и домашних овец ( $2n = 54$ ). *Ammotragus levvia* и *Ovis vignei* ( $2n = 58$ ) имели кариотипы и Г-полосы пар двуплечных аутосом неразличимый.

C. F. NADLER, R. S. HOFFMANN and A. WOOLF<sup>14</sup>

Department of Medicine, Northwestern University Medical School, 303 East Chicago Avenue, Chicago (Illinois 60611, USA);  
Museum of Natural History, University of Kansas, Lawrence (Kansas 66045, USA); and  
Rachelwood Wildlife Research Preserve, New Florence (Pennsylvania 15944, USA), 15 January 1974.

<sup>13</sup> W. L. HARD, Mammal. Chromos. Newsl. 10, 228 (1969).

<sup>14</sup> Supported by National Science Foundation Grants No. GB 32114X and 29131X, the Sprague Foundation, and a grant from the Trustees of the Rachelwood Wildlife Research Preserve. We thank Dr. T. C. Hsu for valuable suggestions, Mr. ARTHUR POPHAM for obtaining the specimens from Iran, and the Highland Park Zoo, Pittsburgh, Pennsylvania, for permission to biopsy their specimen.

### Effects of Abscisic Acid on Photosynthesis of Protoplasts from *Petunia hybrida*

Protoplasts from plants are suitable experimental objects for investigations in which substances are taken up by cells, since chemicals can reach the cell surface directly. Therefore, it is convenient to use them for experiments concerned with effects of phytohormones on metabolism. So far, some work on the application of auxins<sup>1-3</sup> has been described, and recently we have made some investigations on the influence of zeatin on photosynthesis and respiration of isolated mesophyll protoplasts from *Petunia* (KULL and HOFFMANN, in preparation).

Application of abscisic acid (ABA) leads to a decrease of photosynthetic rate in treated leaves, very likely due to stomatal closure caused by the phytohormone<sup>4-6</sup>. Some observations<sup>7</sup> suggest that mesophyll photosynthesis is not affected directly. Recently it was found that in illuminated and unilluminated dark grown *Avena* seedlings and in etio-chloroplasts of the same species, ABA reduced the levels of ribulosediphosphate carboxylase

activity<sup>8</sup>. We studied the effect of ABA on the photosynthesis of mesophyll protoplasts from *Petunia*. The influence of ABA on the respiration of these isolated protoplasts was briefly investigated in connection with our work on zeatin.

<sup>1</sup> D. W. GREGORY and E. C. COCKING, J. exp. Bot. 17, 68 (1966).

<sup>2</sup> J. B. POWER and E. C. COCKING, J. exp. Bot. 21, 64 (1970).

<sup>3</sup> M. H. BAYER, Plant Physiol. 51, 898 (1973).

<sup>4</sup> C. J. MITTELHEUSER and R. F. M. VAN STEVENINCK, Nature, Lond. 221, 281 (1969); Planta 97, 83 (1971).

<sup>5</sup> W. R. CUMMINS, H. KENDE and K. RASCHKE, Planta 99, 347 (1971)

<sup>6</sup> J. POSKUTA, R. ANTOSZEWSKI and M. FALTYNOWICZ, Photosynthetica 6, 370 (1972).

<sup>7</sup> R. J. JONES and T. A. MANSFIELD, Physiologia plant. 26, 321 (1972).

<sup>8</sup> F. A. M. WELLBURN, A. R. WELLBURN, J. L. STODDART and K. J. TREHARNE, Planta 111, 337 (1973).

Effect of abscisic acid on the photosynthetic rate of isolated mesophyll protoplasts from *Petunia hybrida*

Time ABA acting (h)	Concentration of ABA (racemate)				
	1 ppm	5 ppm	10 ppm	50 ppm	200 ppm
	3.8 $\mu$ M/l	19 $\mu$ M/l	38 $\mu$ M/l	190 $\mu$ M/l	760 $\mu$ M/l
1	100 $\pm$ 4.5	100 $\pm$ 4.5	100 $\pm$ 4.5	—	—
2	100 $\pm$ 4.5	100 $\pm$ 4.5	104 $\pm$ 4.5	131 $\pm$ 3.0	—
3	100 $\pm$ 4.5	103 $\pm$ 4.5	123 $\pm$ 3.0	198 $\pm$ 3.0	116 $\pm$ 3.0

Controls = 100%.

The mesophyll protoplasts from *Petunia hybrida* were produced by methods already described<sup>9,10</sup>. In scintillator vials 10<sup>6</sup> protoplasts in 1 ml medium (0.4 M mannitol/0.23 M NaHCO<sub>3</sub>, ratio 95:5; with 0.22 M HCl brought to pH 7.2) were incubated with ABA (racemate) of different concentrations (Table) for different times (1–3 h). Then 2  $\mu$ Ci of NaH<sup>14</sup>CO<sub>3</sub> in 100  $\mu$ l 0.4 M mannitol were added and the probes were illuminated for exactly 10 min in an illuminated thermostate (8  $\times$  40 W Osram-L-Warnton lamps, ca. 9000 lux, temperature 24 °C). After this, hot ethanol was added and the vials boiled for 2 min. After transferring into 50 ml graduated vessels and filling with 60% ethanol, the liquid was boiled again for 30 min, then the vessels filled up with ethanol. The activity of an equivalent of this final solution was measured in the scintillation counter in dioxane cocktail (gain 83%). Since ABA must be brought in solution with a few droplets of ammonia, the concentration of the latter rises with increasing concentration of the hormone. Therefore the respective concentrations of NH<sub>3</sub> were established in the controls. Always after addition of ABA or – in the controls – of NH<sub>3</sub> alone, the pH was adjusted again to 7.2. Due to the rising quantities of NH<sub>3</sub>, the absolute rates of photosynthesis, measured as amount of <sup>14</sup>CO<sub>2</sub> fixed, are not comparable; in the controls they decline. (In 1 set of control experiments without any NH<sub>3</sub> added, the rate is 2.98  $\mu$ M CO<sub>2</sub>/mg chlorophyll in 10 min; this is in good agreement with our earlier findings). Therefore the results are given as percentage of the photosynthetic rate of the respective controls. Each experiment was done 3 times together with 3 controls.

The results are shown in the Table. A rise of the photosynthetic rate caused by ABA can be seen. A marked effect needs relatively high concentrations of ABA with respect to naked protoplasts. With 1.9  $\times$  10<sup>-4</sup> M/l ABA after 3 h the increase of the photosynthesis reaches nearly 100%; with 7.6  $\times$  10<sup>-4</sup> M/l the effect is considerably lower, probably due to an inhibitory effect of very high concentrations of the phytohormone.

To a certain extent the results are surprising, considering that ABA is a senescence-promoting hormone<sup>11,12</sup>. The concentrations of ABA which cause a rise of the photosynthetic rate of mesophyll protoplasts are a great deal higher than those needed for closure of stomata and the subsequent inhibition of photosynthesis of

leaves<sup>6</sup>. Therefore an unspecific effect of ABA cannot be fully excluded. The results show the effects of a phytohormone on isolated cells in this case to be different from those on the whole leaf as a target. Under normal physiological conditions ABA is effective in whole tissues, so in evolution the function of the phytohormone was determined by selection in respect to the whole system and not to single cells. We think the results are explained best, in agreement with the findings of some other authors, as an effect of ABA on membrane permeability<sup>4,13</sup>, or the activity of ion uptake<sup>4,14–16</sup>. The reported<sup>8</sup> reduction of the activity of ribulosediphosphate carboxylase activity is due to an inhibition of plastid development and is not contrary to our findings with 'adult' cells<sup>17</sup>.

*Zusammenfassung.* Inkubation mit Abscisinsäure verursacht bei Mesophyll-Protoplasten von *Petunia hybrida* einen Anstieg der Photosyntheserate.

F. HOFFMANN<sup>18</sup> and U. KULL

*Lehrstuhl für Botanische Entwicklungsphysiologie, Universität Hohenheim, Emil-Wolff-Strasse, D-7000 Stuttgart 70, and Biologisches Institut der Universität Stuttgart, Ulmer Strasse 227, D-7000 Stuttgart 60 (Federal Republic of Germany). 11 February 1974.*

- <sup>9</sup> I. POTRYKUS and J. DURAND, *Nature New Biol.* 237, 286 (1972).  
<sup>10</sup> F. HOFFMANN, *Z. Pflanzenphysiol.* 69, 249 (1973).  
<sup>11</sup> F. T. ADDICOTT and J. L. LYON, *A. Rev. Plant Physiol.* 20, 139 (1969).  
<sup>12</sup> B. V. MILBORROW, *Sci. Prog., Lond.* 57, 533 (1969).  
<sup>13</sup> Z. GLINKA and L. REINHOLD, *Plant Physiol.* 48, 103 (1971). – Z. GLINKA, *Plant Physiol.* 51, 217 (1973).  
<sup>14</sup> T. A. MANSFIELD and R. J. JONES, *Planta* 101, 147 (1971).  
<sup>15</sup> R. F. M. VAN STEVENINCK, *Z. Pflanzenphysiol.* 67, 282 (1973).  
<sup>16</sup> G. N. BROWN and C. Y. SUN, *Physiologia plant.* 28, 412 (1972).  
<sup>17</sup> This work was supported by the German Bundesministerium für Wissenschaft und Forschung and by the Deutsche Forschungsgemeinschaft. We thank Prof. Dr. D. HESS for a helpful discussion, Mrs. A. KRAUS for technical assistance and Mr. P. CLEARY for correction of the English text.  
<sup>18</sup> New address: Max-Planck-Institut für Pflanzengenetik, Rosenhof, D-6802 Ladenburg.

### Der Einfluss der Aggression auf Lernen und Vergessen beim Kampffisch *Betta splendens* (Pisces, Anabantoidei)

Lern- und Behaltensprozesse werden stark von allem beeinflusst, was das lernende Individuum vor, während und nach dem Lernen tut und erlebt. Konsolidierung und Stabilität der Engramme sind offensichtlich am grössten,

wenn auf den Lernakt Perioden absoluter Passivität, etwa Schlaf, oder aber Ruheperioden folgen<sup>1–9</sup>. Zwischen Lernen und Reproduzieren eingeschobene Aktivitäten scheinen dagegen den Vergessensvorgang zu beschleunigen.