

Unlike their terrestrial counterparts, marine organisms have been found to contain very few structures resulting from enzymic coupling of phenols. Thelepin (5), from the annelid worm *Thelepus setosus*⁸, is the only clear example, although 2,2',4,4',6,6' hexahydroxybiphenyl (6), recently found in several brown algae^{9,10}, may also arise via this pathway.

1 Acknowledgment. We are grateful to Dr Klaus Reutzler for identifying the sponge and one of us (SW) is pleased to acknowledge fellowship support (1-F32-AI-05883-02) from the Public Health Service.

- 2 D.J. Faulkner, *Tetrahedron* 33, 1421 (1977), and references cited therein.
- 3 Aldrich Chemical Co., catalog No. 17, 340-1.
- 4 G. Cimino, S. DeStefano and L. Minale, *Comp. Biochem. Physiol.* 47B, 895 (1974).
- 5 I. Leifertova, N. Hejtmankova, H. Hlava, J. Kudrnacova and F. Santavy, *Sb. Prací pŕir. Fak. palack. Univ. Olomouci* 74, 83 (1975); *Chem. Abstr.* 85, 14611s, (1976).
- 6 J. Forrest, B.G. Overell, V. Petrow and O. Stephenson, *J. Pharm. Pharmac.* 4, 231 (1952).
- 7 E. Widmer, J.W. Meyer, A. Walser and E. Hardegger, *Helv. chim. Acta* 48, 538 (1965).
- 8 T. Higa and P.J. Scheuer, *Tetrahedron* 31, 2379 (1975).
- 9 K.-W. Glombitza, in: *Marine Natural Products Chemistry*, p.191. Ed. D.J. Faulkner and W.H. Fenical. Plenum Press, New York 1977.
- 10 M.A. Ragan and J.S. Craigie, *Can. J. Biochem.* 54, 66 (1976).

Selective effect of noradrenaline on superoxide dismutase activity in the brown adipose tissue and liver of the rat

V.M. Petrović, Tatjana Gudž and Zorica Saičić

Department of Endocrinology, Institute for Biological Research and Institute of Physiology and Biochemistry, University of Belgrade (Yugoslavia), 24 March 1980


Summary. Noradrenaline treatment results in a significant increase of superoxide dismutase activity in the intrascapular brown adipose tissue but not in the liver.

A metallo-protein from a few eukaryotic systems, including bovine erythrocytes, the liver, brain and heart, has been isolated previously, but superoxide dismutase (SOD) activity of these proteins was established only 10 years ago by McCord and Fridovich¹. It was found that this enzyme catalyzes the dismutation or disproportionation of superoxide free radical anions, yielding hydrogen peroxide and oxygen as follows: $O_2 + O_2 + 2 H^+ \rightarrow O_2 + H_2O_2$. The formation of these univalently reduced molecular oxygens is evident in various biological systems. Superoxide dismutase prepared mainly from the liver, erythrocytes and the brain has been intensely studied³. However, there are no data on SOD activity in the brown adipose tissue (BAT). This tissue is known to be under direct control of the sympathetic nervous system. We decided to study the effect of exogenous noradrenaline (NA) on SOD activity in the intrascapular brown adipose tissue (IBAT) and liver of the rat.

Materials and methods. Adult male Mill Hill hooded rats, weighing 180–200 g, and 2 months old, were used for the experiment. One group of 10 animals was treated with noradrenaline (Galenika, 1.6 mg/kg b.wt, i.p.) and the control group of 16 animals was injected with the same volume of saline solution. Animals were killed by decapitation 35 min after the injection of NA. IBAT and liver were taken perfused and homogenized at 4°C in a buffer con-

taining 0.05 M K_2HPO_4 and 10^{-4} M EDTA, pH 7.8, and centrifuged for 90 min at $85,000 \times g$. The supernatant was dialyzed for 20 h at 4°C. SOD activity was determined as described by Misra and Fridovich⁴. Protein was analyzed by the method of Lowry et al.⁵.

Results and discussion. As shown in the figure 1, SOD activity was significantly higher in the liver than in IBAT ($p < 0.005$). NA treatment produced a slight decrease in enzyme activity in the liver. However, the same amount of NA injected produced a significant increase in SOD activity in IBAT ($p < 0.01$). This increase was seen 35 min after the treatment, which was the time when a maximum calorogenic effect of this neurohormone was registered in previous experiments^{6,7}. NA is known to be a potent mediator of nonshivering thermogenesis, which occurs in the cold⁸. Under these conditions, BAT plays an important role in heat production⁹. In ambient cold, as well as under the influence of NA injected, the production of superoxide free radicals should be increased. Thus the increased SOD activity found in our present experiment in the IBAT of NA-treated rats may be particularly important in the protection against the toxicity of free oxygen radicals. In addition the selective effect of NA on SOD activity, found in the present experiments, may result from the higher capacity of the IBAT, as compared to the liver, to take up injected NA from the circulation.



Superoxide dismutase activity in the interscapular brown adipose tissue and liver of control and noradrenaline-treated rats (units/mg protein). Mean \pm SEM of 10 or 16 animals. The difference between the control and NA-treated rats in IBAT has $p < 0.01$.

- 1 J.M. McCord and I. Fridovich, *J. biol. Chem.* 244, 6049 (1969).
- 2 I. Fridovich, *Acc. chem. Res.* 5, 321 (1972).
- 3 R.A. Wiesiger and I. Fridovich, *J. biol. Chem.* 248, 3582, (1973).
- 4 H.P. Misra and I. Fridovich, *J. biol. Chem.* 247, 3170 (1972).
- 5 H.O. Lowry, J.N. Rosebrough, L.A. Farr and J.K. Randall, *J. biol. Chem.* 193, 265 (1951).
- 6 V.M. Petrović, K. Maksimović and L. Marković-Gajda, *Experientia* 32, 1544 (1976).
- 7 V.M. Petrović and K. Maksimović, *Bull. Acad. Sci. Serbe* 66, 81 (1979).
- 8 L. Jansky, *Can. J. Biochem. Physiol.* 41, 1847 (1963).
- 9 R.E. Smith and B.A. Horwitz, *Physiol. Rev.* 49, 330 (1969).