

### A new host of *Hemipteroseius indicus*

K. P. Shahi<sup>1</sup> and S. S. Krishna<sup>2</sup>

Entomology Laboratory, Department of Zoology, University of Gorakhpur, Gorakhpur – 273001 (U. P., India), 29 December 1980

**Summary.** The discovery of the predator mite, *Hemipteroseius indicus* capable of effectively reducing nymphal and adult populations of the phytophagous insect pest, *Dysdercus koenigii* infesting cotton has been reported for the first time.

Current literature on nymphal and adult parasites working as natural enemies of the red cotton bug, *Dysdercus koenigii* damaging cotton and okra in the tropics contains information only about a tachinid fly<sup>3,4</sup> and none on any species of mite. This communication places on record, for the first time, the occurrence of a mite, *Hemipteroseius indicus*<sup>5</sup> as a new and powerful predator attacking nymphs and adults of this bug.

**Observations.** The mite population consisting of all stages was largely confined to the dorsal part of the thoracic region around the scutellum in both male and female individuals of this insect where these comfortably sheltered predators were seen feeding intensely on the bugs. Predation neither inhibited mating between sexes in these hemipterans nor detrimentally affected the production of mature oocytes in the mated females. However, the gravid female, if preyed upon severely, died without laying any eggs although she manifested the typical distended condition of the abdomen caused by the presence of ripe oocytes in her ovaries. Predatory attack on nymphs occurs even in the newborn 1st instar individuals which, like those of later instars, harbour these acarines in the ventral region of their

thoraces, especially at the base of the legs. Nymphs of any instar heavily preyed upon succumbed without progress in their development.

**Scope.** Future investigations can explore the possibility of considering this acarine species as a potential biological agent for inclusion in pest suppression programs formulated to check the multiplication and eventual establishment of *D. koenigii* in cotton fields.

- 1 Acknowledgment. This author conveys his grateful thanks to the Council of Scientific and Industrial Research, New Delhi, for the award of a Fellowship which made it possible for him to undertake this work.
- 2 To whom request for reprint should be made.
- 3 G. S. Sohi, in: Entomology in India, p. 111. Ed. N. C. Pant. The Entomological Society of India, New Delhi 1964.
- 4 T. J. Crowe, in: Diseases, pests and weeds in tropical crops, p. 298. Ed. J. Kranz, H. Schmutterer and W. Koch. Paul Parey, Berlin 1977.
- 5 Identification of the species made possible through the courtesy of the staff of the Zoological Survey of India, Calcutta.

### Functions of nuclear bodies as revealed by ultrastructural autoradiography and cytochemistry

G. Yasuzumi, N. Yabumoto, T. Shirai, Y. Nakai and F. Yasuzumi

Academy of Medical Technology and Nursing, Hanwa Memorial Hospital, Karita-7, Osaka 558 (Japan); Department of Dermatology, Nara Medical University, Kashihara 634, Department of Anatomy, Showa University School of Medicine, Tokyo 142 (Japan), and Department of Anatomy, Fukuoka University School of Medicine, Fukuoka 814 (Japan), 14 October 1980

**Summary.** The simple nuclear body containing a few RNA particles appears through the nuclear pores in the cytoplasm, originating from the nucleolus. The complex nuclear body consisting mainly of RNA components is highly active in the incorporation of RNA precursors. Accordingly, the appearance of nuclear bodies may be related either to transport to the cytoplasm of nucleolar components or to the enhancement of rRNA synthesis.

Nuclear specific components appearing often in cancer and precancer cells were first designated 'nuclear bodies' by Weber et al.<sup>1</sup>. Their fine structure and chemical components, both simple and complex types, according to Bouteille et al.<sup>2</sup>, are now well known. To our knowledge their function remains unsettled as yet, although suggestions have been made by several authors<sup>1-6</sup>.

**Material and methods.** Biopsy materials were obtained by operative excision from the following patients: 4 male and 2 female patients (42–50 years old) with Bowen's disease; precancerous dermatosis with chronic atypical epithelial proliferation of the skin, and 4 male patients (50–60 years old) with senile keratosis; actinic keratosis induced by sunlight or X-rays. The materials were cut into tiny blocks, fixed in 2.5% glutaraldehyde alone, or fixed in 2.5% glutaraldehyde and postfixed in 1.0% osmium tetroxide, and embedded in epoxy epon resin by routine methods. Ultrathin sections were stained with saturated uranyl acetate

followed by lead citrate and examined with an electron microscope, either HU-11D-S or JEM-100C.

Sections of samples fixed only with glutaraldehyde were used for extraction of DNA using Bernhard's EDTA technique<sup>7</sup>. Since nuclear bodies of several types appear in lymphoblasts in the germinal center of normal mouse spleen<sup>8</sup>, we have tried to observe similar materials by an autoradiographic electron microscopic technique. Swiss F26 strain mice, 12 weeks old, weighing 28–30 g, were injected i.p. with 0.9 mCi of uridine (New England Nuclear, NET-367; sp. act. 37.6 Ci/mmol). The injected animals were sacrificed after 1 h. The autoradiographic technique of Granboulan<sup>9</sup> was used with Ilford L4 emulsion diluted 1:5 in distilled water.

**Results and discussion.** Many of the nuclear bodies originating from the nucleolus<sup>2,6</sup> in Bowen's disease migrate to the periphery of the nucleoplasm to attach to the nuclear envelope, where the content of the nuclear body seems to