

- 8 Powell, R. G., and Smith, C. R. Jr, in: *Alkaloids*, vol. 2, p. 149. Ed. S. W. Pelletier. John Wiley & Sons, New York 1984.
- 9 Reider, P. J., and Roland, D. M., in: *The Alkaloids*, vol. 23, p. 71. Ed. A. Brossi. Academic Press, Orlando 1984.
- 10 Spjut, R. W., Cassady, J. M., McCloud, T., Suffness, M., Norris, D. H., Cragg, G. M., and Edson, C. F., *Econ. Bot.* 42 (1988) 62.
- 11 Spiess, L. D., Lippincott, B. B., and Lippincott, J. A., *J. Hattori Bot. Lab.* 55 (1984) 67.
- 12 Caldwell, B. A., Hugedorn, C., and Denison, W. C., *Microbial Ecol.* 5 (1979) 91.
- 13 Crass, S. M., and Scheirer, D. C., *Bryologist* 84 (1981) 348.
- 14 Fenton, J. H. C., *Trans. Br. mycol. Soc.* 60 (1983) 415.

0014-4754/90/010117-04\$1.50 + 0.20/0
© Birkhäuser Verlag Basel, 1990

Corrigendum

Due to a regrettable technical error parts of the multi-author review (Phylogeny and Function of the Pineal/*Experientia* 45/10) contribution, 'The pineal and melatonin: Regulators of circadian function in lower vertebrates' by H. Underwood, were misplaced in the final text. We are therefore reprinting the corrected article in its entirety. When citing, please refer to both publications:

Experientia 45 (1989) 914–922; 46 (1990) 120–128.

The pineal and melatonin: Regulators of circadian function in lower vertebrates

H. Underwood

Department of Zoology, North Carolina State University, Raleigh (North Carolina 27695, USA)

Summary. The pineal has been identified as a major circadian pacemaker within the circadian system of a number of lower vertebrates although other pacemaking sites have been implicated as well. The rhythmic synthesis and secretion of the pineal hormone, melatonin, is suggested as the mechanism by which the pineal controls circadian oscillators located elsewhere. Both light and temperature cycles can entrain the pineal melatonin rhythm. The pineal, therefore, acts as a photo and thermoendocrine transducer which functions to synchronize internal cycle with cycles in the environment. A model is presented which portrays the pineal as a major component of a 'multioscillator' circadian system and which suggests how these multiple circadian clocks are coupled to each other and to cycles of light and temperature in the external world.

Key words. Pineal; melatonin; circadian rhythm.

Organisms, from unicellulars to vertebrates, are structured in time as well as in space. Many, if not most, biochemical, physiological and behavioral parameters exhibited by organisms show daily fluctuations. Significantly, most of these daily rhythms will persist under constant conditions, with periods near 24 h in length, showing that they are driven by an internal daily or 'circadian clock'. Under natural conditions these rhythms are 'entrained' by environmental stimuli (such as light and temperature) so that the 'right' events occur at the 'right time of day'.

The use of the term 'circadian clock' does not mean that a single discrete 'clock' is responsible for driving all of an organism's daily rhythms. In recent years it has become appreciated that multicellular organisms are multi-oscillator in nature; that is, more than one circadian clock

may exist within a single organism. Normally, however, all of an organism's multiple circadian rhythms exhibit fixed phase relationships with each other under both entrained and freerunning conditions. Accordingly, when multiple clocks exist within an individual organism, they must be coupled together in some manner.

Several areas have been implicated as being important to circadian organization in vertebrates; these include the pineal organ, the suprachiasmatic nuclei (SCN) of the hypothalamus, and the lateral eyes. Interesting similarities, as well as significant differences, seem to exist between species in the relative roles that these areas play within an animal's circadian system.

Among lower vertebrates most of the studies to date have focused on the role of the pineal organ. Embryologically, the pineal arises as an evagination of the roof of the

diencephalon and, in lower vertebrates, the pineal is situated on the brain's surface and remains attached to the brain by a stalk^{26, 36, 45}. Pineal organs are present in virtually all vertebrates and, in addition, a second component of the pineal 'system', generally termed a parapineal organ, is often present in lower vertebrates which arises from a more rostral differentiation of the diencephalic roof^{26, 36, 45}. During development the parapineal component (called a frontal organ in anuran amphibians or a parietal eye in lizards) migrates to a position close to the roof of the skull. Parapineal organs are absent in birds, mammals, and in some lower vertebrates (such as snakes and toads).

The pineal organs of lower vertebrates typically contain several types of cells including photoreceptors and nerve cells (Falcón and Collin, this issue)^{26, 36, 45}. The photoreceptive cells resemble the photoreceptors of the lateral eyes and have been found in petromyzontides, fish, amphibians, chelonians, and lacertilians. The photoreceptive cells may possess a well-defined outer segment with stacked lamellar disks as well as an inner segment and a cell body with a synaptic pedicle which establishes contact with one or more secondary neurons, or they may be more 'rudimentary' in appearance and exhibit disorganized or rudimentary outer segments and lack contact with secondary neurons. Rudimentary photoreceptors are characteristic of the pineal organs of chelonians and lacertilians. The pineal photoreceptors are hypothesized to be the precursors of the (secretory) pinealocyte which is the predominant cell type of ophidians, birds, and mammals.

The principal cell type in the parapineal organs of lower vertebrates is also photosensory in appearance. The ultrastructural evidence of photoreception by the parapineal (and the pineal) organs of lower vertebrates is supported by electrophysiological studies²⁶. In general, pineal organs show an achromatic electrical response to illumination whereas parapineal organs are often capable of wavelength discrimination (chromatic response). In fish, amphibians and reptiles, nerve cells found within the pineal and parapineal organs project axons carrying photic information to pretectal and tegmental centers^{26, 45}. From lower to higher vertebrates there is a definite phylogenetic trend toward a decrease in the intrapineal neurons and their pinealofugal fibers. Also, whereas pinealopedal (sympathetic) innervation, originating in the superior cervical ganglia, has been well documented in birds and mammals, pinealopedal innervation in lower vertebrates has been poorly studied. The pineals of some lower vertebrates appear to contain sympathetic (efferent) nerves but the origin of these nerves is uncertain.

Despite the photosensory appearance of the main cell type in the pineals of lower vertebrates, ample evidence exists that these cells are also capable of considerable biosynthetic activity. For example, immunohistochemical and autoradiographic studies suggest that the pineal

photosensory-type cells are the sites of synthesis of indoleamines, including melatonin (Falcón and Collin, this issue)^{15, 25, 67}.

Multiple photic inputs to the circadian system

Removal of the eyes does not prevent entrainment (synchronization) of the circadian system by 24 h light-dark cycles in any of the nonmammalian vertebrates examined so far^{1, 2, 4, 12, 24, 33, 51, 59, 61, 62}. Extraretinal photoreceptors, therefore, exist in fish, amphibians, reptiles and birds which are fully capable of mediating entrainment. The photosensory capacities of the pineal systems of lower vertebrates have made them obvious candidates as the extraretinal photoreceptors. In a variety of fish and reptiles, however, removal of the pineal organ, or the parapineal organ, failed to abolish entrainment in blinded animals^{33, 59, 61, 62}. It has been suggested that the frontal organ of frogs may be the extraretinal photoreceptor mediating entrainment since removal of the frontal organ was reported to abolish entrainment of blinded green frogs (*Rana clamitans*)². The small sample size and the 'noisiness' of the activity rhythm, however, precludes any firm conclusions about the role of the frontal organ in *R. clamitans*. In another anuran, *Xenopus laevis*, entrainment of the activity rhythm of blinded *Xenopus* to LD cycles was unaffected by removal of the frontal organ⁴. Although these data show that entrainment can persist after removal of the eyes and pineal system – showing the universal existence of extrapineal-extraretinal receptors in lower vertebrates – the conclusion that pineal photoreception is not involved in mediating photic influences on the circadian system is unwarranted. In view of the major role of the pineal organ in circadian system in some vertebrates (see below), and in view of the fact that light can directly affect pineal biochemistry^{11, 20, 28, 37, 38}, it seems probable that direct pineal photoreception is important to circadian organization, at least in some lower vertebrates.

The nature of the role of extraretinal photoreceptors (ERRs) and the eyes as photic inputs to the circadian system of lower vertebrates has been poorly studied. The (extrapineal) ERRs have been localized to the brain in fish, amphibians and reptiles, but their location(s) within the brain are not known^{1, 61, 62}. Several observations suggest that the lateral eyes have a photic input to the circadian system of lower vertebrates including: a) Removal of the eyes of lizards (*Sceloporus olivaceus*) entrained to very dim LD cycles can cause them to freerun⁶¹. b) Blinding affects several light-dependent circadian properties in the lizard *Sceloporus occidentalis* including range of entrainment, rate of reentrainment to shifted LD cycles, and the sensitivity to light pulses⁵⁵. c) In trout, *Salvelinus fontinalis*, entrainment to natural LD cycles is weaker in blinded fish at certain times of year^{13, 14}. d) In the lake chub, *Couesius plumbeus*, blinded fish have a more restricted spectral sensitivity and a

higher energy requirement for entrainment than sighted fish³³.

One important question that arises is: Why do multiple photic inputs to the circadian system of lower vertebrates exist? The fact that ERRs exist, in addition to the lateral eyes, suggests that the different photoreceptors may be processing different kinds of photic information. This idea is supported by studies in lizards in which it has been shown that the eyes and ERRs are extracting different kinds of information from the photic environment. Diurnal animals typically exhibit a shorter freerunning period when exposed to constant light (LL) than when exposed to constant dark (DD). This empirical generalization has been termed 'Aschoff's Rule'. The lizard *Lacerta sicula* obeys Aschoff's Rule when sighted but, after blinding, the lizard's freerunning period is unaffected when the lizard is switched from DD to LL⁶¹. Blind *L. sicula*, however, are still entrainable by LD cycles. These data show that the eyes mediate the continuous (parametric) effects of light on the velocity of the circadian oscillator but ERRs can mediate entrainment via the light to dark (transitional or non-parametric) effects of light. Clearly, in this species, different photoreceptors (eyes versus ERRs) are involved in extracting different kinds (parametric versus non-parametric) of photic information. Other species of lizards, however, can obey Aschoff's Rule even after blinding showing that the parametric effects of light can be mediated via ERRs in these species^{51, 61}. The reasons for these differences among species are unknown but may well be related to the particular ecological niches to which these animals have evolved.

Pineal as a pacemaker

A major role for the pineal within the circadian systems of lower vertebrates is revealed by observing the effects of pinealectomy on the freerunning activity rhythms (fig. 1). Pinealectomy can cause: 1) marked changes in the freerunning periods of fish (e.g., *Couesius plumbeus*, *Lota lota*, *Catostomus commersoni*) and lizards (e.g., *Sceloporus olivaceus*, *Sceloporus occidentalis*)^{31, 32, 34, 50, 51}, 2) splitting of the activity rhythm into two major circadian components in a fish (*Catostomus commersoni*) and a lizard (*S. olivaceus*)^{30, 50}, or 3) arrhythmicity in lizards (e.g., *Anolis carolinensis*, *S. olivaceus*, *S. occidentalis*) and fish (e.g., *Heteropneustes fossilis*)^{19, 50, 51, 53}. Pinealectomy of the lizard *S. occidentalis* may also cause a major distortion of the phase response curve to light pulses⁵². Note that a given species may exhibit more than one kind of response to pinealectomy. These kinds of results are compatible with the hypothesis that the pineal can act as a pacemaker within a multi-oscillator circadian system. Obviously, circadian oscillators must also exist outside the pineal since, in some lizards and fish, rhythmicity can persist after pinealectomy. Both the inter- and intraspecies differences in the effects of pinealectomy could be explained by individual




Figure 1. Effects of pinealectomy (P) on the circadian activity rhythms of several species of lizard (A) and fish (B). Pinealectomy has various effects including splitting of the activity pattern into two components (*S. olivaceus*, *C. commersoni*), arrhythmicity (*A. carolinensis*, *H. fossilis*), and period changes (*S. occidentalis*, *L. lota*). The original records from which these patterns were generated can be found in: (50, *S. olivaceus*), (53, *A. carolinensis*), (51, *S. occidentalis*), (30, *C. commersoni*), (19, *H. fossilis*), (32, *L. lota*).

variations in the strength of coupling among these extrapineal circadian oscillators. For example, if coupling among extrapineal oscillators is weak (or absent) pinealectomy will cause splitting or arrhythmicity. If the extrapineal oscillators remain coupled following pinealectomy, a change in the period of the activity rhythm will be observed.

The pineal hormone, melatonin, appears to play an important role in the circadian system of lower vertebrates. Immunohistochemical and autoradiographic studies suggest that the pineal photoreceptive cells are the likely sites of melatonin production (Falcón and Collin, this issue)^{15, 25}. Melatonin is highly lipid soluble and appears to be rapidly secreted into the blood by passive diffusion. Pineal melatonin production is rhythmic; melatonin levels at night are higher than those occurring during the day^{17, 20, 38, 54, 63}. In higher vertebrates the daily rhythm in melatonin is believed to result from a rhythm in activity of the rate-limiting enzyme N-acetyltransferase (NAT), which converts serotonin to N-acetylserotonin. The terminal enzyme in the melatonin synthesizing pathway, hydroxyindole-O-methyltransferase (HIOMT), is not believed to contribute to melatonin rhythmicity in higher vertebrates since its activity remains relatively constant throughout the day (Sugden, this issue). However, in some lower vertebrates (e.g., lamprey, *Lampetra planeri*; salmon, *Oncorhynchus tshawytscha*; lizard, *Lam-*

propholas guichenoti), but not others (e.g., pike *Esox lucius*), HIOMT activity or amount can also show significant daily (and seasonal) fluctuations^{8, 9, 16, 25, 29}. The contribution of NAT versus HIOMT activity to melatonin rhythmicity in these cases has not been well studied. A role for melatonin in circadian organization is shown by the fact that: a) continuous melatonin administration can elicit period changes or arrhythmicity in some lizards⁵¹, b) daily melatonin injections can entrain the activity rhythm of the lizard *S. occidentalis*⁶⁰, c) a phase response curve to melatonin exists in a lizard (*S. occidentalis*)⁵⁶, and d) melatonin pulses can phase-shift celestial orientation in the tiger salamander, *Ambystoma tigrinum*³ (accurate celestial orientation relies on the circadian clock to compensate for the apparent movement of celestial cues).

These data suggest that the daily rhythm of melatonin secretion may be the mechanism by which the pineal communicates with the rest of the circadian system. The entrainment of circadian clocks by entraining stimuli such as light (or melatonin) requires that the circadian clock exhibit a diurnal change in sensitivity to that stimulus; that is, a phase response curve to that stimulus must exist⁴³. All known phase response curves are qualitatively similar. If it is hypothesized that the phase response curve to melatonin differs (i.e., quantitatively) among extrapineal clocks, the daily melatonin pulse could, at least theoretically, not only entrain these clocks but also exert phase control over them so that the myriad overt rhythms they control (i.e., locomotor activity, hormone levels, enzyme activities) would occur at the appropriate time of day.

In vitro studies of pineal melatonin secretion have been conducted in fish and lizards (fig. 2). In the trout *Salmo gairdneri* melatonin production is cyclical under 24-h LD cycles, showing that the organ cultured pineal is capable of direct photoreception, but no rhythmicity is observed in DD²⁰. Similarly, in the iguanid lizard *Dipsosaurus dorsalis* a rhythm is seen in organ cultured pineals exposed to LD but not in pineals held in DD²⁸. In another iguanid lizard, *Anolis carolinensis*, however, a temperature compensated rhythm of melatonin secretion persists for up to 10 circadian cycles in DD; this rhythm is directly entrainable by LD cycles³⁸. A circadian rhythm in melatonin production in vitro is also seen in the iguanid lizard *S. occidentalis* in DD³⁷. The studies on *A. carolinensis* and *S. occidentalis* show that a circadian pacemaker must reside within their pineals which is capable of driving a rhythm of melatonin secretion. The results in *A. carolinensis* and *S. occidentalis* are compatible with the hypothesis that the pineal can act as a circadian pacemaker within a multioscillator circadian system. Such a role is also supported by the observation that pinealectomy in these two species induces period changes in the activity rhythm, or arrhythmicity, in DD^{51, 53}. Interestingly, pinealectomy of another iguanid lizard, the desert iguana *D. dorsalis*, has little effect on the freerunning




Figure 2. The in vitro pattern of melatonin secretion from individual pineal glands of several species of iguanid lizard (*A. carolinensis*, *S. occidentalis*, *D. dorsalis*) and one species of fish (*S. gairdneri*) held in either 24-h LD cycles (left panels) or in DD (right panels). The figures are redrawn from the following sources: (38, *A. carolinensis*), (37, *S. occidentalis*, courtesy The Ciba Foundation), (28, *D. dorsalis*, courtesy D. Janik), (20, *S. gairdneri*, courtesy Academic Press).

activity rhythm and, as mentioned previously, the pineal of this species is also incapable of expressing a persistent circadian rhythm of melatonin production in vitro²⁸.

In those species in which a circadian rhythm of melatonin production is absent in vitro but a rhythm can be generated by exposing the pineal to LD cycles several possibilities can be entertained. 1) A circadian pacemaker may be absent in the pineal of these species but a rhythm in melatonin production can be exogenously generated by the periodic input of a photic, neural or hormonal signal. That is, the pineal is not capable of acting as a pacemaker but may be part of a driven system. 2) A pacemaker exists but the culture conditions are inadequate to allow expression of a rhythm. 3) A pacemaker exists but is not coupled to melatonin production.

Clearly, more studies are needed to determine, within the same species, the role of the pineal in the circadian system (i.e., by observing the effects of pinealectomy on the activity rhythm) and the behavior of the pineal in vitro. That is, are those species in which pinealectomy is ineffective in disturbing circadian rhythmicity the same spe-

cies in which the pineal is incapable of expressing a circadian rhythm in melatonin secretion? The current data do, however, suggest that, in at least some species, the pineal can act as a circadian pacemaker and it is coupled to the rest of the system via the rhythmic secretion of the hormone, melatonin.

The role of potential neural inputs to the pineals of lower vertebrates has not been studied. Efferent (sympathetic) nerve terminals have been seen but their origin and function is unknown. Speculatively, these nerves could serve as routes by which extrapineal pacemakers are coupled to the pineal or they might be routes by which light perceived by the eyes or ERRs could influence the pineal. Also unstudied is the role of neural outputs from the pineal (or parapineal). These afferent nerves are undoubtedly transmitting photic information to pretectal and tegmental centers^{26, 36}, but the role of these photic inputs (if any) within the circadian system is unknown.

The pineal: A photothermoendocrine transducer

The pineal can function as a transducer of both thermal and photic information in poikilotherms. Studies have shown that either light or temperature cycles can entrain the pineal (or plasma) melatonin rhythm and aspects of the light or temperature cycles (such as the duration of the dark and light portions of the LD cycles or the amplitude of the temperature cycle) can influence aspects of the pineal melatonin rhythm (such as its amplitude, phase, or duration)^{18, 21, 22, 54, 58, 63–65}. When presented together, an interaction between light and temperature cycles on the pineal (or plasma) melatonin rhythm can be observed^{17, 18, 22, 58} (fig. 3). The annual changes in the daily melatonin rhythm that are observed in animals exposed to natural conditions (fig. 4) undoubtedly reflect the interactive effects of both light and temperature on pineal melatonin synthesis. Light and temperature can undoubtedly *directly* affect the pineal melatonin rhythm, as revealed by *in vitro* studies^{20, 28, 38}, but the possibility that photic (or thermal) inputs can reach the pineal indirectly from other brain areas cannot be discounted. One possible way in which light and temperature may control the phase, amplitude, or duration of the pineal melatonin rhythm could reside in the putative 'multi-oscillator' nature of the pineal itself. Dispersed cell cultures of chicken pineals, for example, can exhibit light-entrainable circadian rhythms in NAT activity showing that the pineal, itself, can be multioscillator in nature¹¹. If a number of individual cells (i.e., the photo-sensory cells?) within the pineals of lower vertebrates possess circadian properties, the mutual phase relationships among these oscillators will determine the phase, duration, and amplitude of the cumulative output of these cells (i.e. melatonin). Since both light and temperature can act as entraining agents for circadian clocks in poikilotherms, the way in which these stimuli entrain the multiple pineal oscillators will determine the phase,




Figure 3. Interaction of light and temperature in the control of the pineal melatonin rhythm of the lizard *Anolis carolinensis*. The lizards were exposed to a temperature cycle in which the cool phase of the cycle occurred either during the day (left panels) or night (right panels) of a LD 12:12 light cycle. In the 'abnormal' light/temperature conditions (left panels: cool days/warm nights) the temperature cycles controlled the phase of the melatonin rhythm when its amplitude was 12°C (top), both light and temperature interacted to control phase when the amplitude of the temperature cycle was 5°C (middle), but the light cycle controlled phase when the amplitude of the temperature cycle was 2°C (bottom).




Figure 4. Annual changes in the daily rhythm of pineal melatonin content in the tortoise, *Testudo hermanni*, held under natural conditions of photoperiod and temperature (Adapted from 63, courtesy Academic Press).

amplitude, and duration of the melatonin rhythm. Such a model could also be applied to pineals which do not exhibit persistent circadian rhythmicity *in vitro* (such as the trout or desert iguana) if the melatonin producing cells, although lacking persistent circadian rhythmicity, can act as damped oscillators which require daily stimulation by light or temperature to generate a melatonin rhythm. In this way the non-autonomous pineal could still act as a photothermal transducer but it would be expected that the role of the pineal within the total circadian system would be less.

The effect of light and temperature cycles on the pineal melatonin rhythm, therefore, offers a mechanism by which these stimuli can entrain the circadian system appropriately. That is, aspects of the pineal melatonin profile (i.e., phase, duration) directly reflect ambient photic and thermal conditions. The pineal melatonin rhythm, in turn, can phase extrapineal oscillators appropriately so that the overt rhythms they drive occur at the 'right time of day'.

One feature of the response of the pineals of at least some lower vertebrates to light is remarkably different from that observed in birds and mammals. In higher vertebrates light pulses presented at night cause a rapid (within minutes) fall in pineal melatonin levels⁴⁷. This fall may not represent a 'direct' response to light (which in mammals is perceived exclusively via the eyes) but may reflect the way light shifts the phase of the circadian clocks driving NAT activity²⁷. That is, light may cause a rapid shift of the phase of the clock to a phase at which it does not call for melatonin synthesis. In lower vertebrates, however, night breaks can depress melatonin levels in some (e.g. trout, *Salmo gairdneri*) but not in others (e.g. lizard *A. carolinensis*; turtle, *Terrapene carolina triunguis*)^{20, 57, 65}. In the pike (*Esox lucius*) night breaks induced inactivation of roughly half of the maximal NAT activity indicating the existence of one photolabile and one photostable enzymatic component¹⁶. Clearly, additional studies are needed to determine the basis of the different kinds of response to night breaks – are these differences due to different forms of the NAT enzyme or to the way in which light pulses shift the phase of the oscillators driving the melatonin rhythm? Also, what is the adaptive value of the different kinds of response of pineal melatonin to night breaks?

Extrapineal melatonin production

Although it was originally believed that melatonin was a unique pineal product since HIOMT activity was thought to be uniquely localized in pineal tissue, it is now recognized that some extrapineal sites can generate melatonin⁴⁶. In at least some lower vertebrates melatonin synthesis can occur in the eyes, the parapineal organ, and possibly the Harderian glands^{17, 36, 41, 46, 66–68}. Although it has been shown that blood levels of melatonin in higher vertebrates reflect mainly pineal melatonin se-

cretion^{46, 67} comparable studies in lower vertebrates are scarce^{21, 23, 28}. In some species, such as the trout, the tiger salamander, and the desert iguana, pinealectomy does not completely eliminate blood-borne melatonin^{21, 23, 28}. Clearly, in those species in which melatonin is important to circadian organization the potential exists for extrapineal melatonin-producing sites to play a role within the circadian system.

Extrapineal pacemakers

SCN. In mammals a host of studies have shown that the SCN of the hypothalamus are the site of a neural circadian pacemaker which drives many (if not all) of the animals' circadian rhythms³⁹. Although fewer in number, available evidence suggests that the SCN may play a similar role in birds as well⁴⁹. A direct retinohypothalamic (RH) tract to the SCN is the major route by which lighting information reaches the mammalian SCN³⁹. In lower vertebrates, nuclei presumed to be homologous to the mammalian and avian SCN have been described, as well as the existence of a RH pathway^{28, 44, 48}. However, only a couple of studies have examined the role of the hypothalamus in circadian organization in lower vertebrates. Surgical ablation of the optic tectum, the anterior portions of the telencephalon, and the dorsal portions of the telencephalon-diencephalon of the hagfish, *Eptatretus burgeri*, failed to affect the freerunning activity rhythm⁴⁰. However, ablation of the ventromedial part of the telencephalon-diencephalon, which included the hypothalamus, caused hagfish to become arrhythmic in DD⁴⁰. Interestingly, the hagfish is one of the most primitive vertebrates and it is one of the few animals in which a pineal organ is reported to be absent. Even if there is 'pineal equivalent' tissue in the dorsal diencephalon of the hagfish, it does not seem to have a pacemaking function since removal of the dorsal portions of the diencephalon did not abolish the circadian activity rhythm⁴⁰. Lesions of the SCN of the desert iguana, *D. dorsalis*, abolish the circadian activity rhythm²⁸ (fig. 5).

The data from lower vertebrates, although sparse, suggest that the suprachiasmatic area of the hypothalamus is important for circadian organization and may act as a neural circadian pacemaker. The SCN, therefore, may be a site for at least some of the extrapineal pacemakers that exist in lower vertebrates.

Eyes. A small but growing body of evidence suggests that the eyes of some vertebrates may play a role within the circadian system which is not merely a photosensory one. There is some evidence that the eyes may be the loci of circadian pacemakers. Numerous studies have demonstrated that events within the eyes of lower vertebrates, such as photomechanical movements of rods and cones, shedding of discs from photoreceptor outer segments, visual sensitivity, and NAT activity show daily rhythms and, in some cases, these rhythms have been shown to persist in constant conditions^{5–7, 10, 35, 42}. In one study




Figure 5. Locomotor activity pattern of a lizard (*Dipsosaurus dorsalis*) held under constant conditions and subjected to a lesion (L) of the suprachiasmatic area of the hypothalamus. The lizard showed a well-defined circadian rhythm of activity until the day of the lesion. Subsequently, the activity became arrhythmic (28, courtesy D. Janik).

the site of the oscillator driving an ocular rhythm was shown to reside within the eye itself⁷. This study utilized an in vitro eye explant preparation from the African clawed frog, *Xenopus laevis*, and showed that a rhythm in NAT activity could persist in the isolated eye in constant conditions⁷.

Some rather paradoxical effects of blinding on the circadian system of the lizard *S. occidentalis* also suggest that the eyes may play an important role within the lizards' circadian system⁵⁵. Blinding *S. occidentalis* by optic nerve section actually renders them more sensitive to the effects of light since: 1) single light pulses cause a larger shift in the phase of the freerunning activity rhythm in blind lizards than in sighted ones, 2) blind lizards reentrain more rapidly than sighted lizards to a shift in the phase of an entraining LD cycle, and 3) the limits of entrainment are larger in blind lizards than in sighted lizards⁵⁵. One possible model to explain these results would postulate that the eyes can act as oscillators within a multioscillator circadian system. Uncoupling the ocular oscillators from the rest of the system renders the remaining components more sensitive to the phase shifting effects of light.

In some lower vertebrates the eyes are capable of synthesizing melatonin and there is some evidence that the photoreceptors within the eye are the sites of this melatonin synthesis^{7, 29, 41, 42, 66–68}. In some species, such as *Xenopus*, melatonin does appear to play a role in controlling some daily rhythms within the eye itself⁴² but it is not currently known if retinally produced melatonin is capa-

ble of acting on the circadian system at sites outside the eyes.

Circadian organization in lower vertebrates: A model

The results from studies on the circadian systems of a number of species of lower vertebrates can be summarized in the following model (fig. 6). Light and temperature, the primary entraining stimuli for poikilotherms, can act at several different levels of the circadian system. First, both light and temperature can be transduced into a hormonal signal (melatonin) at the level of the pineal. Second, light can act via extraretinal-extrapineal receptors (ERRs) to entrain central oscillators and, since temperature cycles can entrain the activity rhythm of pinealectomized lizards (Underwood, unpubl.), temperature can also directly entrain extrapineal oscillators. Third, light (and possibly temperature) can also act on the circadian system via the eyes.

The pineal, itself, may be autonomously rhythmic and comprised of a population of coupled oscillators. Light and temperature can entrain (and phase) the output of these oscillators (melatonin) so a particular melatonin profile (i.e., phase, amplitude, or duration) reflects the particular environmental stimuli to which the animal is exposed. The pineal melatonin rhythm, in turn, is responsible for entraining circadian clocks located elsewhere (possibly in the SCN) so that these oscillators assume phase-relationships appropriate to those environmental conditions. The coupling strength between the extrapineal oscillators determines the behavior of the system after pinealectomy (i.e. whether splitting, arrhythmicity, or period changes occur). Whether feedback from extrapineal oscillators to the pineal occurs is unknown. Some data suggest that the eyes may also be the loci of circadian oscillators but the importance of these ocular clocks within the animals' total circadian system is unknown.

In some species (e.g., the iguanid lizard *D. dorsalis*) the pineal may play only a minor role within circadian organization and, in these species, the pineal may be unable to sustain a melatonin rhythm in the absence of periodic inputs.

According to the model shown in figure 6 environmental stimuli can act at more than one level on the circadian system. Some as yet poorly understood selective advantages must accrue to such a system. Undoubtedly, aspects of this kind of system must also reflect the organism's evolutionary history. For example, extraretinal photoreceptors may well have evolved early and then been retained after the evolution of the eyes. Or the pineal may have acted as a primitive photoreceptor with melatonin acting only within the pineal itself. Later, melatonin may have been secreted into the blood and evolved a more 'systemic' role within the animals' total circadian 'system'.




Figure 6. A model for the circadian system of lower vertebrates. Light and temperature conditions (i.e., summer versus winter conditions) are transduced by the (multioscillator) pineal into a melatonin pulse which, in turn, controls both the period and phase of circadian oscillators located elsewhere (i.e., in the SCN?). Other components of the system (shown in the left panel only) include: 1) the eyes which act as photoreceptors and

(possibly) as biological clocks, 2) 'direct' entrainment of extrapineal oscillators by temperature and light (via extraretinal photoreceptors) and, 3) possible links between central oscillators and oscillators in the pineal and eyes. This system allows coordination between internal circadian clocks, and between those clocks and the external world, so that the right events occur at the 'right time of day'.

Circadian organization in the lower vertebrates clearly involves a number of components including the pineal, the eyes, and the SCN. However, it is becoming increasingly apparent that the relative roles that these sites play between species can vary. This variation can be seen even at the level of the family (e.g. lizards of the family Iguanidae). Little is known about the reasons for this variation but they probably reflect the different selection pressures operating on animals which occupy diverse ecological and temporal niches.

The major weakness of the model depicted in figure 6 arises from the fact that it represents a composite drawn from the results of experiments on a variety of species but, in no case, has the role of all of the elements depicted been thoroughly examined in a single species. Clearly, a major goal of future research should be a comprehensive examination of the roles of the individual elements of the circadian 'system' within individual species while paying close attention to the particular niche which the species occupies. These data would increase our confidence in the model and would yield insights into why different species have opted to emphasize one element over another.

- Adler, K., Extraoptic phase shifting of circadian locomotor rhythm in salamanders. *Science* 164 (1969) 1290–1292.
- Adler, K., Pineal end organ: role in extraoptic entrainment of circadian locomotor rhythm in frogs, in: *Biochronometry*, p. 342–350. Ed. M. Menaker. National Academy of Sciences, Washington, D.C. 1971.
- Adler, K., and Taylor, D. H., Melatonin and thyroxine: influence on compass orientation in salamanders. *J. comp. Physiol.* 136 (1980) 235–241.
- Anderson, K., and Turek, F., The influence of blinding and removal of the frontal organ on the activity of *Xenopus laevis*. *Am. Zool.* 23 (1983) 884.
- Bassi, C. J., and Powers, M. K., Circadian rhythm in goldfish visual sensitivity. *Invest. Ophthalmol. vis. Sci.* 28 (1987) 1811–1815.
- Bernstein, S. A., Breeding, D. J., and Fisher, S. K., The influence of light on cone disk shedding in the lizard, *Sceloporus occidentalis*. *J. Cell Biol.* 99 (1984) 379–389.
- Besharse, J. C., and Iuvone, P. M., Circadian clock in *Xenopus* eye controlling retinal serotonin N-acetyltransferase. *Nature* 305 (1983) 133–135.

- Birks, E. K., and Ewing, R. D., Photoperiod effects on hydroxyindole-O-methyltransferase activity in the pineal gland of chinook salmon (*Oncorhynchus tshawytscha*). *Gen. comp. Endocr.* 43 (1981) 277–283.
- Birks, E. K., and Ewing, R. D., Seasonal changes in pineal melatonin content and hydroxyindole-O-methyltransferase activity in juvenile chinook salmon, *Oncorhynchus tshawytscha*. *Gen. comp. Endocr.* 64 (1986) 91–98.
- Dearry, A., and Barlow, R. B. Jr., Circadian rhythms in the green sunfish retina. *J. gen. Physiol.* 89 (1987) 745–770.
- Deguchi, T., A circadian oscillator in cultured cells of chicken pineal gland. *Nature* 282 (1979) 94–96.
- Demian, J. J., and Taylor, D. H., Photoreception and locomotor rhythm entrainment by the pineal body of the newt, *Notophthalmus viridescens* (Amphibia, Urodela, Salamandridae). *J. Herpet.* 11 (1977) 131–139.
- Eriksson, L. O., Tagesperiodik geblendeter Bachsaiblinge. *Naturwissenschaften* 59 (1972) 219–220.
- Eriksson, L. O., Die Jahresperiodik augen- und pinealorganloser Bachsaiblinge *Salvelinus fontinalis* Mitchell. *Aquilo Ser. Zool.* 13 (1972) 8–12.
- Falcón, J., and Collin, J.-P., In vitro uptake and metabolism of [³H] indole compounds in the pineal organ of the pike. II. A radioautographic study. *J. Pineal Res.* 2 (1985) 357–373.
- Falcón, J., Guerlotti, J. F., Voisin, P., and Collin, J.-P., Rhythmic melatonin biosynthesis in a photoreceptive pineal organ: a study in the pike. *Neuroendocrinology* 45 (1987) 479–486.
- Firth, B. T., and Kennaway, D. J., Melatonin content of the pineal, parietal eye and blood plasma of the lizard, *Trachydosaurus rugosus*: effect of constant and fluctuating temperature. *Brain Res.* 404 (1987) 313–318.
- Firth, B. T., Kennaway, D. J., and Rozenblids, M. A. M., Plasma melatonin in the scincid lizard, *Trachydosaurus rugosus*: diel rhythm, seasonality, and the effect of constant light and constant darkness. *Gen. comp. Endocr.* 37 (1979) 493–500.
- Garg, S. K., and Sundararaj, B. I., Role of pineal in the regulation of some aspects of circadian rhythmicity in the catfish, *Heteropneustes fossilis* (Bloch). *Chronobiologia* 13 (1986) 1–11.
- Gern, W. A., and Greenhouse, S. S., Examination of in vitro melatonin secretion from superfused trout (*Salmo gairdneri*) pineal organs maintained under diel illumination or continuous darkness. *Gen. comp. Endocr.* 71 (1988) 163–174.
- Gern, W. A., and Norris, D. O., Plasma melatonin in the neotenic tiger salamander (*Ambystoma tigrinum*): effects of photoperiod and pinealectomy. *Gen. comp. Endocr.* 38 (1979) 393–398.
- Gern, W. A., Norris, D. O., and Duvall, D., The effect of light and temperature on plasma melatonin in neotenic tiger salamanders (*Ambystoma tigrinum*). *J. Herpet.* 17 (1983) 228–234.
- Gern, W. A., Owens, D. W., and Ralph, C. L., Persistence of the nyctohemeral rhythm of melatonin secretion in pinealectomized or optic

- tract-sectioned trout (*Salmo gairdneri*). J. exp. Zool. 205 (1978) 371–376.
- 24 Goudie, C. A., Davis, K. B., and Simco, B. A., Influence of the eyes and pineal gland on locomotor activity patterns of channel catfish *Ictalurus punctatus*. Physiol. Zool. 56 (1983) 10–17.
 - 25 Guerlotti, J., Falcón, J., Voisin, P., and Collin, J.-P., Indoles in the photoreceptor cells of the lamprey pineal complex. Annls Endocr., Paris 47 (1986) 62–64.
 - 26 Hamasaki, D. I., and Eder, D. J., Adaptive radiation of the pineal system, in: Handbook of Sensory Physiology, p. 497–548. Ed. F. Crescitelli. Springer, New York 1977.
 - 27 Illnerová, H., and Vanček, J., Circadian rhythm in inducibility of rat pineal N-acetyltransferase after brief light pulses at night: control by a morning oscillator. J. comp. Physiol. A154 (1984) 739–744.
 - 28 Janik, D. S., Circadian organization in the desert iguana. Ph.D. dissertation. University of Oregon, Eugene 1987.
 - 29 Joss, J. M. P., A rhythm in hydroxyindole-O-methyltransferase (HIOMT) activity in the scincid lizard, *Lampropholis guichenoti*. Gen. comp. Endocr. 36 (1978) 521–525.
 - 30 Kavaliers, M., The pineal organ and circadian organization of teleost fish. Rev. Can. Biol. 38 (1979) 281–292.
 - 31 Kavaliers, M., Pineal involvement in the control of circadian rhythmicity in the lake chub, *Couesius plumbeus*. J. exp. Zool. 209 (1979) 33–40.
 - 32 Kavaliers, M., Circadian locomotor activity rhythms of the Turbot, *Lota lota*: seasonal differences in period length and the effect of pinealectomy. J. comp. Physiol. 136 (1980) 215–218.
 - 33 Kavaliers, M., Retinal and extraretinal entrainment action spectra for the activity rhythms of the lake chub, *Couesius plumbeus*. Behav. neural Biol. 30 (1980) 56–67.
 - 34 Kavaliers, M., Circadian organization in white suckers *Catostomus commersoni*: the role of the pineal organ. Comp. Biochem. Physiol. 68A (1981) 127–129.
 - 35 Lythgoe, J. N., and Shand, J., Endogenous circadian retinomotor movements in the neon tetra (*Paracheirodon innesi*). Invest. Ophthalm. vis. Sci. 24 (1983) 1203–1210.
 - 36 McNulty, J. A., Functional morphology of the pineal complex in cyclostomes, elasmobranchs, and bony fishes, in: Pineal Research Reviews, vol. 2, p. 1–40. Ed. R. J. Reiter. Alan R. Liss, Inc., New York 1984.
 - 37 Menaker, M., Eyes – the second (and third) pineal glands?, in: Photoperiodism, Melatonin and the Pineal, p. 78–87. Eds D. Evered and S. Clark. Pitman, London 1985.
 - 38 Menaker, M., and Wisner, S., Temperature-compensated circadian clock in the pineal of *Anolis*. Proc. natl Acad. Sci. USA 80 (1983) 6119–6121.
 - 39 Moore, R. Y., and Card, J. P., Visual pathways and the entrainment of circadian rhythms, in: The Medical and Biological Effects of Light, p. 123–133. Eds R. J. Wurtman, M. J. Baum and J. T. Potts, Jr. New York Academy of Sciences, New York 1985.
 - 40 Ooka-Souda, S., and Kabasawa, H., Circadian rhythms in locomotor activity of the hagfish, *Eptatretus burgeri* III. Hypothalamus: a locus of the circadian pacemaker? Zool. Sci. 5 (1988) 437–442.
 - 41 Pang, S. F., Shiu, S. Y. W., and Tse, S. F., Effect of photic manipulation on the level of melatonin in the retinas of frogs (*Rana tigrina regulosa*). Gen. comp. Endocr. 58 (1985) 464–470.
 - 42 Pierce, M. E., and Besharse, J. C., Melatonin and dopamine interactions in the regulation of rhythmic photoreceptor metabolism, in: Pineal and Retinal Relationships, p. 219–237. Eds P. J. O'Brien and D. C. Klein. Academic Press, New York 1986.
 - 43 Pittendrigh, C. S., On the mechanism of the entrainment of a circadian rhythm by light cycles, in: Circadian Clocks, p. 277–297. Ed. J. Aschoff. North-Holland, Amsterdam 1965.
 - 44 Prasada Rao, P. D., and Sharma, S. C., Retinofugal pathways in juvenile and adult channel catfish, *Ictalurus (Ameiurus) punctatus*: an HRP and autoradiographic study. J. comp. Neurol. 210 (1982) 37–48.
 - 45 Quay, W. B., The parietal eye-pineal complex, in: Biology of the Reptilia, vol. 9, p. 245–406. Ed. C. Gans. Academic Press, New York 1979.
 - 46 Ralph, C. L., Melatonin production by extrapineal tissues, in: Melatonin: Current Status and Perspectives, p. 35–46. Eds N. Birau and W. Schlot. Pergamon Press, New York 1981.
 - 47 Reiter, R. J., Action spectra, dose-response relationships, and temporal aspects of light's effects on the pineal gland, in: The Medical and Biological Effects of Light, p. 215–230. Eds R. J. Wurtman, M. J. Baum and J. T. Potts, Jr. New York Academy of Sciences, New York 1985.
 - 48 Repérant, J., Rio, J. P., Miceli, D., and Lemire, M., A radioautographic study of retinal projections in type I and type II lizards. Brain Res. 142 (1978) 401–411.
 - 49 Takahashi, J. S., and Menaker, M., Role of the suprachiasmatic nuclei in the circadian system of the house sparrow, *Passer domesticus*. J. Neurosci. 2 (1982) 815–828.
 - 50 Underwood, H., Circadian organization in lizards: the role of the pineal organ. Science 195 (1977) 587–589.
 - 51 Underwood, H., Circadian organization in the lizard *Sceloporus occidentalis*: the effects of pinealectomy, blinding, and melatonin. J. comp. Physiol. 141 (1981) 537–547.
 - 52 Underwood, H., Circadian pacemakers in lizards: phase-response curves and effects of pinealectomy. Am. J. Physiol. 244 (1983) R857–R864.
 - 53 Underwood, H., Circadian organization in the lizard *Anolis carolinensis*: a multioscillator system. J. comp. Physiol. A 152 (1983) 265–274.
 - 54 Underwood, H., Pineal melatonin rhythms in the lizard *Anolis carolinensis*: effects of light and temperature cycles. J. comp. Physiol. A 157 (1985) 57–65.
 - 55 Underwood, H., Extraretinal photoreception in the lizard *Sceloporus occidentalis*: phase response curve. Am. J. Physiol. 248 (1985) R407–R414.
 - 56 Underwood, H., Circadian rhythms in lizards: phase response curve for melatonin. J. Pineal Res. 3 (1986) 187–196.
 - 57 Underwood, H., Light at night cannot suppress pineal melatonin levels in the lizard *Anolis carolinensis*. Comp. Biochem. Physiol. 84A (1986) 661–663.
 - 58 Underwood, H., and Calaban, M., Pineal melatonin rhythms in the lizard *Anolis carolinensis*: I. Response to light and temperature cycles. J. Biol. Rhythms 2 (1987) 179–193.
 - 59 Underwood, H., and Gross, G., Vertebrate circadian rhythms: retinal and extraretinal photoreception. Experientia 38 (1982) 1013–1021.
 - 60 Underwood, H., and Harless, M., Entrainment of the circadian activity rhythm of a lizard to melatonin injections. Physiol. Behav. 35 (1985) 267–270.
 - 61 Underwood, H., and Menaker, M., Extraretinal photoreception in lizards. Photochem. Photobiol. 23 (1976) 227–243.
 - 62 van Veen, T., Hartwig, H. G., and Muller, K., Light-dependent motor activity and photonegative behavior in the eel (*Anguilla anguilla* L.). Evidence for extraretinal and extrapineal photoreception. J. comp. Physiol. 111 (1976) 209–219.
 - 63 Vivien-Roels, B., Arendt, J., and Bradtke, J., Circadian and circannual fluctuations of pineal indoleamines (serotonin and melatonin) in *Tesudo hermanni* Gmelin (Reptilia, Chelonia) 1. Under natural conditions of photoperiod and temperature. Gen. comp. Endocr. 37 (1979) 197–210.
 - 64 Vivien-Roels, B., and Pévet, P., The pineal gland and the synchronization of reproductive cycles with variations of the environmental climatic conditions, with special reference to temperature, in: Pineal Research Reviews, vol. 1, p. 91–143. Ed. R. J. Reiter. Alan R. Liss, Inc., New York 1983.
 - 65 Vivien-Roels, B., Pévet, P., and Claustrat, B., Pineal and circulating melatonin rhythms in the box turtle, *Terrapene carolina triunguis*: effect of photoperiod, light pulse, and environmental temperature. Gen. comp. Endocr. 69 (1988) 163–173.
 - 66 Vivien-Roels, B., Pévet, P., Dubois, M. P., Arendt, J., and Brown, G. M., Immunohistochemical evidence for the presence of melatonin in the pineal gland, the retina and the Harderian gland. Cell Tissue Res. 217 (1981) 105–115.
 - 67 Wiechmann, A. F., Melatonin: parallels in pineal gland and retina. Exp. Eye Res. 42 (1986) 507–527.
 - 68 Wiechmann, A. F., Bok, D., and Horwitz, J., Melatonin binding in the frog retina: autoradiographic and biochemical analysis. Invest. Ophthalm. vis. Sci. 27 (1986) 153–163.