

## Studies on Mesoionic Compounds. IX.<sup>1)</sup> Synthesis of Bicyclic Mesoionic Compounds

KATSUTADA MASUDA, JUN ADACHI, TATSUMI SHIBATA, and KEIICHI NOMURA

*Faculty of Pharmaceutical Sciences, Toyama Medical and Pharmaceutical University<sup>2)</sup>*

(Received December 18, 1978)

Reaction of the mesoionic 4-amino-1,2,4-triazolium-3-thiolates (I) with cyanogen bromide gave bicyclic mesoionic compounds (III) related to the product (II) obtained on treatment of I with phosgene. Benzoylation, tosylation, and nitrosation at the exocyclic nitrogen of III were also investigated.


**Keywords**—mesoionic compound; 4-amino-1,2,4-triazolium derivative; cyanogen bromide; bicyclic mesoionic compound; nitrosation

It is known that thioacylhydrazines give mesoionic heterocycles on treatment with phosgene, thiophosgene,<sup>3)</sup> 2,2-dichloroacrylates,<sup>4)</sup> and dichlorides of imidic<sup>5)</sup> and acylimidic acids.<sup>6)</sup> Based on the observation that mesoionic 4-amino-1,5-diphenyl-1,2,4-triazolium-3-thiolate<sup>7)</sup> (Ia) possesses a moiety similar to that of thioacylhydrazine, Lazaris *et al.*<sup>8)</sup> synthesized the novel bicyclic mesoionic compounds, 1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-olate (IIa) and -2-benzoylamidine (Va), by the reaction of Ia with phosgene and benzoyl-carbonimidic dichloride, respectively. In connection with our studies on reactions of 4-amino-1,2,4-triazolium-3-thiolates (I),<sup>1)</sup> we synthesized similar bicyclic mesoionic compounds by an alternative procedure.

When the amino compounds (Ia, b)<sup>1,7,9)</sup> were treated with cyanogen bromide, the hydrobromides of the bicyclic mesoionic imino compounds (III) were obtained directly. It was considered that in this reaction the intermediate nitrile formed at the first stage of the reaction was cyclized with the aid of hydrogen bromide. These compounds (III) were characterized by reaction with benzoyl chloride, giving the chlorides of the N-benzoyl derivatives (IV). The salts (IV) readily gave the free mesoionic compounds (V) on treatment with alkali. In the infrared (IR) spectra, absorptions of the carbonyl stretching of the compounds (V) were at lower frequencies than those of the salts (IV). For example,  $\nu_{\text{C=O}}$  1560  $\text{cm}^{-1}$  for IVa and  $\nu_{\text{C=O}}$  1470  $\text{cm}^{-1}$  for Va were observed. N-*p*-Toluenesulfonyl derivatives (VI) were also prepared similarly. Nuclear magnetic resonance (NMR) and mass spectra of the above bicyclic compounds were consistent with these structures. Furthermore, the compounds (III) afforded N-nitroso derivatives (VII) upon treatment with sodium nitrite

- 1) Part VIII: K. Masuda, J. Adachi, T. Shibata, and K. Nomura, *Chem. Pharm. Bull.* (Tokyo), **27**, 1683 (1979).
- 2) Location: *Sugitani, Toyama 930-01, Japan.*
- 3) A.R. McCarthy, W.D. Ollis, and C.A. Ramsden, *Chem. Commun.*, **1968**, 499; K.T. Potts and C. Sapino, *ibid.*, **1968**, 672; R. Grashey, M. Baumann, and W.D. Lubos, *Tetrahedron Lett.*, **1968**, 5877, 5881.
- 4) R. Grashey, M. Baumann, and R. Hamprecht, *Tetrahedron Lett.*, **1970**, 5083.
- 5) W.D. Ollis and C.A. Ramsden, *Chem. Commun.*, **1971**, 1222.
- 6) R. Grashey, R. Hamprecht, N. Keramaris, and M. Baumann, *Tetrahedron Lett.*, **1972**, 2943; A. Ya. Lazaris, S.M. Shmuilovich, and A.N. Egorochkin, *Khim. Geterotsikl. Soedin.*, **1973**, 1345 [*C.A.*, **80**, 27175 (1974)].
- 7) A. Ya. Lazaris, S.M. Shmuilovich, and A.N. Egorochkin, *Zh. Org. Khim.*, **8**, 2621 (1972) [*C.A.*, **78**, 72015 (1973)].
- 8) A. Ya. Lazaris, S.M. Shmuilovich, and A.N. Egorochkin, *Zh. Org. Khim.*, **10**, 2236 (1974) [*C.A.*, **82**, 43270 (1975)].
- 9) W.D. Ollis and G. Rawson, *J.C.S. Chem. Commun.*, **1976**, 440.

in acetic acid. It is known that pyrolysis of N-nitroso-sydnone imine<sup>10)</sup> and N-nitroso-1,2,3,4-oxatriazolium-5-aminide<sup>11)</sup> give the corresponding oxo-type mesoionic compounds. Thus, in order to obtain II thermolysis of VII in several solvents was attempted. Though II was detected on thin layer chromatography, these reactions resulted in the formation of many complex materials. This may occur because the thiadiazole ring of II easily undergoes ring opening. For example, the oxo-type compounds (II) were changed to the monocyclic mesoionic esters (VIII) on heating in ethanol. A similar reaction of mesoionic 1,3,4-thiadiazolium-2-olate with aniline giving the semicarbazide derivative has been reported.<sup>12)</sup>



### Experimental<sup>13)</sup>

**2-Amino-5,6-diphenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium Bromide (IIIa)**—A mixture of 4-amino-1,5-diphenyl-1,2,4-triazolium-3-thiolate (Ia)<sup>1)</sup> (1.5 g) and BrCN (1.2 g) in CHCl<sub>3</sub> (35 ml) was stirred for 2.5 hr with cooling in an ice bath. After concentration, the oily residue was crystallized from ether, and recrystallized from CH<sub>3</sub>CN to give 840 mg (40%) of IIIa, colorless fine needles, mp > 280°. IR  $\nu_{\max}^{\text{KBr}}$  cm<sup>-1</sup>: 1600. UV  $\lambda_{\max}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 274 (4.31). MS  $m/e$ : 293 (M<sup>+</sup> - HBr). Anal. Calcd. for C<sub>15</sub>H<sub>12</sub>BrN<sub>5</sub>S: C, 48.14; H, 3.23; N, 18.71. Found: C, 48.17; H, 3.00; N, 18.61.

**2-Amino-5-methyl-6-phenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium Bromide (IIIb)**—A solution of BrCN (4 g) in CHCl<sub>3</sub> (40 ml) was added to a cooled solution of 5-methyl-1-phenyl-1,2,4-triazolium-3-thiolate (Ib)<sup>1)</sup> (1.5 g) in CHCl<sub>3</sub> (40 ml) with stirring. The mixture was allowed to stand at room temperature overnight. After evaporation, MeOH was added to the residue and the mixture was filtered. Concentration of the filtrate and recrystallization of the residue from MeOH-isopropyl alcohol gave 922 mg (40%) of IIIb, colorless needles, mp > 280°. IR  $\nu_{\max}^{\text{KBr}}$  cm<sup>-1</sup>: 3400—2800. UV  $\lambda_{\max}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 256 (4.21). NMR (CD<sub>3</sub>OD)  $\delta$ : 2.8 (3H, s, Me), 7.6 (5H, s, arom.). Anal. Calcd. for C<sub>10</sub>H<sub>10</sub>BrN<sub>5</sub>S: C, 38.53; H, 3.23; N, 22.47. Found: C, 38.72; H, 3.33; N, 22.49.

**5,6-Diphenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-benzoylamide (Va)**—A mixture of IIIa (100 mg) and benzoyl chloride (3 ml) was heated at 110° for 5 hr. After cooling, the precipitates were collected

10) L.E. Kholodov and V.G. Yashunsky, *Dokl. Akad. Nauk SSSR*, **179**, 366 (1968) [*C.A.*, **69**, 106607 (1968)].

11) C. Christophersen and S. Trepndahl, *Acta Chem. Scand.*, **26**, 858 (1972).

12) R. Grashey, M. Baumann, and W.D. Lubos, *Tetrahedron Lett.*, **1968**, 5877.

13) All melting points are uncorrected. IR spectra were recorded on a JASCO IRA-1 spectrophotometer, and UV spectra were measured with a Hitachi 124 spectrophotometer. NMR spectra were obtained with a JEOL JNM-PMX-60 spectrometer with tetramethylsilane as an internal standard. Mass spectra (MS) were obtained with a JEOL JMS-01SG instrument.

and washed with ether to give 94 mg of IVa, mp 265—270° (dec.). IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1560. The above salt (IVa) was stirred in saturated aqueous  $\text{NaHCO}_3$ . Extraction with  $\text{CHCl}_3$  and usual work-up of the extract gave 40 mg (37%) of Va, pale yellow cubes, mp 254—259° ( $\text{CH}_3\text{CN}$ -isopropyl alcohol) (lit.<sup>8</sup>) mp 240—240.5°. IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1470. UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 285 (4.13), 329 (4.45). NMR ( $\text{CDCl}_3$ )  $\delta$ : 7.1—7.8 (13H, m, arom.), 8.25—8.6 (2H, m, arom.). MS  $m/e$ : 397 ( $\text{M}^+$ ). Anal. Calcd. for  $\text{C}_{22}\text{H}_{15}\text{N}_5\text{OS}$ : C, 66.48; H, 3.80; N, 17.62. Found: C, 66.51; H, 3.52; N, 17.36.

**5-Methyl-6-phenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-benzoylamide (Vb)**—A mixture of IIIb (50 mg) and benzoyl chloride (2 ml) was heated at 150° for 1 hr. After cooling, the precipitates were collected to give 43 mg of IVb. IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1540. The salt (IVb) was treated with  $\text{NaHCO}_3$  as described above to afford 25 mg (47%) of Vb, pale yellow prisms, mp 272—275° ( $\text{EtOH}$ - $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1450. UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 313 (4.49). NMR ( $\text{CDCl}_3$ )  $\delta$ : 2.8 (3H, s, Me); 7.0—7.6 (3H, m, arom.), 7.45 (5H, s, arom.), 8.0—8.2 (2H, m, arom.). MS  $m/e$ : 335 ( $\text{M}^+$ ). Anal. Calcd. for  $\text{C}_{17}\text{H}_{13}\text{N}_5\text{OS}$ : C, 60.88; H, 3.91; N, 20.88. Found: C, 60.96; H, 3.87; N, 20.62.

**5,6-Diphenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-*p*-toluenesulfonylamide (VIa)**—A solution of KOH (64 mg) in water (0.5 ml) was added to a mixture of IIIa (100 mg) and *p*-toluenesulfonyl chloride (100 mg) in water (3 ml). The mixture was then heated at 100° for 7 hr. After cooling, extraction with  $\text{CHCl}_3$  and usual work-up of the extract gave 32 mg (27%) of VIa, colorless powder, mp >280° ( $\text{EtOH}$ ). IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1500, 1140, 1085. UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 291 (4.31). MS  $m/e$ : 447 ( $\text{M}^+$ ). Anal. Calcd. for  $\text{C}_{22}\text{H}_{17}\text{N}_5\text{O}_2\text{S}_2$ : C, 59.04; H, 3.83; N, 15.65. Found: C, 59.33; H, 3.85; N, 15.46.

**5-Methyl-6-phenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-*p*-toluenesulfonylamide (VIb)**—A solution of KOH (40 mg) in water (0.5 ml) was added to a cooled mixture of IIIb (100 mg) and *p*-toluenesulfonyl chloride (100 mg) in water (3 ml), and the mixture was stirred at room temperature for 6 hr. The precipitates were collected and recrystallized from  $\text{EtOH}$  to give 56 mg (45.5%) of VIb, colorless powder, mp 260—265°. IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1500, 1130, 1080. UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 282 (4.40). MS  $m/e$ : 385 ( $\text{M}^+$ ). Anal. Calcd. for  $\text{C}_{17}\text{H}_{15}\text{N}_5\text{O}_2\text{S}_2$ : C, 52.97; H, 3.92; N, 18.17. Found: C, 52.86; H, 3.88; N, 18.16.

**5,6-Diphenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-nitrosoamide (VIIa)**—A solution of  $\text{NaNO}_2$  (140 mg) in water (0.5 ml) was added dropwise to a cooled solution of IIIa (432 mg) in  $\text{AcOH}$  (15 ml) and water (5 ml) at 0—5°. The mixture was stirred for 6 hr, and the resulting precipitates were collected and recrystallized from  $\text{EtOH}$ - $\text{CHCl}_3$  to give 250 mg (66%) of VIIa, fine yellow needles, mp 156—157° (dec.). IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1510, 1380. UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 319 (4.29). Anal. Calcd. for  $\text{C}_{15}\text{H}_{10}\text{N}_6\text{OS}$ : C, 55.89; H, 3.13; N, 26.07. Found: C, 55.76; H, 3.34; N, 25.97.

**5-Methyl-6-phenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-nitrosoamide (VIIb)**—A solution of  $\text{NaNO}_2$  (60 mg) in water (0.3 ml) was added to a cooled solution of IIIb (100 mg) in  $\text{AcOH}$  (3 ml) and water (1 ml). The mixture was kept in a refrigerator overnight. The precipitates were collected and recrystallized from  $\text{MeOH}$ - $\text{CH}_3\text{CN}$  to give 33 mg (40%) of VIIb, yellow plates, mp 180—181°. IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1520, 1380. UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 312 (4.03). Anal. Calcd. for  $\text{C}_{10}\text{H}_8\text{N}_6\text{OS}$ : C, 46.15; H, 3.10; N, 32.29. Found: C, 46.25; H, 3.11; N, 32.01.

**5,6-Diphenyl- and 6-Methyl-5-phenyl-1,2,4-triazolo[3,4-*b*]-1,3,4-thiadiazolium-2-olates (IIa and IIc)**—Using the reported method,<sup>8</sup> IIa and IIc were prepared as follows. A solution of  $\text{COCl}_2$  (1.6 g) in  $\text{CHCl}_3$  (5 ml) was added to a solution of Ia, c<sup>1</sup> (5 mmol) in  $\text{CHCl}_3$  (5 ml) at 0—5°. The mixture was refluxed for 1 hr, and the precipitates were filtered off. Concentration of the filtrate and recrystallization of the residue gave IIa, c.

IIa: 880 mg (60%), colorless cubes, mp 192—194° (benzene) (lit.<sup>9</sup>) mp 181°. IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1680. UV  $\lambda_{\text{max}}^{\text{EtOH}}$  nm (log  $\epsilon$ ): 280 (4.15). NMR ( $\text{CDCl}_3$ )  $\delta$ : 7.0—8.0 (m, arom.). MS  $m/e$ : 294 ( $\text{M}^+$ ). Anal. Calcd. for  $\text{C}_{15}\text{H}_{10}\text{N}_4\text{OS}$ : C, 61.21; H, 3.42; N, 19.04. Found: C, 61.03; H, 3.25; N, 18.92.

IIc: 1.1 g (87%), colorless needles, mp 154—156° (isopropyl alcohol). IR  $\nu_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 1680. NMR ( $\text{CDCl}_3$ )  $\delta$ : 4.1 (3H, s, Me), 7.3—8.0 (5H, m, arom.). Anal. Calcd. for  $\text{C}_{10}\text{H}_8\text{N}_4\text{OS}$ : C, 51.71; H, 3.47; N, 24.12. Found: C, 51.48; H, 3.39; N, 24.20.

**Alcoholysis of II with Ethanol**—a) A suspension of IIa (1 g) in  $\text{EtOH}$  (5 ml) was refluxed for 1 hr. After cooling, the precipitates were collected and recrystallized from  $\text{EtOH}$  to give 670 mg (58%) of VIIIa, colorless cubes, mp 199—200°. This product was identical with authentic VIIIa<sup>1</sup> (mp 199—201°) by comparison of IR and NMR spectra.

b) A mixture of IIc (100 mg) and  $\text{EtOH}$  (3 ml) was refluxed for 30 min. Removal of the solvent by evaporation, followed by recrystallization of the residue from isopropyl alcohol gave 98 mg (82%) of VIIIc, colorless sticks, mp 94—95°. The IR spectrum of this product was identical with that of authentic VIIIc<sup>1</sup> (mp 94—96°).

**Acknowledgement** The authors are indebted to Mr. M. Morikoshi and Mr. H. Hori of this University for mass spectral measurements and microanalyses, respectively.