

Communications to the Editor

[Chem. Pharm. Bull.]
30(5)1925—1928(1982)


1,6-DIHYDRO-3(2*H*)-PYRIDINONES AS SYNTHETIC INTERMEDIATES.
STEREOSELECTIVE SYNTHESIS OF (±)-CORYNANTHEIDOL AND (±)-QUININE

Takeshi Imanishi, Makoto Inoue, Yasuaki Wada, and Miyoji Hanaoka*
Faculty of Pharmaceutical Sciences, Kanazawa University
Takara-machi, Kanazawa 920, Japan


A new synthetic method is described for the *cis*-3,4-disubstituted piperidine compounds starting from ethyl 1,6-dihydro-3(2*H*)-pyridinone-1-carboxylate (**1**). Total synthesis of (±)-corynantheidol (**3**) and formal synthesis of (±)-quinine (**4**) were accomplished by an application of this method.

KEYWORDS — dihydropyridinone; Claisen rearrangement; hydroboration-oxidation; oxymercuration-demercuration; *cis*-3,4-disubstituted piperidine; meroquinene; indole alkaloid; corynantheidol; quinine

Since the first preparation of *N*-substituted 1,6-dihydro-3(2*H*)-pyridinone (**1**) was reported,¹⁾ we have been demonstrating various alkaloid syntheses using the dihydropyridinone as a common synthon.²⁾ At present we are beginning to synthesize many alkaloids and related compounds bearing the *cis*-3,4-disubstituted piperidine moiety, e.g. reserpine, corynantheidine, and quinine, utilizing the same synthon. Here we wish to present a new method for construction of the piperidine (**2**) with functionalized *cis*-dialkyl pendants at 3- and 4-position, and its application to a new stereoselective synthesis of (±)-corynantheidol (**3**) and (±)-quinine (**4**).




It had been found that ethyl 3-(2-ethylenedioxyethyl)-1,2,3,6-tetrahydropyridine-1-carboxylate (**5**) gave the 3-hydroxy compound (**6**) along with a small amount of the 4-hydroxy one (**7**) by the hydroboration-oxidation process.^{2b)} Oxymercuration-demercuration of **5**, however, yielded an approximately equal amount of **6** and **7**, and furthermore, epoxidation of **5** followed by reduction with diborane afforded **7** as a major isomer in 49% yield (see Table). Replacement of the nitrogen protective group of **7** from ethyl ester into benzyl ester was performed by basic hydrolysis and subsequent treatment with carbobenzoxy chloride, yielding **8** from **5** in overall yield of 32%. The compound **8** was also obtained in overall 36% yield by initial conversion of **5** into the benzyl urethane (**10**) *via* the amine (**9**) and subsequent epoxidation followed by reduction with diborane.

Table. Hydration of the Olefin (5) into 6 and 7

Entry	Conditions	Products (%)	
		<u>6</u>	<u>7</u>
<u>1</u> 2b)	1) B ₂ H ₆ /THF 0°C 3 h 2) H ₂ O ₂ , aq. NaOH 0°C 3 h	52	13
<u>2</u>	1) Hg(OAc) ₂ /THF-H ₂ O r.t. 13 h 2) NaBH ₄ , aq. NaOH r.t. 30 min	40	38
<u>3</u>	1) MCPBA/CH ₂ Cl ₂ r.t. 15 h 2) B ₂ H ₆ , NaBH ₄ /THF r.t. 3 h	32	49


Oxidation of 8 with PCC³⁾ afforded the ketone (12), which was condensed with the ylid of triethyl phosphonoacetate in benzene to give the unsaturated ester (13) [60% yield from 8; mp 78-80°C]. Catalytic hydrogenation of 13 over platinum oxide in ethyl acetate gave the saturated ester (14) as a single stereoisomer in 93% yield. From the inspection of the Dreiding model it is clear that the hydrogenation proceeded exclusively from the side opposite to the adjacent bulky



substituent to give only the *cis*-3,4-disubstituted piperidine as depicted in 14. The acetal group of 14 was hydrolyzed with 1% hydrochloric acid in acetone to give the aldehyde (15), which was reduced with sodium borohydride in methanol into the alcohol (16) in 91% yield from 14. Mesylation of 16 followed by reduction with zinc-sodium iodide in boiling 1,2-dimethoxyethane⁴⁾ furnished 18 via 17 in 71% yield. Hydrogenolysis of 18 over 5% palladium on carbon in ethanol yielded the amine (19) [84% yield; ν 3300, 1720; δ 0.90 (3H, t, $J = 6$ Hz), 1.23 (3H, t, $J = 7$ Hz), 2.00 (1H, s), 2.23 (2H, d, $J = 1.5$ Hz), 4.06 (2H, q, $J = 7$ Hz)], which was found to be identical with (+)-cincholoipon ethyl ester⁵⁾ by means of TLC and IR comparison. In this stage the stereochemistry of the product (14) obtained on the hydrogenation of 13 was completely confirmed.

On condensation with tryptophyl bromide⁶⁾ in dioxane the secondary amine (19) was converted into the tertiary amine (20) in 88% yield. Oxidative cyclization⁷⁾ of 20 using mercuric acetate-EDTA·2Na and subsequent sodium borohydride reduction yielded an inseparable mixture of the cyclized products (21 and 22) in total 37% yield. The mixture was reduced with lithium aluminum hydride in ether to furnish (\pm)-corynantheidol (3) [52% yield; mp 162-164°C; ν 3470, 3300, 2790, 2730; δ 0.92 (3H, t, $J = 6$ Hz), 3.75 (2H, t, $J = 7$ Hz), 7.0-7.5 (4H), 7.88 (1H, br s); m/e 298 (M^+), 297 (base), 225, 184, 170] and its structural isomer⁸⁾ (23) [34% yield; mp 137-140°C; ν 3460, 3300, 2800, 2760; δ 0.83 (3H, t, $J = 7$ Hz), 3.47 (1H, s), 3.78 (2H, t, $J = 7$ Hz), 7.0-7.5 (4H), 7.59 (1H, br s); m/e 298 (M^+), 297 (base), 253, 197, 184, 170]. The synthetic (\pm)-corynantheidol was proved by the spectral comparison to be identical with the authentic sample.

On the other hand, hydrogenation of 13 over 5% palladium on carbon in ethanol gave the saturated amine (24) as a sole isomer in 78% yield. The benzamide (25), prepared in the usual manner from 24 in 99% yield, was subjected to an acidic hydrolysis to afford the aldehyde (26), which was reduced with sodium borohydride in ethanol to the alcohol (27) [97% yield from 24; ν 3400, 1720, 1615; δ 1.25 (3H, t, $J = 7$ Hz), 2.03 (1H, s), 2.31 (2H, br), 4.13 (2H, q, $J = 7$ Hz), 7.39 (5H, s); m/e 319 (M^+), 105 (base)]. Chlorination of 27 with phosphorus oxychloride in pyridine followed by basic hydrolysis provided the carboxylic acid (29) [79% yield; mp 110-114°C; ν 3600-2400, 1705, 1615; δ 2.27 (2H, br), 7.27 (5H, s), 9.91 (1H, s); m/e 309 (M^+), 105 (base)]. According to the known method,⁹⁾ 29 was converted into



(±)-*N*-benzoylmeroquinene methyl ester (**31**) [56% yield; ν 1725, 1616; δ 2.26 (2H, br), 3.66 (3H, s), 4.9-5.2 (2H, m), 5.79 (1H, ddd, $J = 16, 11, 9$ Hz), 7.34 (5H, s); m/e 287 (M^+), 105 (base)], which was proved by TLC, IR, and 1H -NMR comparison to be identical with (+)-*N*-benzoylmeroquinene methyl ester.¹⁰⁾ Since the ester (**31**) has already been transformed into (±)-quinine (**4**),¹¹⁾ the present synthesis of **31** constitutes a formal synthesis of (±)-quinine (**4**).

Thus, we have established a novel and stereoselective method for construction of the piperidine bearing functionalized alkyl substituents at the 3- and 4-positions with the *cis* relationship. By applying the method to alkaloid syntheses, (±)-corynantheidol (**3**) and (±)-quinine (**4**) have been synthesized. The successive synthetic studies on more complex alkaloids are now in progress.

ACKNOWLEDGEMENT We wish to thank Prof. T. Fujii, Kanazawa University for a gift of (+)-cincholoipon ethyl ester, Prof. S. Takano, Tohoku University, for sending the spectra of (±)-corynantheidol and (±)-3-epicorynantheidol, and Dr. M. R. Uskoković, Hoffmann-La Roche Inc., for supplying (±)-meroquinene *tert*-butyl ester *d*-tartarate.

REFERENCES AND NOTES

- 1) T. Imanishi, I. Imanishi, and T. Momose, *Syn. Commun.*, **8**, 99 (1978).
- 2) a) T. Imanishi, H. Shin, M. Hanaoka, T. Momose, and I. Imanishi, *Heterocycles*, **14**, 1111 (1980); b) T. Imanishi, H. Shin, N. Yagi, and M. Hanaoka, *Tetrahedron Lett.*, **21**, 3285 (1980); c) T. Imanishi, N. Yagi, and M. Hanaoka, *ibid.*, **22**, 667 (1981); d) T. Imanishi, A. Nakai, N. Yagi, and M. Hanaoka, *Chem. Pharm. Bull.*, **29**, 901 (1981); e) T. Imanishi, N. Yagi, H. Shin, and M. Hanaoka, *Tetrahedron Lett.*, **22**, 4001 (1981); f) T. Imanishi, Y. Wada, M. Inoue, and M. Hanaoka, *Heterocycles*, **16**, 2133 (1981); g) T. Imanishi, K. Miyashita, A. Nakai, M. Inoue, and M. Hanaoka, *Chem. Pharm. Bull.*, in press.
- 3) E.J. Corey and J.W. Suggs, *Tetrahedron Lett.*, **1975**, 2647.
- 4) Y. Fujimoto and T. Tatsuno, *ibid.*, **1976**, 3325.
- 5) V. Prelog and E. Zalan, *Helv. Chim. Acta*, **27**, 535 (1944); T. Fujii, S. Yoshifuji, and M. Tai, *Chem. Pharm. Bull.*, **23**, 2094 (1975).
- 6) J.L. Neumeyer, U.V. Moyer, and J.E. Leonard, *J. Med. Chem.*, **12**, 450 (1969).
- 7) J. Gutzwiller, G. Pizzolato, and M. Uskoković, *J. Am. Chem. Soc.*, **93**, 5907 (1971); P.A. Wender, J.M. Schaus, and A.W. White, *ibid.*, **102**, 6157 (1980).
- 8) Its structure could be assigned as depicted from the facts that **23** was found to be not identical with (±)-3-epicorynantheidol and showed the Bohlmann bands in its IR spectrum.
- 9) M.R. Uskoković, T. Henderson, C. Reese, H.L. Lee, G. Grethe, and J. Gutzwiller, *J. Am. Chem. Soc.*, **100**, 571 (1978).
- 10) (+)-*N*-Benzoylmeroquinene methyl ester was derived from (+)-meroquinene *tert*-butyl ester *d*-tartarate according to the literature [M.R. Uskoković, D.L. Pruess, C.W. Despreaux, S. Shiuey, G. Pizzolato, and J. Gutzwiller, *Helv. Chim. Acta*, **56**, 2834 (1973)].
- 11) J. Gutzwiller and M. Uskoković, *J. Am. Chem. Soc.*, **92**, 203 (1970).

(Received April 21, 1982)