

EFFICIENT SOLID PHASE PEPTIDE SYNTHESIS ON A PHENACYL-RESIN BY A METHANESULFONIC ACID α -AMINO DEPROTECTING PROCEDURE¹⁾

Yoshiaki KISO,* Tooru KIMURA, Yoichi FUJIWARA, Hiroshi SAKIKAWA and Kenichi AKAJI

Department of Medicinal Chemistry, Kyoto Pharmaceutical University, Yamashina-ku, Kyoto 607, Japan

We have developed an efficient method for solid phase peptide synthesis which consists of N^α -selective deprotection by dilute methanesulfonic acid, *in situ* neutralization, and rapid coupling reaction using benzotriazol-1-yl-oxy-tris(dimethylamino)phosphonium hexafluorophosphate or new 2-(benzotriazol-1-yl)oxy-1,3-dimethylimidazolidinium hexafluorophosphate. This method was successfully used to synthesize several peptides using a new derivative, Boc-Tyr(Dpp) (Dpp: diphenylphosphinyl) and Boc-Arg(HCl) on a phenacyl-resin. For this method, we employed a fluoride ion final deprotection strategy based on a two-dimensional orthogonal protection scheme.

KEYWORDS solid phase peptide synthesis; fluoride ion final deprotection; *in situ* neutralization; methanesulfonic acid; coupling reagent; O-diphenylphosphinyl-tyrosine

We reported previously that the methanesulfonic acid (MSA) system [0.5 M MSA in dichloromethane-dioxane(9:1)]²⁻⁴⁾ is suitable for the removal of N^α -*tert*-butyloxycarbonyl (Boc) groups in solid phase peptide synthesis (SPPS), and is superior to trifluoroacetic acid (TFA) systems in terms of stability of semipermanent side chain-protecting groups and undesired pyroglutamyl formation from N-terminal glutamine in peptide-resin.³⁾ Herein we report an efficient method for solid phase peptide synthesis using this MSA deprotection system on an acid-stable phenacyl ester linkage (Pac)-resin⁵⁾ cleavable with fluoride ion.^{2,6)} This method consists of *in situ* neutralization and the rapid coupling reaction using benzotriazol-1-yl-oxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP)⁷⁾ or new 2-(benzotriazol-1-yl)oxy-1,3-dimethylimidazolidinium hexafluorophosphate (BOI)¹⁾ reagent activation. For this method, we used a fluoride ion final deprotection strategy based on a two-dimensional orthogonal protection scheme.^{2,4)}

First, we synthesized bradykinin potentiating peptide 5a (BPP5a; [Glu-Lys-Trp-Ala-Pro]⁸⁾) by this method. The termination in the amino acid Pac-resin occurs during N^α -deprotection and neutralization cycle of SPPS^{9,10)} (Fig. 1), and diketopiperadine formation and loss of loaded peptides from the Pac-resin are also observed.⁹⁾ These side reactions result from the high susceptibility of phenacyl linkage to aminolysis. To reduce the side reactions, we omitted the base-wash neutralization cycle using diisopropylethylamine (DIEA). After N^α -Boc deprotection by 0.5 M MSA, Boc-amino acid (2.5eq), BOP (2.5eq) and DIEA (4.5eq) were added to the N^α -deprotected resin, in which the α -amino group was masked as its MSA salt (Fig. 2). Then appropriate amounts of DIEA were added manually to adjust the reaction mixture to pH 7-8. After the coupling reaction, 0.3 M decanoic anhydride was used for capping. The protected BPP5a-resin [Glu-Lys(Fmoc)-Trp(Ppt)-Ala-Pro-O-CH₂CO-C₆H₄-resin] (Fmoc: 9-fluorenylmethyloxycarbonyl; Ppt: diphenylphosphinothiyl) was obtained from Boc-Ala-Pro-Pac-resin²⁾ using a




Fig. 1




Fig. 2. Coupling Pathway of Efficient SPPS Employing *in situ* Neutralization

Beckman 990E synthesizer at 25°C employing amino acid derivatives bearing side-chain protecting groups removable with fluoride ion in combination with the acid-labile N^α-Boc group, *i.e.*, Boc-Lys(Fmoc)²⁾ and Boc-Trp(Ppt).^{4,11)} The protected BPP5a-resin was deprotected by fluoride ion and purified as previously described.²⁾ The homogeneous peptide¹²⁾ was obtained in excellent yield (72% based on starting Boc-Ala-Pro-Pac-resin) compared with the previous synthesis using carbodiimide.²⁾

To apply this *in situ* neutralization method to the fully automated synthesis (Table), we employed an additional pyridine-wash (step 6) to remove the trace amount of free MSA remaining on the resin. After this treatment, 1 eq of MSA remains in the resin as the salt to mask the α-amino function. The protected BPP5a-resin was synthesized according to the Table using a Biosearch 9500 synthesizer, and pure BPP5a was obtained by the same manner (overall yield 72%). This *in situ* neutralization method reduced side reactions such as cyclization, because the terminal amino group was less exposed as a nucleophile by MSA masking and the rapid coupling reaction using BOP or BOI reagent.

By this method, we successfully synthesized Leu-enkephalin (Tyr-Gly-Gly-Phe-Leu) and neuromedin N (Lys-Ile-Pro-Tyr-Ile-Leu)¹³⁾ using a new Tyr derivative with the fluoride ion-labile O-diphenylphosphinyl (Dpp) group, Boc-Tyr(Dpp)-OH, [mp. 69-70°C; [α]_D¹⁷+16.2°(c=0.56, MeOH); satisfactory elemental analyses were obtained for C₂₆H₂₈NO₆P·0.5H₂O] prepared by the reaction of Boc-Tyr-OPac

Table. Schedule for Efficient Solid Phase Peptide Synthesis

1	DCM (x3)	0.3 min
2	0.5M MSA in DCM-dioxane(9:1)	1 min and 20 min
3	dioxane-DCM(1:2) (x3)	0.7 min
4	DCM-DMF(1:1) (x2)	0.3 min
5	DCM (x3)	0.3 min
6	2% pyridine in DCM (x2)	0.7 min
7	DCM (x4)	0.3 min
8	DCM-DMF(1:1)	1.5 min
9	Boc-amino acid(2.5eq), BOP or BOI(2.5eq) and DIEA(4.5eq) in DCM-NMP(1:2)	60min or 90 min
10	DCM-DMF(1:1) (x2)	0.3 min
11	DCM (x2)	0.3 min
12	if recoupling was necessary then go to step 8	
13	DCM-DMF(1:1) (x2)	0.3 min
14	0.3M decanoic anhydride in DCM-NMP(1:1)	30 min
15	DCM-DMF(1:1) (x2)	0.3 min

DCM: dichloromethane, DMF: *N,N*-dimethylformamide, NMP: *N*-methylpyrrolidone.




Fig. 3. A Two-Dimensional Orthogonal Protection Scheme for α -Neo-endorphin Synthesis by a Fluoride Ion Final Deprotection Strategy

and diphenylphosphinyl chloride (1.2 eq) in the presence of triethylamine (1.2 eq), followed by zinc-acetic acid treatment in DMF. The O-Dpp group was stable to such conditions as TFA (r.t., 24 h), 0.5 M MSA system (r.t., 24 h) and 5% DIEA in DMF (r.t., 24 h), while it could be readily removed with 0.1 M tetra-*n*-butylammonium fluoride trihydrate in DMF within 5 min at r.t. The protected peptide-resins¹⁴⁾ prepared from Boc-Leu-Pac-resin according to Table were treated with 0.5 M MSA system to remove the terminal Boc group, prior to fluoride ion final deprotection. The deprotected peptides were purified in the same way as BPP5a. The purified Leu-enkephalin and neuromedin N were obtained in 79% and 77% overall yield (based on the starting Boc-Leu-Pac-resin), respectively. The synthetic peptides had properties identical with authentic samples.

We applied this new method to the synthesis of an arginine-containing peptide, α -neo-endorphin (Tyr-Gly-Gly-Phe-Leu-Arg-Lys-Tyr-Pro-Lys),¹⁵⁾ using Boc-Arg(HCl)-OH, the guanidine group of which was protected by protonation. The fully protected α -neo-endorphin-resin¹⁶⁾ prepared from Boc-Lys(Fmoc)-Pac-resin according to the Table using a two-dimensional orthogonal protection scheme as shown in Fig. 3 was deprotected and purified as described above. The homogeneous α -neo-endorphin identical with an authentic sample was obtained in 46% yield [based on the starting Boc-Lys(Fmoc)-Pac-resin].

This method reduces side reactions such as cyclizations resulting from acid- or base-catalyzed intramolecular aminolysis. These excellent results show the potential of the method for SPPS on phenacyl ester linkage-resin in combination with fluoride ion final deprotection.

REFERENCES AND NOTES

- 1) Presented in part at the 26th Symposium on Peptide Chemistry, Oct. 24-26, 1988, Tokyo, Japan: "Peptide Chemistry 1988," p. 123, p. 129 and the 11th American Peptide Symposium, Jul. 9-14, 1989, La Jolla, California: Abstracts, P-329.
- 2) Y. Kiso, T. Kimura, Y. Fujiwara, M. Shimokura and A. Nishitani, *Chem. Pharm. Bull.*, **36**, 5024(1988).
- 3) Y. Kiso, A. Nishitani, M. Shimokura, Y. Fujiwara and T. Kimura, "Peptide Chemistry 1987," eds. T. Shiba and S. Sakakibara, Protein Res. Found., Osaka, 1988, p. 291.
- 4) Y. Kiso, T. Kimura, M. Shimokura and T. Narukami, *J. Chem. Soc., Chem. Commun.*, **1988**, 287.
- 5) F. Weygand and R. Obermeier, *Z. Naturforsch., Teil. B*, **23**, 1390(1968); T. Mizoguchi, K. Shigezane and N. Takamura, *Chem. Pharm. Bull.*, **18**, 1465(1970).
- 6) M. Ueki, K. Kai, M. Amemiya, H. Horino and H. Oyamada, *J. Chem. Soc., Chem. Commun.*, **1988**, 414.
- 7) B. Castro, J. R. Dormoy, E. Evin and C. Selve, *Tetrahedron Lett.*, **14**, 1219(1975); D. Le-Nguyen, A. Heitz and B. Castro, *J. Chem. Soc., Perkin Trans. 1*, **1987**, 1915.
- 8) S. H. Ferreira, D. C. Dartelt and T. L. Greene, *Biochemistry*, **9**, 2583(1970).
- 9) G. Barany and R. B. Merrifield, "The Peptides," Vol. 2, eds. E. Gross and J. Meienhofer, Academic Press, New York, 1980, p. 84; C. Birr, M. Wengert-Müller and A. Buku, "Peptides" Proc. 5th American Peptide Symposium, eds. M. Goodman and J. Meienhofer, Wiley, New York, 1977, p. 510.
- 10) J. P. Tam, W. R. Cunningham-Rundles, B. W. Erickson and R. B. Merrifield, *Tetrahedron Lett.*, **46**, 4001(1977).
- 11) Y. Kiso, M. Shimokura, T. Narukami, and T. Kimura, "Peptide Chemistry 1987," eds. T. Shiba and S. Sakakibara, Protein Res. Found., Osaka, 1988, p. 219.
- 12) D-Pro was not detected. In the case of the synthesis starting from Boc-Pro-Pac-resin, 26.7% of D-Pro was observed at the stage of dipeptide. Occurrence of racemization in proline phenacyl ester is described (S. Kubo, H. Kuroda, N. Chino, T. Kimura, and S. Sakakibara, Abstracts of 27th Symposium on Peptide Chemistry of Japan, 1989, p. 47).
- 13) N. Minamino, K. Kangawa and H. Matsuo, *Biochem. Biophys. Res. Commun.*, **122**, 542(1984).
- 14) Amounts of D-Leu were less than 0.2% and 1.7% at the stages of Phe-Leu and Ile-Leu, respectively.
- 15) K. Kangawa, N. Minamino, N. Chino, S. Sakakibara and H. Matsuo, *Biochem. Biophys. Res. Commun.*, **99**, 871(1981).
- 16) Amounts of D-Lys were less than 0.6% at the stage of dipeptide.

(Received October 23, 1989)