

Spiro-Substituted Piperidines as Neurokinin Receptor Antagonists. I. Design and Synthesis of (\pm)-*N*-[2-(3,4-Dichlorophenyl)-4-(spiro[isobenzofuran-1(3*H*),4'-piperidin]-1'-yl)butyl]-*N*-methylbenzamide, YM-35375, as a New Lead Compound for Novel Neurokinin Receptor Antagonists

Hirokazu KUBOTA,* Masahiro FUJII, Ken IKEDA, Makoto TAKEUCHI, Tadao SHIBANUMA, and Yasuo ISOMURA

Institute for Drug Discovery Research, Yamanouchi Pharmaceutical Co., Ltd., 21, Miyukigaoka, Tsukuba, Ibaraki 305, Japan. Received June 17, 1997; accepted October 13, 1997

Analysis of the structural requirements of compound 1 (SR48968), a potent NK₂ receptor antagonist, revealed that the 4-phenyl group of the piperidine is essential for binding with the NK₂ receptor and occupies an equatorial position. Energy calculation of a variety of substituted 4-phenyl piperidines revealed that spiro[isobenzofuran-1(3*H*),4'-piperidine] possesses a conformationally restricted equatorial phenyl group. Our compound 12 (YM-35375) possessing this spiro-substituted piperidine bound to the NK₂ receptor with an IC₅₀ value of 84 nM and to the NK₁ receptor with an IC₅₀ value of 710 nM. It showed more potent inhibitory activity (ID₅₀ 41 μg/kg (i.v.)) against [β-Ala⁸]-NKA(4–10)-induced bronchoconstriction in guinea pigs than (\pm)-SR48968 (ID₅₀ 68 μg/kg (i.v.)). YM-35375 may be a new lead compound for novel NK₂ receptor antagonists or NK₁-NK₂ dual antagonists.

Key words spiro[isobenzofuran-1(3*H*),4'-piperidine]; NK₂ antagonist; dual antagonist; YM-35375; SR48968

Neurokinins are peptides comprising ten or eleven amino acids and possess in common the sequence –Phe–X–Gly–Leu–Met–NH₂ at their C-termini. Among them, substance P (SP),¹ neurokinin A (NKA)² and neurokinin B (NKB),² isolated from mammals, exhibit a wide variety of biological responses³ through their receptors. Neurokinin receptors are now classified into the following three subtypes, NK₁, NK₂ and NK₃, which have high affinity for SP, NKA and NKB, respectively.⁴ Recently, Barnes *et al.* proposed that SP and NKA may play important roles in the pathogenesis of asthma.⁵ Namely, SP and NKA are released from the endings of sensory nerves by a variety of stimulants and induce the pathological features of asthma, such as microvascular leakage, mucus hypersecretion and bronchoconstriction.⁵ In the airway, SP causes microvascular leakage and mucus hypersecretion,⁶ and NKA induces bronchoconstriction.^{6,7} Based on these mechanisms, NK₁ and NK₂ receptor antagonists could prevent these functions in asthmatic patients and may be of clinical benefit for treatment of asthma.

Recently, various potent and selective non-peptide NK₁ receptor antagonists have been reported⁸ and their clinical efficacy has been evaluated. As regards non-peptide NK₂ receptor antagonists, only SR48968^{9,10} (**1**, Fig. 1) has been clinically evaluated to our knowledge. SR48968 has a characteristic structure, 4-acetamido-4-phenylpiperidine, which may be important for binding to the NK₂ receptor.

In this paper, we report the molecular design, synthesis and pharmacological properties of (\pm)-*N*-[2-(3,4-dichlorophenyl)-4-(spiro[isobenzofuran-1(3*H*),4'-piperidin]-1'-yl)butyl]-*N*-methylbenzamide (**12**, YM-35375) as a new lead compound for novel antiasthmatic drugs.

Chemistry

(\pm)-*N*-[2-(3,4-Dichlorophenyl)-4-(4-phenylpiperidino)butyl]-*N*-methylbenzamide (**2**) and (\pm)-*N*-[4-(4-acetamidopiperidino)-2-(3,4-dichlorophenyl)butyl]-*N*-methylbenzamide (**3**) were synthesized according to the literature.¹⁰

Spiro[isobenzofuran-1(3*H*),4'-piperidine] (**8**) was synthesized from 2-bromobenzyl alcohol (**4**) as shown in Chart 1. Compound **4** was converted to a dianion with *n*-butyllithium (*n*-BuLi)¹¹ in tetrahydrofuran (THF)-diethyl ether (Et₂O) and treated with 1-ethoxycarbonyl-4-piperidone (**5**) to give ethyl 4-hydroxy-4-(2-hydroxyethylphenyl)piperidine-1-carboxylate (**6**). The primary hydroxyl group of compound **6** was selectively tosylated by treatment with *p*-toluenesulfonyl chloride (TsCl), and




Fig. 1. Structure of SR48968 (**1**)




Chart 1

* To whom correspondence should be addressed.





Chart 2

Table 1. NK₂ Receptor Binding Affinities and Antagonistic Activities *in Vivo* of *N*-[2-(3,4-Dichlorophenyl)-4-(4-substituted piperidin-1'-yl)butyl]-*N*-methylbenzamides

	NK ₂ binding ^{a)} IC ₅₀ (nM)	NK ₁ binding ^{b)} IC ₅₀ (nM)	NK ₂ selectivity index ^{c)}	Bronchoconstriction in guinea pigs ^{d)} ID ₅₀ (μg/kg i.v.)
1 [(±)-SR48968]	4.1	> 1000	> 240	68
2	> 1000	N. T.	—	N. T.
3	180	N. T.	—	N. T.
12 (YM-35375)	84	710	8.5	41

^{a)} The binding affinities for hamster urinary bladder NK₂ receptor. See experimental section. ^{b)} The binding affinities for guinea pig urinary bladder NK₁ receptor. See experimental section. ^{c)} IC₅₀ to NK₁ receptor/IC₅₀ to NK₂ receptor. ^{d)} See experimental section. N. T.: Not tested.

the resultant tosylate was cyclized in the presence of pyridine to give the protected spiro-substituted piperidine (7). Compound 7 was treated with sodium hydroxide to give compound 8. Synthesis of YM-35375 is shown in Chart 2. (±)-4-Amino-3-(3,4-dichlorophenyl)butanol (9) was prepared according to the literature.¹²⁾ Compound 9 was treated with a mixture of formic acid and acetic anhydride (Ac₂O), followed by reduction with borane-THF complex to give the *N*-methyl derivative (10). Compound 10 was treated with benzoyl chloride in the presence of triethylamine (Et₃N), followed by hydrolysis with sodium hydroxide to give (±)-*N*-[2-(3,4-dichlorophenyl)-4-hydroxybutyl]-*N*-methylbenzamide (11). Treatment of compound 11 with methanesulfonyl chloride (MsCl), followed by substitution with compound 8 in the presence of Et₃N gave compound 12 (YM-35375).

Results and Discussion

Thus obtained compounds were evaluated for their binding affinities¹³⁾ to hamster urinary bladder NK₂ receptor and guinea pig urinary bladder NK₁ receptor.

As shown in Table 1, (±)-SR48968 exhibited high affinity for the NK₂ receptor with an IC₅₀ value of 4.1 nM. The removal of the phenyl group of 4-acetamido-4-phenylpiperidine (2) caused complete loss of the potency in contrast to the analog without the acetamido group (3), which retained the potency with an IC₅₀ value of 180 nM. These facts suggested that the phenyl group at the




Fig. 2. Putative 3-Dimensional Structures^{a)} of 4-Acetamido-1-methyl-4-phenylpiperidine (13) and 1'-Methylspiro[isobenzofuran-1(3*H*),4'-piperidine] (14)

^{a)} The 2-dimensional structures of these molecules were constructed using Sybyl (6.22) and converted to initial 3-dimensional structures with Concord 3.2.1. These initial structures were relaxed using the Tripos force field.

4-position of the piperidine may be crucial to bind to the NK₂ receptor and that the acetamido group may be necessary for increasing the affinity for the NK₂ receptor. To analyze the conformation of the 4-acetamido-4-phenylpiperidine moiety of SR48968, a putative 3-dimensional structure of 4-acetamido-1-methyl-4-phenylpiperidine (13), which was adopted as a simplified molecule, was constructed by energy minimization. As shown in Fig. 2A, the phenyl group was placed in an equatorial position and the acetamido group in an axial position. This conformation was 1.68 kcal/mol more stable than the other one with an axial phenyl group and an equatorial acetamido group. We speculated that the phenyl group may be immobilized in the equatorial position by

the axial acetamide group and that compounds with more restricted equatorial phenyl groups may show higher affinity for the NK₂ receptor. We designed 1'-methylspiro[isobenzofuran-1(3*H*),4'-piperidine] (**14**) as a conformationally restricted 4-phenylpiperidine. Conformational analysis of compound **14** suggested that the phenyl group lies in an equatorial position (Fig. 2B). This conformation was 2.51 kcal/mol more stable than the other one with an axial phenyl group. Based on these results, it was expected that YM-35375, which possesses this spiro-substituted piperidine instead of 4-acetamido-4-phenylpiperidine of SR48968, would bind to the NK₂ receptor with high affinity. Unfortunately, it was found that YM-35375 showed affinity for the NK₂ receptor with an IC₅₀ value of only 84 nM, which was a 20-fold decrease in potency relative to (±)-SR48968. One of the reasons for this decreased affinity for the NK₂ receptor was considered to be the lack of the acetamide group, which would affect the angle of the equatorial phenyl group and interact with the NK₂ receptor, *e.g.*, through hydrogen bonding.

As shown in Table 1, (±)-SR48968 inhibited [β-Ala⁸]-NKA(4—10)-induced bronchoconstriction in guinea pigs^{14,15} with an ID₅₀ value of 68 μg/kg (*i.v.*). In contrast to the binding assay, YM-35375 showed more potent inhibitory activity with an ID₅₀ value of 41 μg/kg (*i.v.*) than (±)-SR48968 in this model. These results suggested that the substitution of spiro[isobenzofuran-1(3*H*),4'-piperidine] for 4-acetamido-4-phenylpiperidine was favorable for NK₂ receptor-antagonistic activity *in vivo* and that YM-35375 may be a new lead compound for further research to find a novel NK₂ receptor antagonist. Furthermore, YM-35375 also showed weak NK₁ receptor affinity with an IC₅₀ value of 710 nM, and its NK₂ receptor selectivity index was 8.5 in contrast to over 240 of (±)-SR48968. This result suggested that the substitution of spiro[isobenzofuran-1(3*H*),4'-piperidine] for 4-acetamido-4-phenylpiperidine caused a change in selectivity and that the discovery of a novel NK₁-NK₂ dual antagonist¹⁶ might be possible by suitable modifications of YM-35375.

In conclusion, spiro[isobenzofuran-1(3*H*),4'-piperidine] was designed as a conformationally restricted piperidine with an equatorial phenyl group with the aim of obtaining a potent NK₂ receptor antagonist. Although YM-35375 which possessed this spiro-substituted piperidine was less potent than (±)-SR48968 in binding assay, it was more potent than (±)-SR48968 *in vivo*. As YM-35375 also showed weak NK₁ receptor affinity, we anticipate that further structural modifications of this compound may alter the selectivity between the NK₁ and NK₂ receptors and give rise to not only highly selective NK₂ receptor antagonists, but also NK₁-NK₂ dual antagonists. We are currently searching for novel neurokinin antagonists by utilizing YM-35375 as a lead compound. The results will be reported in the near future.

Experimental

All melting points were determined on a Yanagimoto MP-3 melting point apparatus without correction. ¹H-NMR spectra were taken on a JEOL JNM-EX400 spectrometer or a JEOL JNM-A500 spectrometer. Chemical shifts are given in ppm relative to that of Me₄Si (δ: 0) in CDCl₃ or dimethylsulfoxide-*d*₆ (DMSO-*d*₆) as an internal standard. The

abbreviations of signal patterns are as follows: s, singlet; brs, broad singlet; d, doublet; t, triplet; dd, double doublet; q, quartet; m, multiplet. Column chromatography was carried out on silica gel (Wakogel C-200 or Merck Silica gel 60). FAB-MS were obtained with a JEOL JMS-DX300 mass spectrometer, and electron impact (EI)-MS with a Hitachi M-80 mass spectrometer or a Hewlett-Packard 5890 GC-5970 MSD.

Ethyl 4-Hydroxy-4-(2-hydroxymethylphenyl)-1-piperidinecarboxylate (6) A 1.6 M solution of *n*-BuLi (705 ml, 1.13 mol) in hexane was added dropwise to a solution of 2-bromobenzyl alcohol (**4**) (101 g, 537 mmol) in THF (500 ml) and Et₂O (1000 ml) at -78 °C under an argon atmosphere. The mixture was stirred for 1 h at -78 °C, then 1-ethoxycarbonyl-4-piperidone (**5**) (101 g, 591 mmol) and THF (100 ml) were added at the same temperature. The mixture was stirred for 16 h at room temperature, H₂O was added and the whole was extracted with ethyl acetate (AcOEt). The extract was washed with saturated brine, dried over anhydrous MgSO₄ and concentrated *in vacuo*. The residue was purified by column chromatography (hexane:AcOEt = 1:1) to give the diol (**6**, 69.3 g, 46%) as a pale yellow oil. ¹H-NMR (CDCl₃) δ: 1.27 (3H, t, *J* = 7.1 Hz), 1.85—1.91 (2H, m), 1.96—2.04 (2H, m), 3.23—3.38 (2H, m), 3.98—4.07 (2H, m), 4.13 (2H, q, *J* = 7.1 Hz), 4.89 (2H, s), 7.20—7.30 (4H, m). EI-MS *m/z*: 279 (M⁺).

Ethyl Spiro[isobenzofuran-1(3*H*),4'-piperidine]-1'-carboxylate (7) A solution of TsCl (51.6 g, 271 mmol) in CH₂Cl₂ (160 ml) was added to a solution of compound **6** (68.7 g, 246 mmol), pyridine (43.7 ml, 541 mmol) and CH₂Cl₂ (800 ml) at 0 °C. The mixture was stirred for 18 h at room temperature, then pyridine (19.9 ml, 246 mmol) and TsCl (46.9 g, 246 mmol) were added at 0 °C. The whole was stirred for 22 h at room temperature. Further pyridine (19.9 ml, 246 mmol) and TsCl (46.9 g, 246 mmol) was added to the reaction mixture at 0 °C, and stirring was continued for an additional 18 h at room temperature. The mixture was diluted with CHCl₃ and washed with H₂O, 5% aqueous NaHCO₃, 10% aqueous citric acid and saturated brine. The organic layer was dried over anhydrous MgSO₄ and concentrated *in vacuo*. The residue was purified by column chromatography (hexane:AcOEt = 4:1) to give the carboxylate (**7**, 52.2 g, 81%) as a pale yellow solid. ¹H-NMR (CDCl₃) δ: 1.29 (3H, t, *J* = 7.4 Hz), 1.70—1.76 (2H, m), 1.79—1.89 (2H, m), 3.16—3.29 (2H, m), 4.06—4.22 (4H, m), 5.08 (2H, s), 7.07—7.10 (1H, m), 7.20—7.24 (1H, m), 7.26—7.30 (2H, m). EI-MS *m/z*: 261 (M⁺).

Spiro[isobenzofuran-1(3*H*),4'-piperidine] (8) A mixture of compound **7** (51.6 g, 197 mmol), 5N NaOH (100 ml) and EtOH (500 ml) was heated to reflux for 20 h, then concentrated *in vacuo*. The residue was diluted with brine and extracted with Et₂O. The organic layer was washed with saturated brine, dried over anhydrous MgSO₄ and concentrated *in vacuo* to give the piperidine (**8**, 33.0 g, 89%) as a colorless solid. ¹H-NMR (CDCl₃) δ: 1.71—1.77 (2H, m), 1.80—1.89 (2H, m), 1.99 (1H, s), 2.99—3.10 (4H, m), 5.07 (2H, s), 7.11—7.16 (1H, m), 7.19—7.22 (1H, m), 7.25—7.30 (2H, m). EI-MS *m/z*: 189 (M⁺).

A 4N HCl solution in 1,4-dioxane (0.476 ml, 1.90 mmol) was added to a mixture of compound **8** (300 mg, 1.59 mmol), AcOEt (3 ml) and MeOH (0.3 ml) at 0 °C, and the whole was stirred for 20 min at the same temperature. The resulting precipitate was collected by filtration, washed with AcOEt and recrystallized from MeOH-AcOEt to give the hydrochloride of compound **8** (230 mg, 64%) as colorless crystals. mp 191—193 °C. ¹H-NMR (DMSO-*d*₆) δ: 1.74—1.81 (2H, m), 2.16—2.27 (2H, m), 3.04—3.12 (2H, m), 3.27—3.36 (2H, m), 5.04 (2H, s), 7.15—7.20 (1H, m), 7.31—7.37 (3H, m). EI-MS *m/z*: 189 (M⁺).

(±)-N-[2-(3,4-Dichlorophenyl)-4-hydroxybutyl]-N-methylbenzamide (11) A mixture of Ac₂O (10.0 ml, 112 mmol) and HCOOH (10.0 ml, 265 mmol) was stirred for 30 min at 60 °C, and then a solution of (±)-4-amino-3-(3,4-dichlorophenyl)butan-1-ol (**9**, 2.99 g, 12.8 mmol) in THF (30 ml) was added at 0 °C. The mixture was stirred for 2.5 h at room temperature, poured into H₂O and neutralized with K₂CO₃, and the product was extracted with CHCl₃. The extract was washed with saturated brine, and the organic layer was dried over anhydrous MgSO₄ and concentrated *in vacuo*. The residue was diluted with THF (30 ml), then added to BH₃-THF complex (1.0 M solution in THF, 38 ml, 0.38 mol) at 0 °C, and the mixture was heated to reflux for 5 h. Next, EtOH (30 ml) and diethanolamine (2.69 g, 25.6 mmol) were added at 0 °C, and the mixture was heated again to reflux for 14 h and concentrated *in vacuo*. The residue was diluted with brine, and extracted with AcOEt. The extract was washed with saturated brine, dried over anhydrous MgSO₄ and concentrated *in vacuo*. CH₂Cl₂ (70 ml) and Et₃N (4.46 ml, 32.0 mmol) were added to the residue, and the mixture was cooled to

–30 °C. A solution of benzoyl chloride (3.27 ml, 28.2 mmol) in CH₂Cl₂ (10 ml) was added. The whole was stirred for 5.5 h at room temperature and diluted with CHCl₃. This mixture was washed with 0.5N HCl, saturated NaHCO₃ and saturated brine, dried over anhydrous MgSO₄ and concentrated *in vacuo*. EtOH (50 ml) and 1N NaOH (50 ml) were added to the residue, and the mixture was stirred for 14 h at room temperature and concentrated *in vacuo*. The residue was diluted with brine and extracted with CHCl₃. The extract was washed with saturated brine, dried over anhydrous MgSO₄ and concentrated *in vacuo*. The resultant solid was recrystallized from AcOEt–hexane to give the benzamide (**11**, 3.49 g, 77%) as a colorless powder. ¹H-NMR (DMSO-*d*₆) δ: 1.48–2.33 (3H, m), 2.65–3.13 (3H, m), 3.26–4.00 (5H, m), 6.76–7.24 (3H, m), 7.30–7.45 (5H, m). FAB-MS *m/z*: 352 [(M+H)⁺]. *Anal.* Calcd for C₁₈H₁₉Cl₂NO₂: C, 61.37; H, 5.44; N, 3.98. Found: C, 61.33; H, 5.47; N, 3.88.

(±)-*N*-[2-(3,4-Dichlorophenyl)-4-(spiro[isobenzofuran-1(3*H*),4'-piperidin]-1'-yl)butyl]-*N*-methylbenzamide Monofumarate (**12**) MsCl (0.064 ml, 0.83 mmol) was added to a mixture of compound **11** (233 mg, 0.660 mmol), Et₃N (0.138 ml, 0.991 mmol) and CH₂Cl₂ (10 ml) at 0 °C and the whole was stirred for 50 min at room temperature. It was diluted with AcOEt, washed with H₂O and saturated brine, dried over anhydrous MgSO₄ and concentrated *in vacuo*. The residue was dissolved in *N,N*-dimethylformamide (DMF, 5 ml), then spiro[isobenzofuran-1(3*H*),4'-piperidine] (**8**) (138 mg, 0.727 mmol) and Et₃N (0.276 ml, 1.98 mmol) were added, and the mixture was stirred for 4 h at 70 °C. It was then diluted with AcOEt, washed with saturated brine, dried over anhydrous MgSO₄ and concentrated *in vacuo*. The residue was purified by column chromatography (CHCl₃:MeOH=49:1) to give the free base of compound **12** (156 mg, 45%) as a pale yellow oil. Fumaric acid (221 mg, 1.91 mmol) and MeOH (8 ml) were added to the free base of compound **12** (1.00 g, 1.91 mmol), and the mixture was concentrated *in vacuo*. The residue was crystallized from 2-butanone–Et₂O and recrystallized from MeOH–MeCN to give the monofumarate (**12**, 767 mg, 63%) as a colorless powder. mp 177–178 °C. ¹H-NMR (DMSO-*d*₆) δ: 1.56–1.75 (3H, m), 1.86–2.04 (4H, m), 2.12–2.60 (3H+DMSO, m), 2.67–3.04 (5H, m), 3.16–3.80 (3H, m), 4.97 (2H, s), 6.60 (2H, s), 6.98–7.18 (2H, m), 7.20–7.44 (8H, m), 7.48–7.69 (2H, m). FAB-MS *m/z*: 523 [(M+H)⁺]. *Anal.* Calcd for C₃₀H₃₂Cl₂N₂O₂·C₄H₄O₄: C, 63.85; H, 5.67; N, 4.38. Found: C, 63.95; H, 5.72; N, 4.38.

Energy Minimization The 2-dimensional structures of 4-acetamido-1-methyl-4-phenylpiperidine (**13**) and 1'-methylspiro[isobenzofuran-1(3*H*),4'-piperidine] (**14**) were constructed using the molecular modeling program Sybyl (6.22) on an Indigo Elan Workstation (Silicon Graphics, Inc., Mountain View, CA) and converted to initial 3-dimensional structures with Concord 3.2.1. These initial structures were relaxed using the Tripos force field. The total energy of compound **13** was 4.93 kcal/mol with the equatorial phenyl group and 6.61 kcal/mol with the axial phenyl group. The total energy of compound **14** was 10.00 kcal/mol with the equatorial phenyl group and 12.51 kcal/mol with the axial phenyl group.

Binding Assays Binding studies were carried out according to the method described by Burcher and Buck.¹³⁾ To determine the NK₂ receptor binding affinities of the compounds, ¹²⁵I-NKA and hamster urinary bladder were used, and ¹²⁵I-Bolton–Hunter-SP and guinea pig urinary bladder were employed to test NK₁ binding affinities.

In Vivo Assay Bronchospasm was induced with [β-Ala⁸]-NKA(4–10) (1 nmol/kg i.v.) in urethane-anesthetized guinea pigs under mechanical ventilation.^{14,15)} Inhibitory activities of the compounds were determined by measuring the reduction in the agonist-induced maximal responses after administration. Test compounds were given *via* the i.v.

route 15 min before challenge with the agonist, and lung resistance was measured using a whole-body plethysmogram. The doses required to reduce the responses by 50% (ID₅₀) were determined by probit analysis.

Acknowledgements We are grateful to Drs. K. Murase and T. Mase for their valuable suggestions and encouragement and to the staff of the Molecular Chemistry Research Laboratory for spectral measurements, elemental analyses and energy calculations.

References

- 1) Von Euler U. S., Gaddum J. H., *J. Physiol.*, **72**, 74–86 (1931); Chang M. M., Leeman S. E., Niall H. D., *Nature New Biol.*, **232**, 86–87 (1971).
- 2) Kimura S., Okada M., Sugita Y., Kanazawa I., Munekata E., *Proc. Jpn. Acad. (B)*, **59**, 101–104 (1983).
- 3) Barnes M., Yoshioka K., *Physiol. Rev.*, **73**, 229–308 (1993); Maggi C. A., Patacchini R., Rovero P., Giachetti A., *J. Auton. Pharmacol.*, **13**, 23–93 (1993).
- 4) Nakanishi S., *Annu. Rev. Neurosci.*, **14**, 123–136 (1991); Gerard N. P., Bao L., Ping H. X., Gerard C., *Regul. Pept.*, **43**, 21–35 (1993).
- 5) Barnes P. J., *Lancet*, **1**, 242–245 (1986); *idem*, *Trends Pharmacol. Sci.*, **8**, 24–27 (1987); *idem*, *Arch. Int. Pharmacodyn.*, **303**, 67–82 (1990); Barnes P. J., Belvisi M. G., Rogers D. F., *Trends Pharmacol. Sci.*, **11**, 185–189 (1990).
- 6) Sekizawa K., Tamaoki J., Graf P. D., Basbaum C. B., Borson D. B., Nadel J. A., *J. Pharm. Exp. Ther.*, **243**, 1211–1217 (1987).
- 7) Uchida Y., Nomura A., Ohtsuka M., Hasegawa S., *Am. Rev. Respir. Dis.*, **136**, 718–721 (1987).
- 8) Desai M. C., Lefkowitz S. L., Thadeio P. F., Longo K. P., Snider R. M., *J. Med. Chem.*, **35**, 4911–4913 (1992); Emonds-Alt X., Doutremepuich J.-D., Heaulme M., Neliat G., Santucci V., Steinberg R., Vilain P., Bichon D., Ducoux J.-P., Proietto V., Van Broeck D., Soubrie P., Fur G. L., Breliere J.-C., *Eur. J. Pharmacol.*, **250**, 403–413 (1993); Vassout A., Schaub M., Gentsch C., Ofner S., Schilling W., Veenstra S., *Neuropeptides*, **26** (Suppl. 1), 38 (1994); Tabart M., Peyronel J.-F., *Bioorg. Med. Chem. Lett.*, **4**, 673–676 (1994); Hagiwara D., Miyake H., Igari N., Karino M., Maeda Y., Fujii T., Matsuo M., *J. Med. Chem.*, **37**, 2090–2099 (1994); Harrison T., Williams B. J., Swain C. J., Ball R. G., *Bioorg. Med. Chem. Lett.*, **4**, 2545–2550 (1994); Natsugari H., Ikeura Y., Kiyota Y., Ishiuchi Y., Ishimaru T., Saga O., Shirafuji H., Tanaka T., Kamo I., Doi T., Otsuka M., *J. Med. Chem.*, **38**, 3106–3120 (1995).
- 9) Emonds-Alt X., Vilain P., Goulaouic P., Proietto V., Van Broeck D., Advenier C., Naline E., Neliat G., Le Fur G., Breliere J. C., *Life Sciences*, **50**, PL-101–106 (1992).
- 10) Emonds-Alt X., Goulaouic P., Proietto V., Van Broeck D., *Eur. Pat. Appl.* 474561 (1992) [*Chem. Abstr.*, **117**, 26590b (1992)].
- 11) Pearson W. H., Schkeryantz J. M., *J. Org. Chem.*, **57**, 6783–6789 (1992).
- 12) Emonds-Alt X., Proietto V., Van Broeck D., Vilain P., Advenier C., Neliat G., Le Fur G., Breliere J. C., *Bioorg. Med. Chem. Lett.*, **3**, 925–930 (1993).
- 13) Burcher E., Buck S. H., *Eur. J. Pharmacol.*, **128**, 165–177 (1986).
- 14) Maggi C. A., Patacchini R., Giuliani S., Giachetti A., *Eur. J. Pharmacol.*, **234**, 83–90 (1993).
- 15) Konzett H., Rossler R., *Arch. Exp. Path. Pharm.*, **195**, 71–74 (1940).
- 16) Burkholder T. P., Kudlacz E. M., Le T.-B., Knippenberg R. W., Shatzer S. A., Maynard G. D., Webster M. E., Horgan S. W., *Bioorg. Med. Chem. Lett.*, **6**, 951–956 (1996).