

## Delaumonones A and B, New Antiplasmodial Quassinoids from *Laumoniera bruceadelpha*

Shiori OSHIMI,<sup>a</sup> Aki TAKASAKI,<sup>a</sup> Yusuke HIRASAWA,<sup>a</sup> Takahiro HOSOYA,<sup>a</sup> Khalijah AWANG,<sup>b</sup> A. Hamid A. HADI,<sup>b</sup> Wiwied EKASARI,<sup>c</sup> Aty WIDYAWARUYANTI,<sup>c</sup> and Hiroshi MORITA<sup>\*,a</sup>

<sup>a</sup>Faculty of Pharmaceutical Sciences, Hoshi University; 2–4–41 Ebara, Shinagawa-ku, Tokyo 142–8501, Japan:

<sup>b</sup>Department of Chemistry, Faculty of Science, University of Malaya; 50603 Kuala Lumpur, Malaysia: and <sup>c</sup>Faculty of Pharmacy, Airlangga University; Jalan Dharmawangsa Dalam, Surabaya 60286, Indonesia.

Received February 28, 2009; accepted May 23, 2009; published online May 29, 2009

**New quassinoids, delaumonones A (1) and B (2) have been isolated from the bark of *Laumoniera bruceadelpha* NOOTEBOOM (Simaroubaceae) and the structures were elucidated by 2D NMR analysis and a chemical correlation. Delaumonones showed an antimalarial activity against *Plasmodium falciparum*.**

**Key words** quassinoid; delaumonone; antimalarial activity; *Laumoniera bruceadelpha*

Malaria is one of the crucial infectious disease in the world and continues to cause morbidity and mortality on a large scale in tropical countries.<sup>1)</sup> The antimalarial potential of compounds derived from plants is proven by examples such as quinine from *Cinchona* species and artemisinin from *Artemisia annua*.<sup>2)</sup> The plants belonging to Simaroubaceae are known to contain various quassinoids with biological activities, such as antimalarial, antifeedant, anti-inflammatory, antiulcer, antipyretic, and cytotoxic activities.<sup>3–6)</sup> Recent studies of numerous Simaroubaceae plants have highlighted good antimalarial activity of certain quassinoids against chloroquine-resistant strains of *Plasmodium falciparum*.<sup>7)</sup> These results have prompted us to search for new quassinoids for possible antimalarial action.

With an aim to isolate additional antimalarial natural products,<sup>8–13)</sup> purification of the extracts from the bark of *Laumoniera bruceadelpha* NOOTEBOOM (Simaroubaceae) collected in Malaysia led to isolate two new quassinoids, delaumonones A (1) and B (2). This paper describes the isolation and structural elucidation of 1 and 2 with an antimalarial activity against *P. falciparum*.

The bark of *L. bruceadelpha* (1.4 kg), which was collected in Malaysia, was extracted with MeOH, and the extract was partitioned between CHCl<sub>3</sub>, *n*-BuOH, and H<sub>2</sub>O. CHCl<sub>3</sub>-soluble materials were subjected to a silica gel column (CHCl<sub>3</sub>/MeOH), and then an ODS column followed by ODS HPLC to afford delaumonones A (1, 11.9 mg) and B (2, 5.6 mg) together with isobrucein A (3, 307.0 mg)<sup>14)</sup> and isobrucein B (4, 5.8 mg).<sup>15)</sup> *n*-BuOH-soluble materials were subjected to an HP-20 column, and then a silica gel column (CHCl<sub>3</sub>/MeOH), followed by ODS HPLC to afford delaumonone A (1, 15.0 mg) together with brucein D (5, 1.2 mg)<sup>16)</sup> and yadanzolid A (6, 3.7 mg).<sup>17)</sup>

Delaumonone A (1), colorless solid,  $[\alpha]_D^{20} +20$  ( $c=1.0$ , MeOH), showed molecular formula, C<sub>25</sub>H<sub>32</sub>O<sub>11</sub>, which was determined by HR-ESI-MS [ $m/z$  531.1827 (M+Na)<sup>+</sup>,  $\Delta -1.5$  mmu]. IR absorptions implied the presence of hydroxyl (3439 cm<sup>-1</sup>) and carbonyl (1735, 1665 cm<sup>-1</sup>) functionalities. <sup>1</sup>H- and <sup>13</sup>C-NMR data are presented in Table 1. The <sup>13</sup>C-NMR spectrum revealed 25 carbon signals due to five *sp*<sup>2</sup> quaternary carbons, three *sp*<sup>3</sup> quaternary carbons, one *sp*<sup>2</sup> methine, nine *sp*<sup>3</sup> methines, three methylenes, and four methyls. Among eight quaternary carbons ( $\delta_C$  45.7, 49.0,

81.0, 164.0, 168.2, 172.1, 173.8, 198.4), five quaternary carbons ( $\delta_C$  81.0, 168.2, 172.1, 173.8, 198.4) were ascribed to those bearing an oxygen atom.

The molecular structure of 1 was deduced from extensive analyses of the two-dimensional NMR data, including the <sup>1</sup>H–<sup>1</sup>H COSY, HMQC, and HMBC spectra in CD<sub>3</sub>OD (Fig. 1). The <sup>1</sup>H–<sup>1</sup>H COSY and HOHAHA spectra revealed connectivities of five partial structures **a** (C-3–C-4 and C-18), **b** (C-5–C-7), **c** (C-9 and C-11–C-12), **d** (C-14–C-15), and **e** (C-2'–C-5') as shown in Fig. 1. Connectivity of C-4 to C-5 was implied by an HMBC correlation for H<sub>3</sub>-18 to C-5 ( $\delta_C$  42.4). HMBC correlations were observed for H<sub>3</sub>-19 to C-1 ( $\delta_C$  81.5), C-5, C-9 ( $\delta_C$  41.6), and C-10 ( $\delta_C$  49.0), H-1 to C-2 ( $\delta_C$  198.4), H-3 to C-1, suggesting that C-19 was connected through C-10 in a decalin ring system. HMBC cross peaks for H<sub>2</sub>-20 to C-8 ( $\delta_C$  45.7), C-12 ( $\delta_C$  75.0), and C-13 ( $\delta_C$  81.0) indicated that the presence of an ether bridge between C-13 and C-20. In addition, connection of unit **c** and a carboxylic acid through C-13 was also indicated by the HMBC correlations for H-12 to C-13 and C-21 ( $\delta_C$  172.1). The presence of an isopentanoate at C-15 was implied by the HMBC correlations for H-15 and H-3' to C-1' ( $\delta_C$  173.8). The presence of a  $\delta$ -lactone ring characteristic of C<sub>20</sub> quassinoid skeleton was indicated by an HMBC correlation for H-14 to C-16 ( $\delta_C$  168.2). Thus, the molecular structure of de-




Chart 1

\* To whom correspondence should be addressed. e-mail: moritah@hoshi.ac.jp

laumonone A was elucidated to be **1**.

The relative configuration of **1** was elucidated by NOESY correlations as depicted in the computer-generated three-dimensional drawing (Fig. 2). The NOESY correlations of H-1/H-5 and H-9, and H-9/H-5 and H-15 indicated that these protons at C-1, C-5, C-9, and C-15 were  $\alpha$ -orientated. The  $\beta$ -orientation of H<sub>3</sub>-19, H-7, and H-14, and  $\alpha$ -orientation of H-11 were indicated by NOESY correlations of H-20/H-7 and H-14, and  $^3J_{\text{H-H}}$  coupling constant (3.2 Hz) of H-9/H-11.

Table 1.  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR Data for Delaumonones A and B (**1**, **2**) in  $\text{CD}_3\text{OD}$  at 300 K<sup>a)</sup>

Position	$\delta_{\text{H}}$ (int.; mult.; $J$ (Hz))		$\delta_{\text{C}}$	
	<b>1</b>	<b>2</b>	<b>1</b>	<b>2</b>
1	4.29 (1H, s)	3.89 (1H, s)	81.5	76.3
2			198.4	199.8
3	6.04 (1H, br s)	5.87 (1H, br s)	123.9	125.1
4			164.0	164.6
5	3.00 (1H, br d, 12.4)	3.06 (1H, br d, 12.7)	42.4	39.3
6a	1.96 (1H, m)	1.90 (1H, m)	27.8	28.9
6b	2.34 (1H, ddd, 14.8, 2.8, 2.8)	2.33 (1H, br d, 15.4)		
7	4.90 (1H, m)	4.90 (1H, m)	83.3	83.2
8			45.7	46.9
9	2.45 (1H, br d, 3.2)	2.70 (1H, br s)	41.6	35.3
10			49.0	44.7
11	4.72 (1H, br d, 5.2)	4.26 (1H, br d, 4.7)	73.8	72.5
12	4.20 (1H, br s)	4.20 (1H, br s)	75.0	76.4
13			81.0	84.0
14	3.26 (1H, m)	3.35 (1H, m)	51.2	51.2
15	6.25 (1H, m)	6.22 (1H, m)	68.7	66.0
16			168.2	169.6
18	1.97 (3H, s)	1.95 (3H, s)	21.2	22.5
19	1.19 (3H, s)	1.26 (3H, s)	10.1	14.8
20a	3.71 (1H, d, 7.6)	3.73 (1H, d, 7.6)	72.9	74.7
20b	4.68 (1H, d, 7.6)	4.71 (1H, d, 7.6)		
21			172.1	172.4
1'			173.8	173.1
2'	2.21 (2H, d, 6.8)	2.20 (2H, d, 6.8)	41.9	44.8
3'	2.07 (1H, m)	2.06 (1H, m)	25.2	26.7
4'	0.96 (3H, d, 6.8)	0.97 (3H, d, 6.6)	21.4	22.7
5'	0.97 (3H, d, 6.8)	0.98 (3H, d, 6.6)	21.4	22.7
OMe		3.79 (3H, s)		53.1

a)  $\delta$  in ppm.




Fig. 1. Selected 2D NMR Correlations for Delaumonone A (**1**)




Fig. 3. Selected 2D NMR Correlations and Relative Configuration of Delaumonone B (**2**)

Treatment of **1** with TMS-diazomethane afforded a methyl ester derivative, whose spectroscopic data and specific rotation were identical with those of isobrucein A (**3**).<sup>14</sup> Thus, the absolute configuration of delaumonone A was assigned as **1**.

Delaumonone B (**2**), colorless solid,  $[\alpha]_{\text{D}}^{20} +18$  ( $c=1.0$ , MeOH), showed molecular formula,  $\text{C}_{26}\text{H}_{34}\text{O}_{11}$ , which was determined by HR-ESI-MS [ $m/z$  523.2180,  $(\text{M}+\text{H})^+$ ,  $\Delta -0.1$  mmu]. IR absorptions implied the presence of hydroxyl ( $3477\text{ cm}^{-1}$ ) and carbonyl ( $1743$ ,  $1668\text{ cm}^{-1}$ ) functionalities.  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR data are presented in Table 1. The  $^{13}\text{C}$ -NMR spectrum revealed 26 carbon signals due to five  $sp^2$  quaternary carbons, three  $sp^3$  quaternary carbons, one  $sp^2$  methine, nine  $sp^3$  methines, three methylenes, and five methyls. Among eight quaternary carbons ( $\delta_{\text{C}}$  44.7, 46.9, 84.0, 164.6, 169.6, 172.4, 173.1, 199.8), five quaternary carbons ( $\delta_{\text{C}}$  84.0, 169.6, 172.4, 173.1, 199.8) were ascribed to those bearing an oxygen atom.

$^1\text{H}$ - $^1\text{H}$  COSY, HMQC, and HMBC spectra in Fig. 3 suggested that **2** had the same pentacyclic backbone framework as that of isobrucein A (**3**),<sup>14</sup> although the  $^1\text{H}$ -NMR chemical shifts of H-1 ( $\delta_{\text{H}}$  3.89), H-9 ( $\delta_{\text{H}}$  2.70), and H-11 ( $\delta_{\text{H}}$  4.26) in **2** were remarkably not identical to those [H-1 ( $\delta_{\text{H}}$  4.17), H-9 ( $\delta_{\text{H}}$  2.33), and H-11 ( $\delta_{\text{H}}$  4.73)] in **3**, indicating that it was a stereoisomer of isobrucein A (**3**). The relative configuration of **2** was elucidated by NOESY correlations (Fig. 3). H-1 was assigned to be equatorial. NOESY cross peaks for H-1 to H<sub>3</sub>-19 and H-11 were observed in the case of **2**, but not for **1** and **3**. This suggested that delaumonone B (**2**) was 1-*epi* form of isobrucein A (**3**). Thus, the relative configuration of delaumonone B (**2**) was assigned as shown.

Delaumonones A (**1**) and B (**2**) showed a potent *in vitro* antiplasmodial activity against *P. falciparum* 3D7 ( $\text{IC}_{50}$  **1**: 0.31  $\mu\text{g}/\text{ml}$ ; **2**: 0.60  $\mu\text{g}/\text{ml}$ ) and cytotoxicity against HL-60 cells ( $\text{IC}_{50}$  **1**: 1.6  $\mu\text{g}/\text{ml}$ ; **2**: 2.4  $\mu\text{g}/\text{ml}$ ). Among isolated quassinoids, isobrucein A<sup>14</sup>) ( $\text{IC}_{50}$  **3**: 0.024  $\mu\text{g}/\text{ml}$ ) were more active than delaumonones A and B, but also showed a potent cytotoxicity against HL-60 cells ( $\text{IC}_{50}$  **3**: 0.0068  $\mu\text{g}/\text{ml}$ ).





Fig. 2. Selected NOESY Correlations and Relative Configuration for Delaumonone A (**1**)



Delaumonones A and B had good selectivity index more than isobrucein A, suggesting that the presence of  $\beta$ -orientation of the hydroxyl group at C-1 and the methoxy carbonyl group at C-21 contributed to a cytotoxicity more than an antiplasmodial activity.

In this study, we have isolated two new quassinoids, delau-monones A and B (**1**, **2**) from the bark of *Laumoniera bruceadelpha*. Delaumonones showed an antimalarial activity against *P. falciparum*. Cytotoxic activity against cancer cells was influenced by the nature of the C-21 side chain and of configuration at C-1, which was not much affected by an antimalarial activity against *P. falciparum*.

## Experimental

**General Experimental Procedures** Optical rotations were measured on a JASCO P-1030 polarimeter. UV spectra were recorded on a Shimadzu UV-250 spectrophotometer and IR spectra on a JASCO FTIR-230 spectrometer. Mass spectra were obtained with a Micromass LCT spectrometer.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded on a 600 MHz spectrometer at 300 K, while  $^{13}\text{C}$ -NMR spectra were measured on a 150 MHz spectrometer. Each NMR sample of delau-monones was prepared by dissolving in  $30\ \mu\text{l}$  of  $\text{CD}_3\text{OD}$  in 2.5 mm micro cells (Shigemi Co., Ltd.) and chemical shifts were reported using residual  $\text{CD}_3\text{OD}$  ( $\delta_{\text{H}}$  3.31 and  $\delta_{\text{C}}$  49.0) as internal standard. Standard pulse sequences were employed for the 2D NMR experiments. COSY, HO-HAHA, and NOESY spectra were measured with spectral widths of both dimensions of 4800 Hz, and 32 scans with two dummy scans were accumulated into 1 K data points for each of 256  $t_1$  increments. NOESY and HO-HAHA spectra in the phase sensitive mode were measured with a mixing time of 800 and 30 ms, respectively. For HMQC spectra in the phase sensitive mode and HMBC spectra, a total of 256 increments of 1 K data points were collected. For HMBC spectra with Z-axis PFG, a 50 ms delay time was used for long-range C-H coupling. Zero-filling to 1 K for  $F_1$  and multiplication with squared cosine-bell windows shifted in both dimensions were performed prior to 2D Fourier transformation.

**Plant Material** The bark of *Laumoniera bruceadelpha* was collected at Mersing, Malaysia in 2001. The botanical identification was made by Mr. Teo Leong Eng, Faculty of Science, University of Malaya. Voucher specimens (KL4099) are deposited in the Herbarium of Chemistry Department, University of Malaya.

**Extraction and Isolation** The bark of *L. bruceadelpha* (1.4 kg), which was collected at Malaysia, was extracted with MeOH, and a part (40 g) of the extract (126 g) was partitioned between  $\text{CHCl}_3$ , *n*-BuOH and  $\text{H}_2\text{O}$ .  $\text{CHCl}_3$ -soluble materials were subjected to a silica gel column ( $\text{CHCl}_3/\text{MeOH}$ , 1:0 $\rightarrow$ 0:1), in which a fraction eluted by  $\text{CHCl}_3/\text{MeOH}$  (40:1) was further purified on an ODS column with  $\text{MeOH}/\text{H}_2\text{O}$  (0:1 $\rightarrow$ 1:0) followed by ODS HPLC (55% MeOH) to afford delau-monone B (**2**, 5.6 mg) together with isobrucein A (**3**, 307.0 mg)<sup>14</sup> and isobrucein B (**4**, 5.8 mg).<sup>15</sup> A fraction eluted by  $\text{CHCl}_3/\text{MeOH}$  (0:1) was further purified on ODS HPLC with 55% MeOH containing 0.1% TFA to afford delau-monone A (**1**, 11.9 mg). *n*-BuOH-soluble materials were subjected to an HP-20 column ( $\text{H}_2\text{O}/\text{MeOH}$ , 1:0 $\rightarrow$ 0:1), in which a fraction eluted 80% MeOH was further purified on a silica gel column ( $\text{CHCl}_3/\text{MeOH}$ , 1:0 $\rightarrow$ 0:1) and ODS HPLC (55% MeOH containing 0.1% TFA) to afford delau-monone A (**1**, 15.0 mg) together with brucein D (**5**, 1.2 mg)<sup>16</sup> and yadanzolide A (**6**, 3.7 mg).<sup>17</sup> These compounds could also be isolated from EtOH extract of the same plant.

Delaumonone A (**1**): Colorless amorphous solid,  $[\alpha]_{\text{D}}^{20}$  +20 ( $c=1.0$ , MeOH); IR (KBr)  $\nu_{\text{max}}$  3439, 1735, and 1665  $\text{cm}^{-1}$ ;  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR (Table 1); ESI-MS (neg.)  $m/z$ : 507 ( $\text{M}-\text{H}^-$ ); HR-ESI-TOF-MS (pos.)  $m/z$ : 531.1827 ( $\text{M}+\text{Na}^+$ ), Calcd for  $\text{C}_{25}\text{H}_{32}\text{O}_{11}\text{Na}$ .

Delaumonone B (**2**): Colorless amorphous solid,  $[\alpha]_{\text{D}}^{20}$  +18 ( $c=1.0$ , MeOH); IR (KBr)  $\nu_{\text{max}}$  3477, 1743, and 1668  $\text{cm}^{-1}$ ;  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR (Table 1); ESI-MS (pos.)  $m/z$ : 545 ( $\text{M}+\text{Na}^+$ ); HR-ESI-TOF-MS (pos.)  $m/z$ : 523.2180 ( $\text{M}+\text{H}^+$ ), Calcd for  $\text{C}_{26}\text{H}_{35}\text{O}_{11}$ .

**Conversion of Delaumonone A (**1**) to Isobrucein A (**3**)** To a solution of delau-monone A (2.0 mg) in MeOH (0.2 ml) was added TMS-diazomethane (20  $\mu\text{l}$ ), and the mixture was kept at r.t. for 3 h. After evaporation, a compound (1.9 mg), whose spectroscopic data and  $[\alpha]_{\text{D}}$  value were identical with those of natural isobrucein A (**3**), was obtained.

**Antiplasmodial Activity** Human malaria parasites were cultured according to the method of Trager and Jensen.<sup>18</sup> The antimalarial activity of

the isolated compounds was determined by the procedure described by Budimulja *et al.*<sup>19</sup> In brief, stock solutions of the samples were prepared in DMSO (final DMSO concentrations of <0.5%) and were diluted to the required concentration with complete medium (RPMI 1640 supplemented with 10% human plasma, 25 mM HEPES and 25 mM  $\text{NaHCO}_3$ ) until the final concentrations of samples in culture plate wells were 10; 1; 0.1; 0.01; 0.001  $\mu\text{g}/\text{ml}$ . The malarial parasite *P. falciparum* 3D7 clone was propagated in a 24-well culture plates. Growth of the parasite was monitored by making a blood smear fixed with MeOH and stained with Geimsa stain. The antimalarial activity of each compound was expressed as an  $\text{IC}_{50}$  value, defined as the concentration of the compound causing 50% inhibition of parasite growth relative to an untreated control.

The percentage of growth inhibition was expressed according to following equation: growth inhibition % =  $100 - [(\text{test parasitaemia}/\text{control parasitemia}) \times 100]$ . Chloroquine:  $\text{IC}_{50}$  0.0061  $\mu\text{g}/\text{ml}$ .

**Cytotoxic Activity** HL-60 (human blood premyelocytic leukemia) cell line was seeded onto 96-well microtiter plates at  $1 \times 10^4$  cells per well. Cells were preincubated for 24 h at 37  $^{\circ}\text{C}$  in humidified atmosphere of 5%  $\text{CO}_2$ . Different concentrations of each compound (10  $\mu\text{l}$ ) were added to the cultures, and then the cells were incubated at 37  $^{\circ}\text{C}$  for 48 h. On the third day, 15  $\mu\text{l}$  MTT solution (5 mg/ml) was added into each well of the cultured medium. After further 2 h of incubation, 100  $\mu\text{l}$  of 10% SDS-0.01 N HCl solution was added to each well and the formazan crystals in each well were dissolved by stirring with a pipette. The optical density measurements were made using a micropipette reader (Benchmark Plus microplate spectrometer, BIO-RAD) equipped with a two wavelengths system (550, 700 nm). In each experiment, three replicate of wells were prepared for each sample. The ratio of the living cells was determined based on the difference of the absorbance between those of samples and controls. These differences are expressed in percentage and cytotoxic activity was indicated as an  $\text{IC}_{50}$  value.

**Acknowledgments** This work was partly supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and a Grant from the Open Research Center Project. We also acknowledge the financial support provided by Faculty of Pharmacy, Airlangga University for antimalarial activity.

## References

- 1) Wyler D. J., *Clin. Infect. Dis.*, **16**, 449—456 (1993).
- 2) Peters W., *Br. Med. Bull.*, **32**, 187—192 (1982).
- 3) Bawm S., Matsuura H., Elkhateeb A., Nabeta K., Nonaka N., Oku Y., Katakura K., *Vet. Parasitol.*, **158**, 288—294 (2008).
- 4) Muhammad I., Samoylenko V., *Expert Opinion*, **2**, 1065—1084 (2007).
- 5) Guo Z., Vangapandu S., Sindelar R. W., Walker L. A., Sindelar R. D., *Curr. Med. Chem.*, **12**, 173—190 (2005).
- 6) Klocke J. A., Arisawa M., Honda S., Kinghorn A. D., Cordell G. A., Farnsworth N. R., *Experientia*, **41**, 379—382 (1985).
- 7) Ang H. H., Chan K. L., Mak J. W., *Planta Med.*, **61**, 177—178 (1995).
- 8) Koyama K., Hirasawa Y., Zaima K., Hoe T. C., Chan K. L., Morita H., *Bioorg. Med. Chem.*, **16**, 6483—6488 (2008).
- 9) Hirasawa Y., Miyama S., Kawahara N., Goda Y., Rahman A., Ekasari W., Widyawaruyanti A., Indrayanto G., Zaini N. C., Morita H., *Heterocycles*, **79**, 1107—1112 (2009).
- 10) Hirasawa Y., Arai H., Zaima K., Oktarina R., Rahman A., Ekasari W., Widyawaruyanti A., Indrayanto G., Zaini N. C., Morita H., *J. Nat. Prod.*, **72**, 304—307 (2009).
- 11) Sekiguchi M., Hirasawa Y., Zaima K., Hoe T. C., Chan K.-L., Morita H., *Heterocycles*, **75**, 2283—2288 (2008).
- 12) Sekiguchi M., Hirasawa Y., Zaima K., Hoe T. C., Chan K.-L., Morita H., *Heterocycles*, **76**, 867—874 (2008).
- 13) Zaima K., Matsuno Y., Hirasawa Y., Rahman A., Indrayanto G., Zaini N. C., Morita H., *Heterocycles*, **75**, 2535—2540 (2008).
- 14) Polonsky J., Baskevitch-Varon Z., Sevenet T., *Experientia*, **31**, 1113—1114 (1975).
- 15) Okano M., Fukamiya N., Aratani T., *J. Nat. Prod.*, **48**, 972—975 (1985).
- 16) Li X., Wu L., Konda Y., Iguchi M., Takahashi H., Harigaya Y., Onda M., *J. Heterocycl. Chem.*, **26**, 493—501 (1989).
- 17) Lee K. H., Imakura Y., Sumida Y., Wu R. Y., Iris H., Huang H. C., *J. Org. Chem.*, **44**, 2180—2185 (1979).
- 18) Trager W., Jensen J. B., *Science*, **193**, 673—675 (1976).
- 19) Budimulja A. S., Syafruddin T. P., Wilairat P., Marzuki S., *Mol. Biochem. Parasitol.*, **84**, 137—141 (1997).