

## Novel Aromatization of Inositols in Dimethyl Sulphoxide–Acetic Anhydride

By ALEXANDER J. FATIADI

(Division of Analytical Chemistry, Institute for Materials Research, National Bureau of Standards, Washington, D.C. 20234)

DIMETHYL SULPHOXIDE in acetic anhydride has proved to be an efficient reagent for the oxidation of an isolated or sterically hindered secondary alcoholic group to a ketone group in steroids<sup>1</sup> and partly substituted carbohydrates.<sup>2</sup>

A unique, one-step aromatization of inositols (cyclohexanehexaols) following treatment with dimethyl sulphoxide–acetic anhydride is now reported.

*myo*-Inositol (2 g.) was stirred with a mixture of dry dimethyl sulphoxide (30 ml.), acetic anhydride (10 ml.), and dry pyridine (5 ml.) at 60–65° for 45 min. Treatment of the red-brown reaction mixture with ice-water deposited 1.9–2.1 g. (54–60%) of penta-acetoxybenzene, m.p. 153–158°. Recrystallization from 4:1 (v/v) ethanol–acetic acid raised the melting point to 166–168°. The product was identical (infrared spectrum, mixture m.p., and chromatography) with an authentic sample.<sup>3</sup>

Penta-acetoxybenzene has also been obtained

in good yield from *epi*-inositol, *dextro*-inositol, *laevo*-inositol, *muco*-inositol, and *scyllo*-inositol. Apparently, dimethyl sulphoxide–acetic anhydride is not a stereospecific oxidizing agent, in contrast to catalytic<sup>4</sup> or bacterial<sup>4</sup> oxidants, as *scyllo*-inositol, which has no axial hydroxyl groups and is known to be inert either to catalytic or bacterial oxidation,<sup>4</sup> is converted into penta-acetoxybenzene as readily as are the other inositols.

Surprisingly, penta-acetoxybenzene was not formed from either *myo*-inos-2-ose or *DL*-*epi*-inos-2-ose, the known oxidation products obtained from *myo*-inositol.<sup>5</sup> However, penta-acetoxybenzene was produced when the starting material was “*DL*-xylo-4,5,6-trihydroxycyclohexenediolic acid”, the enolic form of a diketoinositol that is obtained by oxidation of *myo*-inositol with nitric acid.<sup>3b</sup> Thus, the conversion of inositols into penta-acetoxybenzene probably proceeds through a form of the diketoinositol as an intermediate.

(Received, March 17th, 1967; Com. 264.)

<sup>1</sup> J. D. Albright and L. Goldman, *J. Amer. Chem. Soc.*, 1965, **87**, 4214.

<sup>2</sup> B. Lindberg and K. N. Slessor, *Carbohydrate Res.*, 1966, **1**, 492; D. Horton and J. S. Jewell, *ibid.*, **2**, 251.

<sup>3</sup> (a) A. Einhorn, J. Coblner, and H. Pfeiffer, *Ber.*, 1904, **37**, 100; (b) A. J. Fatiadi and H. S. Isbell, *J. Res. Nat. Bur. Stand., Sect. A*, 1964, **68**, 287.

<sup>4</sup> S. J. Angyal and L. Anderson, *Adv. Carbohydrate Chem.*, 1959, **14**, 135.

<sup>5</sup> T. Posternak, *Biochem. Prep.*, 1952, **2**, 57.