


The Total Synthesis of Racemic Isonootkatone (α -Vetivone)

By JAMES A. MARSHALL,* HERMANN FAUBL, and THOMAS M. WARNE, JUN.


(Department of Chemistry, Northwestern University, Evanston, Illinois 60201)

THE recent work of Endo and de Mayo,¹ and independent findings in our own laboratory² have shown that α -vetivone (XIV), long regarded as a relative of vetivazulene (I), must now be formulated as an isomer of nootkatone (II).³ For this reason we have suggested that this compound be renamed isonootkatone.² We now present a total synthesis of racemic isonootkatone by a route (Chart) which fully confirms the suggested structure.



Diethyl isopropylidene malonate,⁴ upon reduction with lithium aluminium hydride afforded the corresponding diol (III) [b.p. 78° at 0.2 mm.; δ (CCl₄) 4.18 (OH), 4.14 (CH₂O), and 1.74 p.p.m. (vinyl CH₃)†]. The dibromide, a powerful lachrymator obtained by treating diol (III) with PBr₃ in ether-hexane-pyridine, yielded the tetra-ester (V) *via* alkylation with an excess of diethyl sodiomalonate in 1,2-dimethoxyethane. Subsequent hydrolysis and decarboxylation converted (V) into diacid (VI) and this compound, in turn, gave the corresponding diester (VII) [b.p. 87–95° at 0.2–0.3 mm.; $\lambda_{\text{max}}^{\text{(nm)}}$ 5.76 (ester CO), 8.35, 8.53, 9.63, and 10.10 μ]. Dieckmann cyclization of the diester with sodium hydride in 1,2-dimethoxyethane afforded methyl 5-isopropylidene-2-oxocyclohexanecarboxylate (VIII) [b.p. 67–75° at 0.03–0.04 mm.; δ (CCl₄) 12.13 (enol OH), 3.76 (OCH₃), and 1.68 p.p.m. (two vinyl CH₃'s)]. Robinson annelation of this cyclic β -keto-ester occurred smoothly upon treatment first with *trans*-pent-3-en-2-one⁵ in the presence of 0.1 molar equivalent of methanolic sodium methoxide at 0° to effect the Michael addition, and then with an excess of the same base at room temperature for eighteen hours to complete the aldol cyclization. The resulting product [b.p. 65–100° (bath temperature) at 0.05–0.1 mm.; $\lambda_{\text{max}}^{\text{(nm)}}$ 5.79 (ester CO), 5.99 (ketone CO), 6.13 (C=C), 8.18, 8.48, 9.10, 10.05, 11.45, and 11.77 μ ; δ (CCl₄) 5.90 (vinyl H), 3.67 (OCH₃), 1.70, 1.76 (two vinyl CH₃'s), and 1.02


p.p.m. (CH₃ doublet, *J* 6 c./sec.)] solidified upon refrigeration and appears to be mainly the bicyclic keto-ester (IX). In our initial synthetic planning we felt that steric and electronic factors in the aforementioned Michael addition would favour the transition state orientation depicted in (XV). This arrangement leads ultimately to the bicyclic keto-ester (IX) with a *cis*-relationship between the methoxycarbonyl and methyl groups. The findings reported here tend to confirm our analysis. Additional work relating to this point is currently in progress.



Unsaturated ketone (IX) afforded the ethylene ketal derivative (X) [$\lambda_{\text{max}}^{\text{(nm)}}$ 5.79 (ester CO), 8.18, 8.83, 9.20, 10.03, 10.49, 11.05, and 12.64 μ ;


† With Me₄Si as internal reference throughout.

$\delta(\text{CCl}_4)$ 5.42 (vinyl H, $w_{h/2}$ 8 c./sec.), 3.88 ($\text{OCH}_2\cdot\text{CH}_2\text{O}$), 3.62 (OCH_3), 1.62, 1.72 (two vinyl CH_3 's), and 0.98 p.p.m. (CH_3 doublet, J 6 c./sec.) upon treatment with ethylene glycol and a trace of toluene-*p*-sulphonic acid in refluxing benzene (Dean-Stark water trap). Reduction of the ester (X) with ethereal lithium aluminium hydride gave the alcohol (XI) [$\lambda_{\text{max}}^{\text{OH}}$ 2.92 (OH), 8.90, 9.16, 9.50,



10.10, 10.50, and 12.58 μ ; $\delta(\text{C}_6\text{H}_5\text{N})$ 5.55 (vinyl H, $w_{h/2}$ 7 c./sec.), 5.03 (OH), 3.88 (CH_2O and $\text{OCH}_2\cdot\text{CH}_2\text{O}$), 1.75, 1.60 (vinyl CH_3 's), and 1.18 p.p.m. (CH_3 doublet, J 6 c./sec.)]. The methanesulphonate derivative (XII) yielded a mixture of the S-O and C-O cleavage products (XI) and (XIII) upon reduction with lithium in ammonia-ethanol. Racemic isonootakatonone (XIV) could be readily isolated by chromatography on silica gel after hydrolysis of the crude ketal mixture. The infrared and n.m.r. spectra and the gas-chromatographic retention time of the material thereby secured exactly matched those of natural isonootakatonone isolated from vetivert acetate.²

Treatment of the methanesulphonate derivative (XII) with ethereal lithium aluminium hydride led to none of the desired hydrogenolysis product (XIII). A cyclopropane derivative [$\delta(\text{CCl}_4)$ 3.50 ($\text{OCH}_2\cdot\text{CH}_2\text{O}$), 1.58 (vinyl CH_3), 0.98 (CH_3 doublet, J 6 c./sec.), and 0.25 p.p.m. (cyclopropane CH_2 , centre of AB quartet)], presumably (XVI), was the predominant product of this reduction.



(Received, June 12th, 1967; Com. 592.)

¹ K. Endo and P. de Mayo, *Chem. Comm.*, 1967, 89.

² J. A. Marshall and N. H. Andersen, *Tetrahedron Letters*, 1967, 1611.

³ W. D. Macleod, jun., *Tetrahedron Letters*, 1965, 4779.

⁴ A. C. Cope and E. M. Hancock, *J. Amer. Chem. Soc.*, 1938, **60**, 2644.

⁵ H. O. House, W. L. Respass, and G. M. Whitesides, *J. Org. Chem.*, 1966, **31**, 2138.