

## Selective Nucleophilic Attack by Peroxy-acid Anions on Nucleic Acid Components

By L. R. SUBBARAMAN, JIJI SUBBARAMAN, and E. J. BEHRMAN\*

(Department of Biochemistry, The Ohio State University, Columbus, Ohio 43210)

ALTHOUGH electrophilic attack by peroxy-acids on cytosine, adenine, and their nucleosides and nucleotides leading to the *N*-oxides has been recognized for some time,<sup>1</sup> it has not been realized that the anions of the same peroxy-acids are capable of nucleophilic attack on uracil, thymine, guanine, and their derivatives. In a study of the rate dependence on pH for the reaction of all of the common nucleic acid components with *m*-chloro-peroxybenzoic acid, reactions were run at 40° with equimolar concentrations of the peroxy-acid and substrate. The kinetics were second-order in all cases. The rate constants have been corrected for the rate of autodecomposition of the peroxy-acid. The rate of reaction for each nucleic acid component showed a characteristic pH maximum. The Table gives the observed second-order rate constant at this maximum and the pH at which it is observed. The reactions fall into two groups: cytosine, adenine, and their derivatives show

*The reaction of m-chloroperoxybenzoic acid with nucleic acid components*

Component	$k_{\text{obs}}$ (mole <sup>-1</sup> min. <sup>-1</sup> )	pH
Adenine .. .. .	4.7	6.2
Adenosine .. .. .	4.0	5.5
Cytosine .. .. .	13.2	6.6
Cytidine .. .. .	4.2	6.0
Deoxycytidine .. .. .	4.5	6.0
Guanosine .. .. .	4.4	8.4
Uracil .. .. .	8.3	8.8
Uridine .. .. .	3.8	8.6
Deoxyuridine .. .. .	3.2	8.5
5-Bromouracil .. .. .	8.3	8.4
6-Methyluracil .. .. .	4.3	8.6
Thymine .. .. .	1.9	8.8
Thymidine .. .. .	0.7	8.6

maxima on the acidic side of the  $pK_a$  of the peroxy-acid (7.6), while uracil, thymine, guanosine, and their derivatives, show maxima on the alkaline side. The kinetic data for the first group are consistent either with attack by the un-ionized

peroxy-acid on the uncharged substrates or with attack by the peroxy-acid anion on the cationic form of the substrate. We favour the first possibility on the basis of substituent effects observed by Dondoni, Modena, and Todesco<sup>2</sup> for the reaction between peroxybenzoic acid and a series of substituted pyridines and the fact that the only products detected in our study were the expected *N*-oxides. However, the two mechanisms described<sup>3</sup> for the oxidation of sulphoxides by peroxy-acids should be borne in mind. The kinetic results for the second group may be interpreted either as a reaction between the peroxy-acid anion and the uncharged substrate, or as a reaction between the un-ionized peroxy-acid and the substrate anion. We favour the first interpretation because of the substituent effects shown by

the uracil derivatives and because of the nature of the products. *N*-Oxides are not formed in the reactions of the second class. Instead, products that can only be attributed to ring-opening are found: from uracil, oxalic, and oxaluric acids and urea; from uridine, ribosylurea; from thymine, urea and hydroxyacetone; from thymidine, deoxyribosylurea; and from guanosine, ribosylurea and formic acid. In the case of the pyrimidines, we visualize attack by the peroxy-acid anion at the 5,6-double bond,<sup>4</sup> epoxide formation, hydrolysis and subsequent reactions similar to those formulated by Jones and his co-workers<sup>5</sup> for permanganate oxidation.

We thank the National Science Foundation for support.

(Received, June 21st, 1968; Com. 815.)

<sup>1</sup> F. Cramer, K. Randerath, and E. A. Schäfer, *Biochim. Biophys. Acta*, 1963, **72**, 150; F. Cramer and H. Seidel, *ibid.*, p. 157; T. J. Delia, M. J. Olsen, and G. B. Brown, *J. Org. Chem.*, 1965, **30**, 2766; H. Seidel, *Biochem. Biophys. Acta*, **138**, 98.

<sup>2</sup> A. Dondoni, G. Modena, and P. E. Todesco, *Gazzetta*, 1961, **91**, 613.

<sup>3</sup> R. Curci, A. Giovine, and G. Modena, *Tetrahedron*, 1966, **22**, 1235; R. Curci and G. Modena, *Tetrahedron Letters*, 1963, 1749.

<sup>4</sup> E. J. Behrman and J. O. Edwards, *Progr. Phys. Org. Chem.*, 1967, **4**, 93; J. B. Lee and B. C. Uff, *Quart. Rev.*, 1967, **21**, 429.

<sup>5</sup> M. H. Benn, B. Chatamra, and A. S. Jones, *J. Chem. Soc.*, 1960, 1014; B. Chatamra and A. S. Jones, *ibid.*, 1963, 811; P. Howgate, A. S. Jones, and J. R. Tittensor, *J. Chem. Soc. (C)*, 1968, 275.