

Reaction of Benzyne with Butadiene

By L. F. HATCH* and D. PETER

(Department of Chemistry, The University of Texas at Austin, Austin, Texas 78712)

BENZYNE has been shown to act as a dienophile with cyclic dienes, but apparently not with acyclic dienes. The reason has been given that benzyne does not survive long enough to permit the open-chain diene to assume the requisite *cis*-like arrangement that the cyclic dienes possess.¹ We have now been able to effect this type of reaction.

Benzyne and buta-1,3-diene, generated by the concurrent thermal decomposition of benzenediazonium-2-carboxylate² and 3-sulpholene (2,5-dihydrothiophen 1,1-dioxide) respectively, react in pentan-2-one at 100° to give a 9% yield of


1,4-dihydronaphthalene. A temperature of 100° is necessary to provide a reasonable, continuous concentration of butadiene from the 3-sulpholene; the method of Friedman and Logullo³ was therefore useful for the preparation and decomposition *in situ* of benzenediazonium-2-carboxylate.

It is suggested that the nascent butadiene from the decomposition of 3-sulpholene would be in the preferred *cis*-conformation, thus enabling reaction to take place.

That 3-sulpholene acts simply as a source of butadiene, and that the SO₂ entity is not involved in a transition state is shown by kinetic data⁴ obtained from the thermal decomposition of 3-sulpholene in the presence and absence of maleic anhydride as the dienophile. The reactions are first order with $k = 2.4 \times 10^{-3}$ (min.⁻¹). At temperatures below the decomposition temperature of 3-sulpholene, no addition product is formed with benzyne.

This research was supported by funds from The Robert A. Welch Foundation.

(Received, September 5th, 1968; Com. 1195.)



¹ J. F. Bunnett, *J. Chem. Educ.*, 1961, **38**, 278.

² M. Stiles and R. G. Miller, *J. Amer. Chem. Soc.*, 1960, **82**, 3802.

³ L. Friedman and F. M. Logullo, *J. Amer. Chem. Soc.*, 1963, **84**, 1549.

⁴ L. F. Hatch and D. Peter, unpublished results.