## Elaeocarpidine, a New Indole Alkaloid from *Elaeocarpus archboldianus* A.C.Sm.

By S. R. Johns, J. A. Lamberton, and A. A. Sioumis

(Division of Applied Chemistry, C.S.I.R.O., Chemical Research Laboratories, Box 4331 G.P.O., Melbourne, Australia)

ELAEOCARPIDINE, m.p. 229—230°  $[\alpha]_D \pm 0^\circ$  in CHCl<sub>3</sub>, the major alkaloid isolated from the leaves of *Elaeocarpus archboldianus* A.C.Sm. and the first indole alkaloid of the family Elaeocarpaceae, has been shown to be (I). Elemental analysis and an  $M^+$  peak at m/e 267 indicated the molecular formula  $C_{17}H_{21}N_3$ , and (I) was characterized spectroscopically as an indole  $[\lambda_{max}$  (EtOH) 226 mμ ( $\epsilon$  39,800), 283 (8,000),  $\lambda$  inflect., 290 (6,750);  $\nu_{max}$  (CHCl<sub>3</sub>) 3510 cm.<sup>-1</sup> and broad NH singlet at  $\delta$  7.68 in the 60 Mc./sec. n.m.r. spectrum\*

from proton exchangeable with  $D_2O$ ] unsubstituted in ring a (four-proton multiplet between 415—455 c./sec.), with only one exchangeable proton ( $M^+$  268 after  $D_2O$  exchange). Strong Bohlmann bands between 2500—2800 cm.<sup>-1</sup> in the i.r. spectrum are consistent with the structure assigned.

Chemical proof of the structure of elaeocarpidine depends upon the formation of 1-ethyl- $\beta$ -carboline on selenium dehydrogenation and on a study of dihydroelaeocarpidine (IIa). The labile nature of the N-CH-N system of (I) is shown by the formation on catalytic hydrogenation in glacial acetic acid solution of dihydroelaeocarpidine (IIa). Dihydroelaeocarpidine, m.p. 123—125,  $M^+$  269, has two exchangeable protons ( $M^+$  271 after D<sub>2</sub>O exchange) and in pyridine-acetic anhydride solution, affords an N-acetyl derivative (IIb) [v<sub>max</sub> (CHCl<sub>3</sub>) 1635 cm.<sup>-1</sup>; three-proton singlet at  $\delta 2.18$ ]. The structure of (IIb) was established by Hofmann degradation (aqueous NaOH at 180°) of the methiodide of (IIb), which afforded N-methylpyrrolidine as the only volatile, basic product.

(Received, February 14th, 1968; Com. 180.)

<sup>\*</sup> N.m.r. spectra were measured in CDCl<sub>3</sub> solution on a Varian A60 spectrometer and chemical shifts are relative to tetramethylsilane.