


## A Total Synthesis of Prostaglandin E<sub>1</sub> and Related Substances *via* *endo*-Bicyclohexane Intermediates


By UDO AXEN,\* FRANK H. LINCOLN, and JOHN L. THOMPSON

(Department of Experimental Chemistry, The Upjohn Company, Kalamazoo, Michigan 49001)

In our studies concerning a prostaglandin synthesis *via* solvolysis of bismethanesulphonates (1)<sup>1</sup> we prepared all four possible glycols (2) (two racemates with *erythro*-configuration and two racemates with *threo*-configuration of the *vic*-glycol) in pure form. These glycols have *exo*-configuration at C-13. In addition, we isolated four minor products from our reaction mixtures with very similar chromatographic




and spectral properties to those of the four major glycols. These minor by-products gave consistently 2–3 times higher yields of (±)-prostaglandin E<sub>1</sub> (PGE<sub>1</sub>) methyl ester when converted into bismethanesulphonates and solvolyzed under standard conditions. The origin and the properties of these by-products allowed only one conclusion about their structure: namely that these were the *endo*-isomers (at C-13) of glycols (2), formed in small amounts either during the carbene addition step<sup>2</sup> or by isomerization in later steps of the synthesis. To prove these structural assignments and to make use of the substantially higher yield in the solvolysis step, we devised a new synthesis which led stereospecifically to the *endo*-bicyclohexane glycols (3) and (4).



Bicyclo[3,1,0]hex-2-ene-6-*endo*-carboxylic acid (5), readily available from norbornadiene,<sup>3</sup> was converted into its methyl ester (6). Hydroboration followed by oxidation gave

alcohols (7) and (8),† which could be separated by column chromatography of their tetrahydropyranyl ethers (9) and (10).  $\text{LiAlH}_4$  reduction of (10) and oxidation with Jones' reagent yielded aldehyde (11), which was converted into ketone (12) by Wittig reaction followed by removal of the protecting group and oxidation with Jones' reagent. Only the *cis*-isomer of (12) was formed in the Wittig reaction ( $J$  11 Hz. for the olefinic protons in the n.m.r. spectrum). Alkylation of (12) with methyl  $\omega$ -iodoheptanoate resulted in



the formation of two alkylated ketones (13) and (14) isomeric at C-8, in a ratio of 1:4 in favour of the  $\alpha$ -isomer (14). This ratio represents approximately the thermodynamic equilibrium, as equilibration studies (potassium *t*-butoxide) with pure (13) and pure (14) have shown, and is in contrast to the *exo*-series<sup>1</sup> (isomeric at C-13), where the thermodynamic equilibrium favours the  $\beta$ -isomer. Hydroxylation of (14) with  $\text{OsO}_4$  yielded two glycols (3) and (4) in approximately equal amounts, which were identical with two of the four minor by-products isolated from the *exo*-series. Conversion of the less polar (silica gel chromatography) glycol (3) into the bismethanesulphonate followed by solvolysis in 2:1 acetone-water at room temperature gave 17–18% yield [based on glycol (3)] of ( $\pm$ )- $\text{PGE}_1$  methyl ester (15) [identical with ( $\pm$ )- $\text{PGE}_1$  methyl ester obtained earlier<sup>1</sup> in spectral properties, m.p. and mixed m.p.] and 19% of ( $\pm$ )-15-*epi*- $\text{PGE}_1$  methyl ester (16). The more polar glycol (4) gave 19% yield of ( $\pm$ )- $\text{PGE}_1$  methyl ester (15) and 16–17% yield of the 15-*epi*-isomer (16). Assuming that hydrolysis of the methanesulphonate group at C-15 proceeds predominantly with inversion at this centre, the less polar glycol (3) must have the same relative configuration at C-8 and C-15 as natural  $\text{PGE}_1$  and the more polar glycol (4) the same relative configuration at these centres as 15-*epi*- $\text{PGE}_1$ . The configuration at C-14 of (3) and (4) is determined by the *erythro*-relationship of the *vic*-glycol system.

The  $\beta$ -isomer (13) was converted into ( $\pm$ )-8-*iso*- $\text{PGE}_1$  methyl ester<sup>4</sup> and its 15-*epi*-isomer in a similar manner.

(Received, January 13th, 1969; Com. 048.)

† No attempt was made to determine the stereochemistry of the hydroxy-group relative to the three-membered ring, but it was assumed that the borane attack occurred from the less hindered side and no change in configuration took place during the peroxide oxidation.

<sup>1</sup> W. P. Schneider, U. Axen, F. H. Lincoln, J. E. Pike, and J. L. Thompson, *J. Amer. Chem. Soc.*, 1968, **90**, 5895.

<sup>2</sup> G. Just, C. Simonovitch, F. H. Lincoln, W. P. Schneider, U. Axen, G. B. Spero, and J. E. Pike, *J. Amer. Chem. Soc.*, (submitted).

<sup>3</sup> J. Meinwald, S. S. Labana, and M. S. Chadha, *J. Amer. Chem. Soc.*, 1963, **85**, 582.

<sup>4</sup> E. G. Daniels, W. C. Krueger, F. P. Kupiecki, J. E. Pike, and W. P. Schneider, *J. Amer. Chem. Soc.*, 1968, **90**, 5894.