

## Oxidation of Hydroxy-steroids

By ROBERT E. HARMON,\* CARMEN V. ZENAROSA, and S. K. GUPTA

(Department of Chemistry, Western Michigan University, Kalamazoo, Michigan 49001)

WE report the first example of the use of alkynylamines<sup>1</sup> for the oxidation of the hydroxy-group in steroids.<sup>2</sup> This new procedure involves the acid-catalysed reaction of *NN*-diethylaminoprop-1-yne with dimethyl sulphoxide and can be illustrated by the oxidation of testosterone and 11-hydroxyprogesterone.

The oxidations were conducted by stirring a mixture of the appropriate hydroxy-steroid (5 mmole), Me<sub>2</sub>SO (5 ml.), *NN*-diethylaminoprop-1-yne (13.5 mmole), and dry benzene (3 ml.). A solution of 100% phosphoric acid (1.59 g.) in Me<sub>2</sub>SC (1 ml.) was added to it dropwise and the reaction mixture was stirred at room temperature for 2 days. It

was then diluted with benzene (200 ml.), washed with 10% sodium hydrogen carbonate solution, and the organic phase was evaporated to dryness. The keto-steroid was isolated in 60% yield by chromatography of the residue over Silica Gel G. The physical constants of the keto-steroids thus obtained were in agreement with the literature values. Thus, the oxidation of testosterone and 11-hydroxyprogesterone furnished androst-4-ene-3,17-dione<sup>3</sup> and 11-oxoprogesterone,<sup>4</sup> respectively.

This work was supported by a Grant from the National Cancer Institute.

(Received, January 21st, 1969; Com. 088.)

<sup>1</sup> H. G. Viehe, *Angew. Chem. Internat. Edn.*, 1967, **6**, 767.

<sup>2</sup> K. E. Pfitzner and J. G. Moffatt, *J. Amer. Chem. Soc.*, 1965, **87**, 5670.

<sup>3</sup> P. N. Rao, *J. Org. Chem.*, 1961, **26**, 2149.

<sup>4</sup> T. Reichstein and H. G. Fuchs, *Helv. Chim. Acta*, 1940, **23**, 688.