




Cleavage of Alkyl Aryl Ethers with Lithium Iodide

By I. T. HARRISON

(Institute of Organic Chemistry, Syntex Research, 3401 Hillview Avenue, Palo Alto, California 94306)

Summary Alkyl aryl ethers are cleaved by lithium iodide to yield the corresponding phenol.

THE halogenolysis of esters by iodide ion is a reaction of considerable utility, allowing the equivalent of ester



hydrolysis in the presence of functions sensitive to either acid or base.¹ Examination of the mechanism (a) of this reaction indicates that a similar cleavage (b) of alkyl aryl ethers should be possible.

We find that these ethers can in fact be cleaved in this manner. Thus a solution of 2-methoxynaphthalene (1 g.) in dry 2,4,6-collidine (4 ml.) containing LiI (1.5 g. dried at 300° under nitrogen) was heated to reflux under nitrogen for 10 hr., acidified, and extracted giving an almost quantitative yield of 2-naphthol. Oestrone methyl ether (1; R = Me) gave oestrone (1, R = H) in 48 hr. Oestrone ethyl ether (1; R = Et) was also cleaved, although more slowly. The reaction mixture becomes mildly basic as the cleavage proceeds but may be buffered by the addition of an equivalent amount of an acid, *e.g.* benzoic acid. Alternatively, the reaction may be carried out with LiI·3H₂O at 180–200° in the absence of solvent. No cleavage was observed with 17-benzyloxyandrostane and hexadecyl methyl ether.

(Received, April 28th, 1969; Com. 569.)

¹ E. Taschner and B. Liberek, *Roczniki Chem.*, 1956, **30**, 323; F. Elsinger, J. Schreiber, and A. Eschenmoser, *Helv. Chim. Acta*, 1960, **43**, 113.