

## Selective Reductions of Steroid Carbonyl Groups with Homogeneous Catalysts Containing Iridium and Rhodium

By JAMES C. ORR,\* M. MERSEREAU, and A. SANFORD

(Huntington Laboratories of Harvard University at the Massachusetts General Hospital and Department of Biological Chemistry, Harvard Medical School, Boston, Massachusetts 02114)

**Summary** Steroid 2-, 3-, and 17-ketones can be reduced selectively with isopropyl alcohol, trimethyl phosphite, and sodium chloroiridate or tris(triphenylphosphine)-rhodium(I) chloride; stereospecific 2- and 3-reduction to the axial alcohol is best attained with the rhodium catalyst.

THE reduction of 5 $\alpha$ -cholestan-3-one to the axial 3 $\alpha$ -alcohol by sodium chloroiridate, aqueous isopropyl alcohol, and trimethyl phosphite has been described by Henbest *et al.*,<sup>1</sup> and its remarkable selectivity for reduction of the 3-keto-group alone under the original conditions has been employed for the preparation of a number of useful 3-axial steroid alcohols.<sup>2</sup>

We have also observed this selectivity and examined some different aspects of the reaction. In addition, a rhodium-containing catalyst of even greater selectivity has been discovered.

The reductions were conveniently effected by maintaining a mixture of the steroid (36  $\mu$ moles, approx. 10 mg.), isopropyl alcohol (0.82 ml.), trimethyl phosphite (0.10 ml., 854  $\mu$ moles), and a solution of sodium chloroiridate (9  $\mu$ moles) in water (0.09 ml.) in a sealed tube at 82° (isopropyl alcohol-water azeotrope) for 24 hr. Preparative runs (0.1–3 g.) were also carried out in sealed tubes with relatively less solvent mixture. The reaction products were examined by t.l.c., n.m.r., and combined g.l.c.–mass spectrometry. Under these conditions the 3-keto-groups of 5 $\alpha$ - and 5 $\beta$ -steroids were completely reduced, predominantly to the axial alcohols. 4-Keto-groups in steroids of both the A/B *cis* and A/B *trans* series and 17- and 20-keto-groups remained unaffected; 30% of 5 $\alpha$ -cholestan-2-one remained unreduced. After 46 hr., approximately 20% of 19-oxocholesterol was reduced to cholest-5-ene-3 $\beta$ ,19-diol. The reduction of 17 $\beta$ -hydroxy-5 $\alpha$ -androstan-2-one was complete after 9 days at 81°, giving 99% of the axial product androstane-2 $\beta$ ,17 $\beta$ -diol, and 1% of the equatorial 2 $\alpha$ ,17 $\beta$ -diol. The two epimers were readily separable by t.l.c.; the n.m.r. absorptions of the 19-H<sub>3</sub> protons and of the C-2 proton allowed clear distinction between them. Pure 3 $\alpha$ -hydroxy-5 $\alpha$ -androstan-17-one and 3 $\beta$ -hydroxy-5 $\beta$ -androstan-17-one were prepared directly from the corresponding diketones in 70–80% yields.

In agreement with Browne and Kirk,<sup>3</sup> we find that isomerization occurs at the 17-position during selective 3-reduction of 5 $\alpha$ -pregnane-3,20-dione. The n.m.r. absorption of the 18-H<sub>3</sub> protons with a 17 $\alpha$ -acetyl side-chain (0.907 p.p.m. from Me<sub>4</sub>Si) clearly distinguishes it from the 17 $\beta$ -epimer (0.605 p.p.m.). The relative areas under the peaks indicate approximately 14% isomerization, less than the equilibrium value,<sup>3</sup> suggesting that milder conditions might cause reduction of the 3-ketone with less isomerization at C-17. Pure 3 $\alpha$ -hydroxy-5 $\alpha$ ,17 $\beta$ -pregnan-20-one (m.p. 174–175°) could not be obtained from the product (m.p. 110–130°) even after eight recrystallizations from

several solvents (to m.p. 164–173°), but was obtained by column chromatography on alumina.

Addition to the reaction mixture described above of acetic acid (1.75 mmoles) caused a slight decrease in the rate of reduction of the 3-keto-group of 5 $\beta$ -androstan-3,17-dione. Sodium hydroxide (0.1 ml. of 1N), although insufficient to neutralize the phosphorous acid, caused approximately a 3-fold increase in rate and gave, after 10 hr. at 82°, complete reduction of the 3-keto-group to the alcohols with no apparent loss of stereospecificity (3 $\beta$ ; 3 $\alpha$  approximately 11:1). In addition, the 17-keto-group had been reduced to the extent of 5%. 3 $\beta$ -Hydroxy-5 $\alpha$ -androstan-17-one was treated at 100° for 7 days in the presence of sodium hydroxide. The product contained approximately 35% of unreduced 17-ketone, 60% of 5 $\alpha$ -androstan-3 $\beta$ ,17 $\beta$ -diol and 5% of the 3 $\beta$ ,17 $\alpha$ -diol. Separation of the diols was accomplished only by t.l.c. on alumina G (Brinkmann) with benzene:ethanol 9:1 and on titanium dioxide with benzene:ether 4:1. The isomers could be distinguished by the n.m.r. absorptions of 18-H<sub>3</sub> at 0.73 p.p.m. (17 $\beta$ -OH) and 0.64 p.p.m. (17 $\alpha$ -OH).

The predominant formation of the  $\psi$ -equatorial 17 $\beta$ -alcohol indicates that formation of the axial alcohol is not obligatory. Inspection of models suggests that in each case the reducing agent approaches from the less hindered side of the carbonyl group. The reduction of the 10-formyl group indicates that the mechanism need not be *via* the hydrogenation of an enol.

Testosterone, under the chloroiridate reducing conditions in the presence of sodium hydroxide for 24 hr., gave starting material (70%), androsta-3,5-diene-17 $\beta$ -ol (27%) and a small amount of an allylic alcohol. The allylic alcohol was not detectable by g.l.c.–mass spectrometry, but showed on t.l.c. as a blue spot that appeared immediately on spraying the plate with sulphuric acid-ethanol.

The possibility of employing the reaction in the oxidative direction was examined. 3 $\alpha$ -Hydroxy-5 $\alpha$ -androstan-17-one was treated with acetone, trimethyl phosphite, water (9:1:1 by volume), and sodium chloroiridate, and after 18 hr. at 100° in a sealed tube, gave 5% yield of the dione, the remainder being unchanged. These conditions of time and temperature were sufficient for complete reaction in the reductive direction.

During the first 30 min. of heating, the chloroiridate reaction mixture turns from deep red-brown to colourless<sup>4</sup> (under reflux in air,<sup>1</sup> the solution becomes yellow); on opening the tube, gas evolution occurs. The possibility that the transfer of hydrogen from isopropyl alcohol to the steroid might be taking place through reversible homogeneous hydrogenation led us to examine the homogeneous hydrogenation catalyst tris(triphenylphosphine)rhodium(I) chloride.<sup>5</sup> Its substitution for sodium chloroiridate resulted in complete reduction of 5 $\beta$ -androstan-3,17-dione to the 3-alcohols with even greater stereospecificity (axial: equatorial 50:1), and with no detectable reduction of the

17-keto-group. The rhodium catalyst caused slow (60% complete in 9 days at 82°) reduction of 17 $\beta$ -hydroxyandrost-2-one entirely to the 2 $\beta$ ,17 $\beta$ -diol; the iridium catalyst under these conditions also caused complete reduction but gave 1% of the equatorial 2 $\alpha$ ,17 $\beta$ -diol. The rhodium catalyst does not cause reduction of the 10-formyl group. The reaction rate at the 3-position with the rhodium catalyst is approximately the same as that of the chloroiridate reduction in the absence of sodium hydroxide. Without trimethyl phosphite, no reduction takes place; tris(triphenylphosphine)rhodium(I) chloride remains as deep red crystals. In its presence a yellow solid is formed and partly dissolved. The rate of the rhodium-catalysed

reaction is not appreciably altered by addition of sodium hydroxide. On a preparative scale, removal of the yellow rhodium catalyst from the steroid requires chromatography on alumina;<sup>6</sup> the iridium catalyst can be removed merely by water-ethyl acetate partition.

This work was supported by the U.S. Public Health Service and a grant from the American Cancer Society. The LKB combined g.l.c.-mass spectrometer was purchased through a special grant from the American Cancer Society, Massachusetts Division, Inc.

We are most grateful to Dr. Lewis L. Engel for many helpful discussions.

(Received, December 4th, 1969; Com. 1848.)

<sup>1</sup> Y. M. Y. Haddad, H. B. Henbest, J. Husbands, and T. R. B. Mitchell, *Proc. Chem. Soc.*, 1964, 361.

<sup>2</sup> P. A. Browne and D. N. Kirk, *J. Chem. Soc. (C)*, 1969, 1653.

<sup>3</sup> M. B. Rubin, *Steroids*, 1963, 2, 561.

<sup>4</sup> Catalytic hydrogenation of double bonds by such catalysts appears to be associated with the direct bonding of hydrogen to the metal atom in solution: J. Trocha-Grimshaw and H. B. Henbest, *Chem. Comm.*, 1968, 544, 757; J. R. Shapley, R. R. Schrock, and J. A. Osborn, *J. Amer. Chem. Soc.*, 1969, 91, 2816.

<sup>5</sup> M. A. Bennett and P. A. Longstaff, *Chem. and Ind.*, 1965, 846.

<sup>6</sup> C. Djerassi and J. Gutzwiller, *J. Amer. Chem. Soc.*, 1966, 88, 4537.