


Synthesis of Some Dimethylpyrano- and 3-Methylbut-2-enyl-4-phenyl- and -4-n-propyl-coumarins

By D. E. GAMES* and N. J. HASKINS

(Department of Chemistry, University College, P.O. Box 78, Cardiff CF1 1XL)

Summary A number of dimethylbenzodipyranones and 3-methylbut-2-enyl-4-n-propyl-5,7-dihydroxycoumarins, including mammeigin and mammea's B/AA and B/AB have been synthesised, showing that MAB5 and MAB6 are mixtures, the structure of ponnalide requires revision, and that a number of these compounds are present in the seed extract of *Mammea americana* L.


THE synthesis of a number of dimethylchromenylated naturally occurring compounds using 3-hydroxy-1,1-dimethoxy-3-methylbutane (I) has been reported by Crombie *et al.*¹ and the reaction has recently been extended to allow the introduction of two 2,2-dimethylpyran groups.² A number of 4-phenyl and 4-alkyl coumarins which occur in the Guttiferae family have this ring system,^{3,4,6-8} and we have found that use of this reagent provides an efficient route to these compounds, enabling a number of these compounds to be detected in natural extracts, and indicating that the structure of ponnalide⁴ requires revision. We have also found that reaction of 2-methylbut-3-en-2-ol and boron trifluoride etherate⁹ with 4-n-propyl-5,7-dihydroxycoumarin results in the introduction of one or more 3-methylbut-2-enyl groups into the coumarin nucleus. This method has been used to synthesise the mammeas B/AA (IIa) and B/AB (IIb), coumarins isolated from *Mammea americana* L.¹⁰ and *Mammea africana* G. Don.⁸



	R ¹	R ²	R ³		R ¹	R ²	R ³
(IIa)	Pr ⁿ	B	A	(IVc)	Pr ⁿ	D	H
(IIb)	Pr ⁿ	C	A	(IVd)	Pr ⁿ	E	H
(IIIa)	Pr ⁿ	H	C	(IVe)	Pr ⁿ	A	H
(IIIb)	Pr ⁿ	H	B	(Va)	Ph	H	C
(IIIc)	Pr ⁿ	H	A	(Vb)	Ph	H	B
(IVa)	Pr ⁿ	C	H	(VIa)	Ph	C	H
(IVb)	Pr ⁿ	B	H	(VIb)	Ph	B	H

Heating of 5,7-dihydroxy-4-phenylcoumarin with (I) in pyridine gave, after chromatography, the crystalline benzotripyranone (VIIa) (m.p. 177.5—178.5°) and the chromens (VIIIa) (11%) (m.p. 225.5—227°) and (XIa) (8%) (m.p. 260—263°). Orientation of (VIIIa) and (XIa) was supported by the shifts observed for 3- and 4-H of the chromen in the n.m.r. spectrum of the acetates of (VIIIa) and (XIa), which agreed with the observations of Merlini and his co-workers.⁵ A similar reaction with 5,7-dihydroxy-4-n-propylcoumarin gave the benzotripyranone (VIIb) and the chromens (VIIIb) and (XIb). Structure (XIa) has been assigned to ponnalide which was isolated from *Calophyllum inophyllum*.⁴ Our data for (XIa) and its acetate are not in agreement with that reported for ponnalide.†


Condensation of the acetal (I) with the acyl coumarins (IIIa,b), (IVa—d), (Va,b), and (VIa,b)⁶ gave the pyranocoumarins (XIIa,b) (IXa—d), (XIIIa,b), and (Xa,b) in yields varying from 30 to 80%. The coumarin (Xb) (m.p. 150—151°) was identical with mammeigin^{6,7} isolated from *Mammea americana* L. and comparisons with a sample of MAB5 and a synthetic sample of MAB6, coumarins recently isolated from *Mammea africana* G. Don,⁸ showed them to be mixtures consisting of (Xa) (m.p. 89.5—92°) and (Xb) for MAB5, and (IXa) (m.p. 97—98.5°) and (IXb) (m.p. 90.5—91.5°) for MAB6. The 2-methylbutyryl isomer was the major component in each case.



(VIIa) R = Ph, (b) R = Prⁿ

A re-examination of the seed extracts of *Mammea americana* L., using m.s., t.l.c., and n.m.r. has resulted in the identification of (Xa), (IXa), and (IXb) in these extracts together with a number of as yet unidentified homologous acylated 4-phenyl- and 4-alkyl-2,2-dimethylpyranocoumarins.

Treatment of 4-n-propyl-5,7-dihydroxycoumarin with an excess of 2-methylbut-3-en-2-ol and boron trifluoride etherate in dioxan solution at room temperature for 24 h,




	R ¹	R ²		R ¹	R ²
(VIIIa)	Ph	H	(XIa)	Ph	H
(VIIIb)	Pr ⁿ	H	(XIb)	Pr ⁿ	H
(IXa)	Pr ⁿ	C	(XIIa)	Pr ⁿ	C
(IXb)	Pr ⁿ	B	(XIIb)	Pr ⁿ	B
(IXc)	Pr ⁿ	D	(XIIIa)	Ph	C
(IXd)	Pr ⁿ	E	(XIIIb)	Ph	B
(Xa)	Ph	C			
(Xb)	Ph	B			

A = Me₂C:CH·CH₂, B = Me₂CH·CH₂·CO, C = MeCH₂·CH(Me)·CO, D = MeCH₂·CH₂·CO, E = Me₂CH·CO.

followed by work-up under non-acidic conditions, and chromatography, gave the coumarins (IIIc) (m.p. 180—182°)

† Professor Seshadri has informed us that he has come to a similar conclusion and the structure of ponnalide is being revised.

(3%) (IVe) (m.p. 166—168°) (9%) (XIV) (m.p. 131—133°) (0.3%) and the α -pyrone (XV) (m.p. 192°) (1%). The



orientation of (IIIc), (IVe) and (XIV) are assigned on the basis of the Gibbs reaction and the conversion of (IIIc) and (IVe) into known 3,3-dimethylchromans.¹¹ With concentrated hydrochloric acid, the pyrone (XV) gave the expected 3,3-dimethylchroman (XVI)¹² and the spectral data obtained for (XV) was consistent with that reported for structurally similar compounds.¹³

Reaction of the acyl coumarins (IVb) and (IVa)⁶ with 2-methylbut-3-en-2-ol under similar conditions gave mamma B/AA (IIa) (m.p. 122.5—123°) (2%) and mamma B/AB (IIIb) (m.p. 118—120°) (2%) which were identical with the natural materials.^{8,10†}

We thank Professor L. Crombie for helpful discussions and Dr. F. Scheinmann for samples of MAB5 and MAB6 and for exchange of information and manuscripts prior to publication.

(Received May 18th, 1971; Com. 806.)

† Seshadri has recently reported¹⁴ the synthesis of the 4-phenylcoumarin mammeisin by a similar method.

¹ W. M. Bandaranayake, L. Crombie, and D. A. Whiting, *Chem. Comm.*, 1969, 970; *J. Chem. Soc. (C)*, 1971, 811.

² W. J. G. Donnelly and P. V. R. Shannon, *Chem. Comm.*, 1971, 76.

³ J. Polonsky, *Bull. Soc. Chim. France*, 1957, 1079; G. H. Stout and K. L. Stevens, *J. Org. Chem.*, 1964, **29**, 3604; S. K. Nigam, C. R. Mitra, G. Kuensch, B. C. Das, and J. Polonsky, *Tetrahedron Letters*, 1967, 2633; K. Kawazu, H. Ohigashi, and T. Mitsui, *ibid.*, 1968, 2383; D. P. Chakraborty and D. Chatterji, *J. Org. Chem.*, 1969, **34**, 3784; G. D. Breck and G. H. Stout, *J. Org. Chem.*, 1969, **34**, 4203.

⁴ D. Adinarayana and T. R. Seshadri, *Bull. Nat. Inst. Sci. India*, 1965, **31**, 90.

⁵ A. Arnone, G. Cardillo, L. Merlini, and R. Mondelli, *Tetrahedron Letters*, 1967, **43**, 4201.

⁶ L. Crombie, D. E. Games, and A. McCormick, *J. Chem. Soc. (C)*, 1967, 2545, 2553.

⁷ R. A. Finnegan and W. H. Mueller, *J. Org. Chem.*, 1965, **30**, 2342.

⁸ I. Carpenter, E. J. McGarry, and F. Scheinmann, *Tetrahedron Letters*, 1970, 3983.

⁹ F. Bohlmann and K. M. Kleine, *Chem. Ber.*, 1966, **99**, 885; A. C. Jain, P. Lal, and T. R. Seshadri, *Tetrahedron*, 1970, **26**, 2631.

¹⁰ L. Crombie, D. E. Games, N. J. Haskins, and G. F. Reed, *Tetrahedron Letters*, 1970, 3979.

¹¹ L. Crombie, D. E. Games, N. J. Haskins, and G. F. Reed, unpublished work.

¹² G. A. Howard, J. R. A. Pollock, and A. R. Tatchell, *J. Chem. Soc.*, 1955, 174.

¹³ G. H. Stout, R. A. Alden, J. Kraut, and D. F. High, *J. Amer. Chem. Soc.*, 1962, **84**, 2653; E. Ritchie and W. C. Taylor, *Tetrahedron Letters*, 1964, 1431; R. Stevens, *Chem. Rev.*, 1967, **67**, 19.

¹⁴ K. R. Bala and T. R. Seshadri, *Phytochemistry*, 1971, **10**, 1131.