

# Journal of The Chemical Society, Chemical Communications

NUMBER 13/1972

5 JULY

## A New Synthesis of Unsaturated Aldehydes

By E. HUNT and B. LYTHGOE\*


(Department of Organic Chemistry, The University, Leeds LS2 9JT)

**Summary** Primary allylic bromides A-Br and 1,3-dithian react to give sulphonium bromides; these form ylides which undergo [2,3]-sigmatropic rearrangement to 2-substituted 1,3-dithians; hydrolysis then gives the  $\beta\gamma$ -unsaturated aldehydes A'-CHO. (A' = rearranged allylic system corresponding to A).

by prototropic isomerisation into an  $\alpha\beta$ -unsaturated aldehyde. We have, for example, obtained  $\gamma$ -cyclocitral<sup>6</sup> (5; R = Me) by the new method, and have demonstrated by spectroscopic methods its conversion into  $\beta$ -cyclocitral (6; R = Me).

By means of the Claisen rearrangement an allylic alcohol A-OH can be converted<sup>1</sup> into the  $\gamma\delta$ -unsaturated aldehyde A'-CH<sub>2</sub>CHO; a valuable feature is that in this way the group -CH<sub>2</sub>CHO can be introduced efficiently into a sterically congested environment.<sup>2</sup> We now describe a related reaction sequence which allows the similar introduction of the aldehyde group itself, and provides a convenient method of converting an allylic bromide A-Br into the  $\beta\gamma$ -unsaturated aldehyde A'-CHO.

The allylic bromide (1; R = H) when treated with 1,3-dithian at 20° gave compound (2; R=H), m.p. 123–124°, in high yield. Treatment of (2; R = H) at -78° in tetrahydrofuran with n-butyl-lithium gave (3; R = H), which was expected, by analogy,<sup>3</sup> to rearrange to the 2-substituted 1,3-dithian (4; R = H). On warming to 20° and normal work-up (4) was obtained in over 80% yield as a homogeneous oil and was characterised by accurate mass and other spectral data [ $\tau$  (CDCl<sub>3</sub>) 5.20 (2H, s) and 5.67 (1H, d,  $J$  9.5 Hz)]. We have similarly prepared the related 1,3-dithians (4; R = Me) and (7). Since good methods are available<sup>4</sup> for the hydrolysis of 1,3-dithians, a new route is thus opened to  $\beta\gamma$ -unsaturated aldehydes. Moreover, when the initial allylic bromide contains a hydrogen atom at the  $\gamma$ -carbon atom, the aldehyde first formed can be converted



Baldwin<sup>6</sup> has recently used a carbene rearrangement sequence in order to convert  $\gamma\gamma$ -dimethylallyl bromide into 2,2-dimethylbut-3-enoic acid (*i.e.* A-Br  $\rightarrow$  A'-CO<sub>2</sub>H), an overall result similar to that achieved by the present method. Baldwin's method should be applicable to second-

ary allylic bromides, whereas ours may be restricted to primary representatives. Our immediate products, the 2 substituted 1,3-dithians, are valuable not only as the precursors of aldehydes, but also because they are readily transformed<sup>4,7</sup> into ketones, hydroxy-ketones, and diketones. We thank the S.R.C. for a Studentship (to E.H.).

(Received, 6th April 1972; Com. 585.)

<sup>1</sup> A. W. Burgstahler and I. C. Nordin, *J. Amer. Chem. Soc.*, 1961, **83**, 198.

<sup>2</sup> E.g. R. F. Church and R. E. Ireland, *J. Org. Chem.*, 1963, **28**, 17.

<sup>3</sup> R. B. Bates and D. Feld, *Tetrahedron Letters*, 1968, 417; G. M. Blackburn, W. D. Ollis, J. D. Plackett, C. Smith, and I. O. Sutherland, *Chem. Comm.*, 1968, 186; J. E. Baldwin, R. E. Hackler, and D. P. Kelly, *ibid.*, p. 537.

<sup>4</sup> E. J. Corey and B. W. Erickson, *J. Org. Chem.*, 1971, **36**, 3553, and references therein.

<sup>5</sup> B. Willhalm, U. Steiner, and H. Schinz, *Helv. Chim. Acta*, 1958, **41**, 1359.

<sup>6</sup> J. E. Baldwin and J. A. Walker, *Chem. Comm.*, 1972, 354.

<sup>7</sup> E. J. Corey and D. Seebach, *Angew. Chem. Internat. Edn.*, 1965, **4**, 1075, 1077.