


Exocyclic Co-ordination of Phosponitriles: The Crystal and Molecular Structure of $N_4P_4(NMe_2)_8, W(CO)_4$

By H. P. CALHOUN, N. L. PADDOCK,* J. TROTTER, and J. N. WINGFIELD

(Department of Chemistry, University of British Columbia, Vancouver 8, B.C., Canada)

Summary In the compound $N_4P_4(NMe_2)_8, W(CO)_4$, the phosphonitrile acts as a bidentate σ -ligand, through one ring nitrogen atom and one exocyclic group.

PHOSPHONITRILE derivatives form complexes with transition metal ions, and, where the geometry has been established crystallographically,¹⁻³ co-ordination occurs through the ring nitrogen atoms. They also react with metal carbonyls,⁴ but hitherto no structures have been determined. The structure of the title compound is unusual in that the metal is co-ordinated to both a ring nitrogen atom and an exocyclic dimethylamido-group. The yellow crystals are monoclinic, $a = 18.274$, $b = 18.594$, $c = 10.533$ Å, $\beta = 90.39^\circ$, $Z = 4$, space group $P2_1/n$. The structure was determined with Mo- K_α diffractometer data from Patterson and electron density maps, and was refined by full-matrix least squares methods to $R = 0.071$ for 2545 observed reflections. Some of the more immediately important bond lengths [$\sigma(P-N) = 0.02$ Å] are shown in the Figure. The four atoms N(1)WN(8)P(2) are nearly coplanar, the co-ordination direction at N(1) being reasonably near the conventional lone pair direction, as it is when co-ordinated to protons^{5,6} and transition metal ions.¹⁻³ In the unco-ordinated ligand,⁷ substantial delocalisation of the lone



pairs on the dimethylamido-groups into the ring shortens the exocyclic P-N bonds from the single bond length⁸ of 1.77 Å to 1.68 Å, the mean sum of the angles round the nitrogen atoms being 353°. In the complex, the tungsten atom is a competitive acceptor; P(2)N(8) is increased to 1.75 Å, and the co-ordination of N(8) is tetrahedral, dona-

tion from this atom to phosphorus being minimal. The compound appears to be a simple *cis*- σ -complex, an interpretation supported by a comparison of its carbonyl stretching frequencies (2000, 1870, 1849, 1809 cm^{-1}) with those⁹ of $\text{enW}(\text{CO})_4$ (2006, 1867, 1852, 1809 cm^{-1}). Coordination has further effects on the geometry of the ligand. The ring is formed from two nearly planar segments N(4)P(1)N(1)P(2) and P(2)N(2)P(3)N(3)P(4)N(4), making

a dihedral angle of 53.7° , the former set being nearly coplanar with N(1)WN(8)P(2).

Three bonds, P(2)N(2), P(3)N(3), and N(4)P(1) are significantly shorter than the other five; a first order perturbation treatment accounts for these differences satisfactorily.¹⁰ We thank the National Research Council of Canada for financial support.

(Received, 5th June 1972; Com. 963.)

¹ W. C. Marsh, N. L. Paddock, C. J. Stewart, and J. Trotter, *Chem. Comm.*, 1970, 1190.

² W. Harrison, N. L. Paddock, J. Trotter, and J. N. Wingfield, *J.C.S., Chem. Comm.*, 1972, 23.

³ J. Trotter and S. H. Whitlow, *J. Chem. Soc. (A)*, 1970, 455.

⁴ N. L. Paddock, T. N. Ranganathan, and J. N. Wingfield, *J.C.S., Dalton I* in the press.

⁵ N. V. Mani and A. J. Wagner, *Chem. Comm.*, 1968, 658.

⁶ J. Trotter and S. H. Whitlow, *J. Chem. Soc. (A)*, 1970, 460.

⁷ G. J. Bullen, *J. Chem. Soc.*, 1962, 3193.

⁸ E. Hobbs, D. E. C. Corbridge, and B. Raistrick, *Acta Cryst.*, 1953, 6, 621; D. W. J. Cruickshank, *Acta Cryst.*, 1964, 17, 671.

⁹ C. S. Kraihanzel and F. A. Cotton, *Inorg. Chem.*, 1963, 2, 533.

¹⁰ W. C. Marsh, T. N. Ranganathan, J. Trotter, and N. L. Paddock, *Chem. Comm.*, 1970, 815.