Precocenes

Juvenile hormones are necessary throughout most stages of the insect life cycle for development of ovaries, sex pheromone production and larval diapause. Application of

exogenous juvenile hormones to insect populations can upset development only during the brief period of metamorphosis that occurs when an immature insect molts to the adult stage.^{2,3}

Bowers and co-workers⁴ have found that introduction of some anti-juvenile hormones (antiallatotropins) into an insect colony terminates juvenile hormone secretion and induces:

- 1) precocious molting of insects to sterile adults,
- inability of larvae which are dependent on a high titer of juvenile hormones to diapause,
- 3) sterility in adult insects which are dependent on juvenile hormone for ovarian development, and
- artificial diapause in species which diapause through a lack of juvenile hormones.

Bowers isolated 7-methoxy-2,2-dimethylchromene and 6,7-dimethoxy-2,2-dimethylchromene from extracts of the plant *Ageratum houstonianum*. Since these compounds were found to induce precocious metamorphosis in insects, he named them **Precocene I** and **Precocene II**, respectively.⁴

In tests performed by Bowers⁴ with the **Precocenes**, milkweed bugs underwent precocious metamorphosis, and in several other species of insects, sterilization was induced. **Precocene II** was also found to induce diapause in Colorado potato beetles.


In these tests, the biological effect of the **Precocenes** is equivalent to that of the surgical removal of the corpora allata which produce juvenile hormones in insects. Indeed, **Precocene II** was found to inhibit the biosynthesis of juvenile hormone by the cockroach corpora allata.⁵ Thus, the **Precocenes** depress juvenile hormone titer and can be highly effective insecticides. Moreover, Bowers,⁴ in some instances, succeeded in reversing the antiallatotropic activity in precocious adults by treatment with exogenous juvenile hormones

We believe that this remarkable discovery of antiallatotropic activity could guide the way to safe, economical and insect-specific pesticides.⁶

References:

- 1) G.M. Chippendale and C.M. Yin, Nature, 246, 511 (1973).
- 2) C.M. Williams, ibid., 178, 212 (1956).
- 3) W.S. Bowers, Bull. W.H.O., 44, 381 (1971).
- W.S. Bowers, T. Ohta, J.S. Cleere, and P.A. Marsella, Science, 193, 542 (1976); Chem. Eng. News, 54 (16), 19 (1976).
- 5) G.E. Pratt and W.S. Bowers, Nature, 265, 548 (1977).
- W.S. Bowers in "Yearbook of Agriculture," Department of Agriculture, Washington, D.C., 1968, pp 94-98.

Pentafluorobenzylhydroxylamine



PFBHA · HCI

O-(2, 3, 4, 5, 6-Pentafluorobenzyl)hydroxylamine hydrochloride (PFBHA · HCl) has recently been introduced as a very sensitive derivatizing agent for the gas chromatographic analysis of ketosteroids using electron capture detection. 1.2 For this type of detection technique to be effective, the compound to be analyzed must contain a strong electron-withdrawing group. Since most ketosteroids do not contain a sufficiently strong electron-withdrawing group, a suitable derivative must be prepared.

Clark and Wotiz³ first suggested the use of heptafluorobutyrate ester derivatives for the chromatography of hydroxysteroids such as testosterone. Although these derivatives are sensitive to electron capture detection, they suffer from the difficulty encountered in controlling the reaction to prepare selectively the 17-hydroxy mono ester, the 3-enol mono ester or the 3-enol-17-hydroxy diester. As a result, one steroid may be resolved into several peaks in the chromatogram, leading to confusion and error in the analysis. Koshy and co-workers have shown **PFBHA·HCl** to be a more effective derivatizing reagent. **PFBHA·HCl** converts ketosteroids to their oximes, and glc conditions can be chosen

such that both the syn and anti oximes appear as one peak.1

Steroid O-(2,3,4,5,6-pentafluorobenzyl) oximes are readily prepared under mild conditions by treating the ketosteroid with a solution of **PFBHA·HCl** in pyridine at 60 to 65°C. After evaporation of the solvent, the oxime is extracted into cyclohexane, and the cyclohexane solution is washed with water and dried over Na₂SO₄. Derivatives can be prepared from less than one nanogram of steroid. PFB-oxime derivatives of testosterone have been detected with samples as small as 5 picograms.

Nambara and co-workers have used **PFBHA·HCl** for the determination of dehydroepiandrosterone in human plasma with satisfactory results.² It has been suggested that this analytical technique may also be applicable to the determination of specific prostaglandins in tissues, and the analysis of ketosteroid levels in meat carcasses to ascertain if they fall within federal standards.⁴

References:

- 1) K.T. Koshy, D.G. Kaiser, and A.L. VanDerSlik, J. Chromatogr. Sci., 13, 97 (1975).
- T. Nambara, K. Kigasawa, T. Iwata, and M. Ibuki, J. Chromatogr., 114, 81 (1975).
- 3) S.J. Clark and H.H. Wotiz, Steroids, 2, 535 (1963).
- 4) Chem. Eng. News, 54 (41), 20 (1976).

19,448-4 *O-*(2,3,4,5,6-Pentafluorobenzyl)- 250mg \$17.00 hydroxylamine hydrochloride 1g \$45.00


Aldrich Chemical Company, Inc.

Craftsmen in Chemistry

ALDRICH

940 West St. Paul Ave Milwaukee Wisconsin 53233 USA Telephone (414) 273-3850

CS Publication/ New/



SPECIAL PUBLICATION No. 29

Herbicides and Fungicides FACTORS AFFECTING THEIR ACTIVITY

Edited by N. R. McFARLANE, Shell Research Ltd.

A wider understanding of the factors affecting the biological activity of herbicides and fungicides provides complex challenges. This volume provides a historical and current perspective of the general areas, including future problems and opportunities, mode of action of some well-known fungicides and their use in practice, criteria affecting efficiency of penetration and translocation of some herbicides. It then goes on to deal with important current experimentation including the influence of adjuvants and humidity, the action of 2,4-dichlorophenoxyacetic acid, inorganic glasses for the slow release of fungicides and herbicides, and spirolactone fungicides related to benzoquinones. The final papers cover theoretical aspects including partition coefficients and systemic activity and the concept of optimal lipophilicity.


Paperbound 156pp 8pp halftones 8½"×5½" £7.50 (CS Members £5.50)

Chemical Education in Europe

Edited by P. J. FARAGO, M. J. FRAZER, and S. D. WALKER

The science of chemistry has always been international and recent developments have added a new reality to this concept. Chemical education in Europe and the United Kingdom can now be regarded as a coherent whole, despite local differences and the exchange of people, both at staff and student levels, is proceeding apace. This book fulfils two needs: it describes the flavour of chemical education in a number of major European countries, based upon the experiences of well-known educators, at both secondary and tertiary levels. In addition it also provides a reference volume, collecting together a mass of information hitherto only available in widely scattered publications, which has been updated through personal interviews and questionnaires. It provides invaluable background and reference material both for those involved in the study and comparison of European education systems, and for those who are thinking of spending some time at an institution in a foreign country.

Paperbound 380pp 84"×53" £3.50 (CS Members £2.75)





MOLECULAR STRUCTURE:

ITS STUDY BY CRYSTAL DIFFRACTION

J. Clare Speakman



MONOGRAPHS FOR TEACHERS No. 30

Molecular Structure:

Its Study by Crystal Diffraction

By J. CLARE SPEAKMAN

Most chemists have become conditioned into believing that the molecule is 'really there'; they develop an image of it, with given size, shape and motions, and with certain relations with its neighbours. Though such simple faith is no doubt open to metaphysical attack, it works well. It receives apparent support from the wealth of information on molecular properties now accessible to physical methods of study. Those methods based on the diffraction, by crystals, of X-rays, and some other radiations, are particularly powerful. Their impact on the chemist may be likened to the experience of actually meeting a person, with whom communication had hitherto been restricted to the telephone. This monograph offers an elementary explanation of crystal-diffraction by X-rays and neutrons, and goes on to illustrate the results obtainable on a variety of molecules.

Paperbound 57pp 8½"×5½" £1.20 (CS Members 90p)

Further information on any title may be obtained from: The Marketing Officer, The Chemical Society, Burlington House, London W1V 0BN.

Orders should be sent to The Chemical Society, Distribution Centre, Blackhorse Road, Letchworth, Herts SG6 1HN

or personal callers may purchase copies from The Library, The Chemical Society, Burlington House, London W1V 0BN.

Organometallic Compounds

With the award of the Nobel Prize to Victor Grignard in 1912 for his pioneering work on alkyl-magnesium halides and for the development of the Grignard synthesis, a phase in the development of organometallic compounds was completed. However, improved experimental techniques soon led to new successes in this field, it became possible to synthesize organometallic compounds with nearly all metals of the main and subgroups of the periodic system. Many of these compounds have industrial applications, e.g. the organometallic mixed catalysts of the Ziegler-Natta type for the polymerization of olefins (for which K. Ziegler and G. Natta received the Nobel Prize in 1963), or organotin compounds used effectively as fungicides and bacteriocides. In "classical" organometallic compounds the metal atom is bound to the carbon atom by a more-or-less strongly polarized σ -bond, while in other groups of carbonmetal compounds, a π -bond is present between the metal and carbon atoms. Well-known examples of the latter are ferrocene and dibenzenechromium.

Due to the continuously increasing number of publications on organometallic compounds it is nearly impossible to maintain a general overview of the subject. In this field, the Gmelin Handbook renders a valuable service with its volumes on organometallic compounds published in the last few years as part of the New Supplement Series. There are several volumes for the transition metals (Cr, Co, Fe, Ni, Pt, Ag, Ta, Ti, V, Zr). The source material, which is widely dispersed in numerous publications, is critically evaluated and clearly presented with the aim of achieving an exhaustive review of the pertinent literature. The cut-off date for the literature coverage is indicated in each volume. In the New Supplement Series organometallic compounds are classified according to their formulae following the number of organicallybound metal atoms in the molecule

(mononuclear compounds, binuclear compounds, etc.) rather than in the standard Gmelin Handbook method. Further classification is made according to the organic ligands, whereby the number of metal-carbon bonds which can form a ligand is the determining factor. If a compound contains different ligands, the Gmelin Principle of the Last Position is applied, i.e. the compound is arranged according to the ligands which form the most metal-carbon bonds.

Bibliographic data, chemical preparation, reactions, physical properties, analysis, utilization, spectroscopy of organometallic compounds are treated in individual chapters with the usual Gmelin thoroughness. There is a formula index for each element. The preface, table of contents and the headings of individual sections appear in both English and German, greatly facilitating the Handbook's use for the non-German reader. Volume 10/11 of the New Supplement Series — Zr and Hf organic compounds, respectively, is entirely in English. Below is a listing of the organometallic volumes of the Gmelin Handbook which have been published to date. These volumes provide the investigator with a key to the latest developments in the science of organometallic chemistry.

ORGANOVANADIUM
COMPOUNDS
ORGANOCHROMIUM
COMPOUNDS
Single Volume
(New Suppl. Series Vol. 2/3):
1971. DM 622,—; US \$273.70

ORGANOCOBALT COMPOUNDS (New Suppl. Series Vol. 5): 1973. DM 696,—; US \$306.30

(New Suppl. Series Vol. 6): 1973. DM 329,—; US \$144.80

1973. DM 169,—; US \$74.40

ORGANOZIRCONIUM COMPOUNDS ORGANOHAFNIUM COMPOUNDS Single Volume (New Suppl. Series Vol. 10/11): ORGANOIRON COMPOUNDS (New Suppl. Series Vol. 14): 1974. DM 516,—; US \$227.10 (New Suppl. Series Vol. 36): 1976. DM 431,—; US \$189.70 (New Suppl. Series Vol. 41): 1977. DM 642,—; US \$282.50

ORGANONICKEL COMPOUNDS (New Suppl. Series Vol. 16): 1975. DM 694,—; US \$305.40 (New Suppl. Series Vol. 17): 1974. DM 618,—; US \$272.00 (New Suppl. Series Vol. 18): 1975. DM 198,—; US \$87.20 Index for Volume 16 + 17

ORGANOTIN COMPOUNDS (New Suppl. Series Vol. 26): 1975. DM 301,—; US \$132.50 (New Suppl. Series Vol. 29): 1975. DM 788,—; US \$346.80 (New Suppl. Series Vol. 30): 1976. DM 327,—; US \$143.90 (New Suppl. Series Vol. 35): 1976. DM 311,—; US \$136.90

ORGANOTITANIUM COMPOUNDS (New Suppl. Series Vol. 40): 1977. DM 481,—; US \$211.70

ORGANOSILVER COMPOUNDS System No. 61 Part B, Section 5 1975. DM 322,—; US \$141.70

Prices are subject to change without notice

Shorter chapters on the organometallic compounds of NIOBIUM, PLATI-NUM, RUTHENIUM, and TANTA-LUM are to be found in the volumes: Niobium System No. 49, Part B, Section 4, Platinum System No. 68, Part D, Ruthenium System No. 63, Ru-Supplement Volume and Tantalum System No. 50, Part B, Section 2.

The GMELIN HANDBOOK of Inorganic Chemistry will be distributed by Springer-Verlag Berlin, Heidelberg, New York. All volumes can be separately ordered from your bookseller.

On request we are eagerly willing to send you the alphabetical reference chart for Gmelin as well as the complete catalogue.

CLASSIFIED ADVERTISEMENTS

HERIOT-WATT UNIVERSITY

Department of Chemistry

POST-DOCTORAL APPOINTMENT

Applications are invited for a post-doctoral appointment, supported by the S.R.C, to collaborate with Professor J. G. Buchanan and Dr R. H. Wightman on aspects of the biosynthesis of C-nucleoside antibiotics.

Applicants should have a sound background in either microbiology or synthetic organic chemistry and a strong interest in biological chemistry. Experience in carbohydrate chemistry or organic radiochemistry would be an advantage.

The appointment is available from 1st September, 1977, and is for one year. The salary will be £3333—£3547 p.a. on the 1A scale, depending on experience, plus U.S.S. benefits. Applications, with full curriculum vitae, details of research experience, and the names of two referees should be sent as soon as possible to: Professor J. G. Buchanan, Department of Chemistry. Heriot-Watt University, Riccarton, Edinburgh EH14 4AS.

UNIVERSITY OF SALFORD

DEPARTMENT OF CHEMISTRY AND APPLIED CHEMISTRY

POST DOCTORAL RESEARCH FELLOWSHIP IN ORGANIC CHEMISTRY

Applications are invited from persons with previous experience in synthetic and/or intermediate chemistry, for a Post Doctoral Fellowship to work with Dr E. F. V. Scriven on a project sponsored by the Science Research Council to study some synthetic and biomimetic reactions or aromatic azides. The appointment, from 1 January 1978 is for 1 year with the possibility of renewal.

Salary on the scale: £3333 — £3975 per annum USS/FSSU. Further particulars and application forms may be obtained from the Registrar, University of Salford, Slaford M5 4WT. to whom applications should be returned by 16 September 1977 quoting reference CH/207/CC.

POSTDOCTORAL POSITION IN DEPARTMENT OF BIOLOGICAL CHEMISTRY

Position open October 1, 1977. To work on total synthesis of natural products; good knowledge and experimental background in synthetic organic chemistry required. Experience in carbohydrate chemistry desirable but not necessary. Send resume and two letters of recommendation to Dr Momcilo Miljkovic, Department of Biological Chemistry, The Milton S. Hershey Medical Center, The Pennsylvania State University, Hershey, PA 17033 U.S.A. Equal Opportunity Employer.

CLASSIFIED ADVERTISING

DISPLAY AND SEMI DISPLAY £1.50 per single col. cm. column width 43 mm (10 ems)

Send your advertisements to:-

Susan Shaw, Classified Advertisements, Burlington House, Piccadilly, London W1V 0BN.

MONOGRAPHS FOR TEACHERS

Elements of Organometallic Chemistry

by F. R. HARTLEY

The two major stumbling blocks to the introduction of organometallic chemistry in school teaching are the lack of suitably simple text and the lack of suitable experiments. It is the aim of this monograph to remedy this situation. It contains chapters covering basic organometallic chemistry in terms of preparation, structures, and bonding in organometallic compounds. It then examines the systematic chemistry of the compounds with particular reference to their industrial applications. The final section contains details of experiments which have been specially modified to enable them to be performed in schools without the use of sophisticated equipment.

Paperbound 103pp 8%" x 5%" ISBN 0 85186 909 2 Price £1,20 (Chemical Society Members 90p)

Obtainable from your local bookseller or: THE CHEMICAL SOCIETY
Blackhorse Road, Letchworth, Herts SG6 1HN.