

Synthesis of Allenes *via* Reduction of Acetylenic Tosylhydrazones

By GEORGE W. KABALKA,* RAY J. NEWTON, Jr., and JAMES H. CHANDLER
(*Department of Chemistry, University of Tennessee, Knoxville, Tennessee 37916*)

and DOMINIC T. C. YANG
(*Department of Chemistry, University of Arkansas at Little Rock, Little Rock, Arkansas 72204*)

Summary The reduction of tosylhydrazones of conjugated acetylenic ketones occurs with migration of one of the acetylenic bonds, providing a mild synthesis of allenes.




THE reduction of tosylhydrazones by catecholborane (CB)¹ and other reagents² provides a mild alternative to the Wolf-Kishner and Clemmensen reductions [equation (1)].

TABLE. Formation of allenes *via* reduction of tosylhydrazones of conjugated acetylenic ketones.

Ketone ^a	Product	Yield ^b /%	ν^c/cm^{-1}	δ^d
Oct-3-yn-2-one	Octa-2,3-diene	78 (64)	1950	1.02 (m,3H), 1.35—2.1 (m,9H), 4.80 (m,1H), 5.50 (m,1H)
But-1-yn-3-one	Buta-1,2-diene	49	1960	1.50 (m,3H), 4.45 (m,2H), 4.80 (m,1H)
4-Phenylbut-3-yn-2-one	1-Phenylbuta-1,2-diene	84 (75)	1960	1.60 (dd,3H), 5.5 (m,1H), 6.12 (m,1H), 7.3 (m,5H)
1-Phenylbut-2-yn-1-one	1-Phenylbuta-1,2-diene	30	1960	1.60 (dd,3H), 5.5 (m,1H), 6.12 (m,1H), 7.3 (m,5H)

^a The ketones were first converted into the tosylhydrazone derivatives. ^b By n.m.r. analysis; yields in parentheses are those of isolated products. ^c Allene stretch region. ^d Solvent was CDCl_3 .

The reduction of tosylhydrazones of α,β -unsaturated carbonyl compounds is accompanied by the regiospecific migration of the double bond³ [equation (2)].



We have found that a regiospecific migration also occurs during the reduction of tosylhydrazones derived from conjugated acetylenic ketones to produce allenes in good yield [equation (3)].

The selectivity of catecholborane⁴ and the mild reaction conditions (room temp.) make this synthesis a valuable addition to the existing routes to allenes.⁵ The results are summarized in the Table.

The reduction of the tosylhydrazone of oct-3-yn-2-one is representative. The tosylhydrazone, m.p. 66–68 °C, was prepared in ethanol by mixing equimolar amounts of tosylhydrazide and the ketone; after 1 h at reflux, the product crystallized on cooling. The tosylhydrazone (5 mmol) was dissolved in CHCl_3 (5 ml) and CB (5 mmol) was added. The reduction was allowed to proceed at room temperature until complete (monitored by n.m.r. spectroscopy). Then, $\text{NaOAc}\cdot 3\text{H}_2\text{O}$ (15 mmol) was added and the mixture was warmed for 1 h. The product (78% yield) was isolated by chromatography on silica gel.

(Received, 10th May 1978; Com. 502.)

i, H_2NNHTs ; ii, CB; iii, $\text{NaOAc}\cdot 3\text{H}_2\text{O}$. Ts = *p*- $\text{MeC}_6\text{H}_4\text{SO}_2$.

¹ G. W. Kabalka and J. D. Baker, *J. Org. Chem.*, 1975, **40**, 1834.

² R. O. Hutchins, C. A. Milewski, and B. E. Maryanoff, *J. Amer. Chem. Soc.*, 1973, **95**, 3662; L. Caglioti, *Tetrahedron*, 1966, **22**, 487.

³ G. W. Kabalka, D. T. C. Yang, and J. D. Baker, Jr., *J. Org. Chem.*, 1976, **41**, 574; D. T. C. Yang and G. W. Kabalka, *Org. Prep. Proc. Int.*, 1977, **9**, 85.

⁴ G. W. Kabalka, J. D. Baker, Jr., and G. W. Neal, *J. Org. Chem.*, 1977, **42**, 512.

⁵ P. Rona and P. Crabbé, *J. Amer. Chem. Soc.*, 1969, **91**, 3289; M. M. Midland, *J. Org. Chem.*, 1977, **42**, 2650.