

## Biomimetic Synthesis of Borreverine and Isoborreverine

By FRANÇOIS TILLEQUIN and MICHEL KOCH\*

*(Laboratoire de Matière Médicale, Faculté de Pharmacie, 4 avenue de l'Observatoire 75006 Paris, France)*

and JEAN-LOUIS POUSSET and ANDRÉ CAVÉ

*(U.E.R. de Chimie Thérapeutique, Centre d'Études, Pharmaceutiques, 92290 Chatenay Malabry, France)*


*Summary* Acidic dimerisation of the isoprene-indole alkaloid borrerine (I) gives borreverine (II) and isoborreverine (III).

THE co-occurrence of the indole alkaloid borrerine (I)<sup>1</sup> and of its dimer borreverine (II) in *Borreria verticillata* suggested a simple chemical relationship between these two compounds and the mechanism shown in the Scheme can be

envisaged for the dimerisation of (I): protonation of borrerine (I) gives two open forms (a) and (b), which, after intermolecular cycloaddition, lead to borreverine (II) and isoborreverine (III) via two *cis-trans* isomeric intermediates (c) and (d).

$C_6H_6$ , 65 °C, 30 min) borreverine (II)† and isoborreverine (III)‡ were isolated in approximately equal amounts and in ca. 80% total yield.

Longer reaction time leads to a change in the ratio of borreverine (II) to isoborreverine (III) in favour of the latter.



SCHEME

In order to verify the above mechanism borrerine (I) was heated with several acids (HCl,  $MeCO_2H$ ,  $CF_3CO_2H$ ) under various conditions. Both dimers were obtained in each case although the rate of the reaction and the yield depended on the acid used. Under optimum conditions ( $CF_3CO_2H$ ,

Furthermore, borreverine (II) can be quantitatively transformed into isoborreverine (III) after 12 h under the above conditions. Protonation of the tertiary nitrogen of borreverine (II) may, in fact, lead to a reversal of the reaction sequence depicted in the Scheme whereas isoborre-

† Identical (u.v., i.r.,  $^1H$  n.m.r., and mass spectroscopy, and t.l.c.) with an authentic sample (ref. 2).

‡ The structure of isoborreverine (III) has been demonstrated by physical and spectral data and by chemical correlation:  $C_{32}H_{40}N$  ( $M^+$  480.3244, calc. 480.3253);  $[\alpha]_{D}^{20}$  0°;  $\lambda_{max}$  (EtOH) (log  $\epsilon$ ) 226 (4.72), 288 (4.15), and 294 (4.15) nm; 3400, 2960, 2925, 2870, 2800, 1460, 1300, 1135, and 745  $cm^{-1}$ ;  $m/e$  480 ( $M^+$ ) (68%), 438 (21), 437 (61), 436 (15), 405 (37), 394 (100), 393 (52), 264 (16), 262 (38), 197 (56), 195 (81), 185 (40), 182 (55), 144 (46), and 130 (18);  $^1H$  n.m.r.  $\delta$  ( $Me_4Si$ ) 0.76 (3H, s), 1.06 (3H, s), 1.69 (3H, s), 2.42 (3H, s), 2.44 (3H, s), 5.40 (2H, m), 6.40–7.80 (8ArH), and 2.35–2.65 and 4.05 (2H+1H,  $D_2O$  exchangeable). Acetylation of (III) in cold pyridine leads to a product identical (u.v., i.r.,  $^1H$  n.m.r. and mass spectroscopy, and t.l.c.) with transposed diacetylborreverine, the structure of which has been shown by X-ray diffraction (ref. 2).

rine (III) should not react under these conditions. A similar rearrangement was observed when borreverine (II) was treated with hot acetic anhydride.<sup>2</sup>

Since the dimerisation of borrerine (I) invariably furnishes borreverine (II) as well as isoborreverine (III), their co-occurrence within the same plant might be expected. In fact, these two dimeric alkaloids were recently isolated from *B. verticillata*<sup>3</sup> and *Flindersia fourniei*.<sup>4</sup>

Our experimental data and the lack of optical activity in both natural and synthetic dimers suggest a non-enzymatic

dimerisation of borrerine (I). The isolation of isoborreverine (III) as the major constituent of *Flindersia fourniei*<sup>4</sup> is in full agreement with this hypothesis. Nevertheless, it is interesting to note that borreverine (II) is the main alkaloid in *B. verticillata*.<sup>3</sup>

We thank Pr. Jean Levy (Faculté de Pharmacie, 51 rue Cognacq-Jay 51095 Reims Cedex, France) for helpful discussions about the borrerine dimerisation mechanism.

(Received, 1st June 1978; Com. 562.)

<sup>1</sup> J. L. Pousset, J. Kerharo, G. Maynard, X. Monseur, A. Cavé, and R. Goutarel, *Phytochemistry*, 1973, **12**, 2308.

<sup>2</sup> J. L. Pousset, A. Cavé, A. Chiaroni, and C. Riche, *J.C.S. Chem. Comm.*, 1977, 261.

<sup>3</sup> J. L. Pousset, unpublished results.

<sup>4</sup> F. Tillequin, M. Bert, T. Sevenet, and M. Koch, *Lloydia*, in the press.