

Electrophilic Cleavage of Unsaturated Thiirans


By PETER HEGGISON MCCABE* and ANGUS STEWART

(Department of Chemistry, University of Glasgow, Glasgow G12 8QQ)

Summary High-yield reactions of unsaturated thiirans with halogens or SO_2Cl_2 proceed by initial electrophilic attack on sulphur and furnish halogeno-sulphides, in some cases *via* isolable disulphides, thereby providing a novel method of creating an intramolecular disulphide link.

SULPHUR-BRIDGED systems are conveniently prepared by the widely applicable condensation¹ of SOCl_2 with olefins but the halogenolysis^{2,3} of unsaturated thiirans as a potential route to dihalogeno-sulphides has not been fully explored. In the latter case, it is not known whether halogen, as an electrophile, first attacks the double bond or the episulphide sulphur atom.


Chlorinolysis of (1) and (2) furnishes^{2,4} respectively, (3) and (6). Similarly, we have shown that (1) reacts³ at room temperature with Br_2 or I_2 forming (4) (97%) or (5) (67%) and that (2) undergoes brominolysis to give (7)⁵ (98%).



To monitor the intermediates of these halogenolyses, the reactions of (1) with 1.1 mol of Br_2 in CDCl_3 or I_2 in CD_2Cl_2 were studied in the n.m.r. probe. At -30°C , direct conversion into (4) or (5) occurred and no intermediates were detected even when the halogens were added in portions.

In contrast, reaction of (8)⁶ at -40°C in the n.m.r. probe with 1.1 mol of Br_2 in CDCl_3 gave (10), m.p. $60-62^\circ\text{C}$ [$\delta(\text{CDCl}_3)$ 3.63—3.83 (*H*-CS) and 4.30—4.72 (*H*-CBr); *m/e* 270, 272, and 274], *via* a clearly detectable unsaturated dibromide intermediate, which formed exclusively with 0.5 mol of Br_2 and could be isolated at room temperature. Treatment of (8) with 0.5 mol of I_2 in CFCl_3 at 0°C furnished an unsaturated di-iodide, closely similar (spectra) to the above dibromide, but this remained unchanged on addition of an excess of iodine. These dihalides, $\text{C}_{12}\text{H}_{16}\text{Br}_2\text{S}_2$, *m/e* 382, 384, and 386 and $\text{C}_{12}\text{H}_{16}\text{I}_2\text{S}_2$, *m/e* 477-87815 (M^+), reacted with bromine to yield (10) and were assigned structures (13), b.p. $125-130^\circ\text{C}$, 0.15 mmHg [$\delta(\text{CDCl}_3)$ 4.37—4.77 (*H*-CBr) and 5.47—5.73 (vinyl H); $\nu(\text{CCl}_4)$ 1661 cm^{-1}] and (14), b.p. $90-100^\circ\text{C}$, 0.04 mmHg [$\delta(\text{CDCl}_3)$ 4.50—4.95 (*H*-Cl); $\nu(\text{CCl}_4)$ 1658 cm^{-1}]. Furthermore, (13) condensed with sodium sulphide in tetrahydrofuran to


give a trisulphide, tentatively formulated as (16)¹¹ [$\delta(\text{CDCl}_3)$ 3.1—3.8 (*H*-CS) and 5.7 (vinyl H); *m/e* 256 (M^+), 190 ($M-\text{H}_2\text{S}_2$), and 158 ($M-\text{H}_2\text{S}_3$)]. In preparative reactions, (13) and (14) were formed from (8) in 95 and 90% yield (of isolated material), respectively, and the conversions of (13) into (10) and (8) into (10) were quantitative.



Reaction of (8) with an excess of Cl_2-CCl_4 gave a 2:5 (by n.m.r. spectroscopy) mixture of (9)⁷ and (11), m.p. $110-115^\circ\text{C}$ [$\delta(\text{CDCl}_3)$ 3.59 (*H*-CSO), 4.04—4.50 (*H*-CCl), and 4.73—5.24 (*H*-CCl); *m/e* 198, 200, and 202], the latter being isolated (21%) by t.l.c., while chlorinolysis of (8) with SO_2Cl_2 (1 mol) in CH_2Cl_2 at 0°C furnished exclusively (9) (94%). With 0.5 mol of SO_2Cl_2 , (8) formed the disulphide (12) (97%), b.p. $95-100^\circ\text{C}$, 0.03 mmHg (CDCl_3) 4.17—4.60 (*H*-CCl); $\nu(\text{CCl}_4)$ 1663 cm^{-1}].

Thus, halogenolysis of (8) proceeds by initial electrophilic ring opening of the thiiran, forming the sulphenyl halide (15), which condenses with unchanged substrate to yield a disulphide [(12), (13), or (14)], the latter reaction being faster. When an excess of Cl_2 or Br_2 is present, the disulphide is cleaved reforming (15), in which intramolecular addition of sulphenyl halide to the double bond occurs in the absence of substrate (8) [reaction of which with (15) must be faster] to form the bicyclic sulphides (9) and (10). Iodinolysis⁸ of the disulphide (14) does not occur and (11) is produced from (9) by chlorine oxidation in presence of air.

By analogy, electrophilic ring opening of the thiiran group of (1) also takes place first but the greater conformational flexibility of the eight-membered ring causes intramolecular condensation of sulphenyl halide with the double bond to be faster than disulphide formation.



Reaction of (17)⁸ with 0.5 mol of SO_2Cl_2 furnished only starting material and (18), while, with 1.1 mol of SO_2Cl_2 , (18)⁹ was formed in 83% yield. Compound (18) underwent partial isomerisation and hydrolysis to (19)⁹ and (21), m.p. 110–112 °C [δ (CDCl_3) 3.71 ($H\text{-CS}$), 4.01–4.47 ($\text{CH}_2\text{-O}$), and 4.54 (d, J 2 Hz, $H\text{-CCl}$), m/e 254, 256, and 258] when the mother liquor of crystallisation of (18) was chromatographed. Bromine treatment of (17) gave a 2:1 mixture of (22) and (20) (combined yield 94%) from which the

former, m.p. 106–108 °C [δ (CDCl_3) 4.75 (d, J 2 Hz, $H\text{-CBr}$), m/e 342, 344, and 346], was isolated (35%) by crystallisation. Distillation (120–130 °C at 0.3 mmHg) of the mother liquor gave no further separation, the structural assignment of (20) being based on its n.m.r. similarity to (18).

The intermediacy of disulphides in the halogenolysis of (8) led us to investigate the possibility of creating an intramolecular disulphide bridge by an appropriate ring opening process. Reaction of (8) in CH_2Cl_2 (10^{-3} M) with SCl_2 at -50 °C gave a dichloro-disulphide [m/e 214, 216, 218 (M^+), 145, 147 ($M\text{-SCl}$), and 113, 115 ($M\text{-S}_2\text{Cl}$), λ_{max} (MeOH) 361 (ϵ 66) and 270 nm (33)], which possessed four distinct resonances (31.9, 34.7, 53.7, and 56.6 ppm) in the noise-decoupled ^{13}C n.m.r. spectrum and showed complex ^1H n.m.r. absorption. These spectroscopic properties are compatible with structure (23)¹⁰ and the formation of this compound represents a new method of preparing a cyclic disulphide (42% yield).

The above results demonstrate that reaction of unsaturated thirans with electrophiles proceeds by initial ring opening of the episulphide group, providing a high-yield route to sulphides and, in certain cases, disulphides.

Satisfactory microanalytical data have been obtained for (10), (12), (13), (21), (22), and (23) and mass measurement for (14). Compounds (12)–(14) and (16) are depicted in only one diastereomeric form.¹¹

We thank the S.R.C. for the award of a studentship to A.S.

(Received, 8th October 1979, Com 1065)

¹ M. Muhlstadt and D. Martinetz, *Z. Chem.*, 1974, **14**, 297.

² F. Lautenschlaeger, *J. Org. Chem.*, 1969, **34**, 3998.

³ P. H. McCabe, C. M. Livingston, and A. Stewart, *J. C. S. Chem. Comm.*, 1977, 661.

⁴ F. Lautenschlaeger, *J. Org. Chem.*, 1968, **33**, 2620.

⁵ S. D. Ziman and B. M. Trost, *J. Org. Chem.*, 1973, **38**, 649.

⁶ F. Lautenschlaeger and N. V. Schwartz, *J. Org. Chem.*, 1969, **34**, 3991.

⁷ E. J. Corey and E. Block, *J. Org. Chem.*, 1966, **31**, 1663.

⁸ For comparison see J. P. Danehy in 'Sulphur in Organic and Inorganic Chemistry,' ed. A. Senning, Marcel Dekker, New York, 1971, Vol. 1, p. 331 and L. Field in 'Organic Chemistry of Sulphur,' ed. S. Oae, Plenum Press, New York, 1977, p. 362.

⁹ P. H. McCabe and W. Routledge, *Tetrahedron Letters*, 1976, 85.

¹⁰ Cf. *syn*-2,3-dithiabicyclo[3.2.1]octan-8-ol. R. M. Wilson, D. N. Buchanan and J. E. Davis, *Tetrahedron Letters*, 1971, 3919.

¹¹ For comparison see G. A. Tolstikov, N. N. Novitskaya, R. G. Kant'yukova, L. V. Spirikhin, N. S. Zefirov, and V. A. Palyulin, *Tetrahedron*, 1978, **34**, 2655.